

## Growth hormone administration and exercise combination increase serum IGF-1 levels in perimenopausal wistar rats

Ni Luh Kadek Alit Arsani<sup>1</sup>, Luh Putu Ratna Sundari\*<sup>2</sup>

<sup>1</sup>Faculty of Medicine, Universitas Pendidikan Ganesha, Bali, Indonesia

<sup>2</sup>Department of Physiology, Faculty of Medicine, Universitas Udayana, Bali, Indonesia

Original Article

### ABSTRACT

#### ARTICLE INFO

##### Keywords:

Growth hormone,  
exercise,  
swimming,  
IGF-1,  
perimenopausal Wistar rats

##### \*Corresponding author:

luhputu\_ratna@unud.ac.id

DOI: 10.20885/JKKI.Vol14.Iss1.art9

##### History:

Received: November 1, 2021

Accepted: March 14, 2023

Online: April 5, 2023

Copyright ©2023 Authors. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International Licence (<http://creativecommons.org/licenses/by-nc/4.0/>).

**Background:** Growth hormone (GH), Insulin-like growth factor 1 (IGF-1), and exercise play a crucial role in female reproductive function. At the perimenopausal age, there is a decrease in GH, estrogen, and testosterone levels. In previous studies, the impact of physical activity on serum IGF-1 levels in geriatric was inconsistent and needed further investigation.

**Objective:** The purpose of this study was to look into the effect of GH and exercise on serum IGF-1 levels in perimenopausal Wistar rats.

**Methods:** It is an experimental study with a randomized posttest-only control group design using 24 perimenopausal Wistar rats aged 14 months as the sample. The control rats (C) were given 0.1 ml of distilled water subcutaneously every day for 30 days. Treatment group 1 (T1) rats swam for 30 minutes at a time five times per week. Meanwhile, rats of T2 were given 0.016 IU/0.1ml GH daily injection subcutaneously only for 30 days and T3 were given combination both of GH injection in same dose and swam for 30 minutes at a time 5 times a week for 30 days Data analysis used One-Way ANOVA.

**Results:** The finding revealed that the mean serum IGF-1 levels in the T3 (588.50 ± 84.04 ng/mL) were significantly higher than the control (242.03 ± 46.08 ng/mL), T1 (334.23 ± 75.90 ng/mL) and T2 (428.69 ± 95.10 ng/mL).

**Conclusion:** Our study shows that GH administration and exercise combination increases levels of IGF-1 serum higher than only administrated GH or exercise.

**Latar Belakang:** Hormon pertumbuhan (GH), insulin-like growth factor 1 (IGF-1) dan olahraga memiliki peran penting dalam fungsi reproduksi wanita. Pada tahap sebelum terjadi keluhan penuaan, terjadi penurunan kadar GH, dan hormon-hormon seks seperti estrogen dan testosteron. Pada riset terdahulu, pengaruh olahraga pada kadar IGF-1 serum pada lansia menunjukkan hasil bervariasi yang memerlukan riset lebih lanjut.

**Tujuan:** Penelitian ini bertujuan untuk mengetahui pengaruh kombinasi GH dan latihan terhadap kadar IGF-1 serum tikus wistar perimenopause.

**Metode:** Studi ini merupakan uji eksperimental dengan rancangan randomized posttest only control group menggunakan 24 ekor tikus wistar perimenopause umur 14 bulan sebagai sampel. Tikus kontrol (C) disuntik secara subkutan dengan akuabides 0,1 ml/hari selama 30 hari. Tikus kelompok intervensi 1 (T1) diberi perlakuan berenang secara bersamaan selama 30 menit, 5 kali/minggu selama 30 hari. Sementara itu, tikus T2 disuntik dengan GH 0,016 IU/0,1 ml subkutan saja, sedangkan tikus T3 diberi kombinasi injeksi GH dengan dosis yang sama dan aktivitas berenang 30 menit, 5 kali/minggu selama 30 hari. Data dianalisis dengan uji One-Way ANOVA

**Hasil:** Hasil penelitian menunjukkan bahwa rerata kadar IGF-1 serum pada kelompok T3 (588,50 ± 84,04 ng/mL) secara bermakna lebih tinggi dibandingkan dengan kelompok kontrol (242,03 ± 46,08 ng/mL), T1 (334,23 ± 75,90 ng/mL) dan T2 (428,69 ± 95,10 ng/mL).

**Kesimpulan:** Dari studi ini dapat disimpulkan kombinasi injeksi GH dan aktivitas berenang meningkatkan level IGF-1 serum lebih tinggi daripada pemberian GH saja maupun latihan saja.

## INTRODUCTION

At the perimenopausal age, there is a decrease in GH, estrogen, and testosterone levels. GH production is cut in half every seven years from the age of eighteen to twenty-five. Therefore, an increase in age leads to a reduction in the amplitude of GH impulse. At the transition stage of the aging process, age thirty-five to forty-five years, hormone levels decrease by 25%. Meanwhile, at a later stage of the age of forty-five years and above or clinical stage, the hormone levels steadily decrease, which includes Dehydroepiandrosterone (DHEA), melatonin, growth hormone, testosterone, estrogen, and thyroid hormone. Every three years, there is a decrease in bone density which leads to a decline in muscle mass by one kilogram resulting in the inability to burn calories. These cause an increase in body fat and weight, which eventually trigger chronic illness.<sup>1</sup>

Growth hormone, a vital modulator of female reproduction, works both directly through its receptors and indirectly through the stimulation of insulin-like growth factor-1 (IGF-1).<sup>2</sup> Previous studies reported that GH is a survival factor for primary ovarian follicles that regulates granulosa cell differentiation and inhibits follicular apoptosis and atresia.<sup>3</sup> Furthermore, GH is reported to modulate follicle-stimulating hormone's (FSH) work on granulosa cells by increasing local synthesis of IGF-1. This IGF-1 increases the working effect of gonadotropin at the level of granulosa cells and theca cells' IGF-1 synergism with FSH. Luteinizing hormone (LH) stimulates estrogen and progesterone production by granulosa cells, thereby functioning as a follicular survival factor.<sup>4</sup> In addition to using chemical drugs, physical exercise has been regarded and supported by the community as the crucial and inseparable factor that contributes to maintaining good physical and psychological health. However, the problem is whether the exercise is correct and sufficient to maintain the body's health.

The goal of exercise in health maintenance is to do enough physical activity that is beneficial to health but not too much that causes stress on the body and mind.<sup>5</sup> Physical activity is a powerful stimulator of GH secretion. However, the exact mechanism of the types of exercise that improve GH secretion still needs to be clarified further.<sup>6</sup> Hypothalamus hormones and neurotransmitters regulate the GH secretion, especially the circulating

IGF-1 as the primary mediator downstream for GH activity. It is a strong reason why studies explore the connection between IGF-1 and GH with physical exercise.<sup>6</sup>

Several studies on the advantages of physical exercise on plasma IGF-1 levels showed inconsistent results. A study reports an increase in IGF-1 levels due to physical exercise but another showed no change in IGF-1 levels.<sup>7-10</sup> The effect of exercise on IGF-1 production is affected by supplementation of energy as well as the intensity, type, and duration. However, this required further investigation.<sup>11</sup>

The population of women approaching the age of menopause and its health problems posed a challenge for the world of health. If the rapid decline in follicle numbers during pre-menopause does not occur, the proposed mathematical model estimates that the ovaries will remain functional until the age of seventy-one years.<sup>12</sup> Therefore, studies need to be carried out on factors that maintain or improve ovarian function before the ovarian follicle expires at menopause. Judging from some advantages of physical exercise on GH and IGF-1 levels and GH and IGF-1 as important modulators for female reproduction, studies will investigate the GH administration and physical in increasing serum IGF-1 levels. In addition, a combination of GH administration and physical exercise is proposed to have more benefits than GH or exercise alone.

## METHODS

### Research design

It was an experimental study with a randomized post-test-only control group design using 24 perimenopausal Wistar rats aged 14 months as a sample. This study was in the Biomedical Integrated Laboratory of the Faculty of Medicine, Udayana University. Furthermore, it has been reviewed by the Health Research Ethics Commission of the Faculty of Medicine, Udayana University/RSUP Sanglah No. 285 / UN.14.2/KEP/2016

### Subject

Fourteen months-old rats with similar calculations with women entering the menopause period were chosen. The inclusion criteria are body weight between 180–200 grams and healthy condition. Using Federer Formula for sample

calculation, minimally 24 rats with six rats in each group. The sample amount is added up to 8 rats per group to prevent sample shortages due to dropout.

### Laboratory treatment

Before the study, researchers observed the number of times rats swam. It turned out that rats could swim for 40 minutes until exhausted. In this research, rats were given swimming exercises for 30 minutes ( $3/4 \times 40 \text{ minutes} = 30 \text{ minutes}$ ) with frequency, intensity, time, and type (FITT) principles. Growth Hormone (GH) in this research used Genotropin® and contains recombinant somatotropin 16 IU per 1 ml.

The dose adjustment to rat from human dose was  $0.018 \times 0.9 \text{ IU/day} = 0.016 \text{ IU/day}$ . Growth hormone daily at 1.6 IU/0.1 ml, was diluted with 9.9 ml aquabidest to get a 0.016 IU/0.1 ml dose. The control rats (C) were given 0.1 ml of distilled water subcutaneously every day for 30 days. Treatment group 1 (T1) rats swam for 30 minutes at a time five times per week. Meanwhile, rats of T2 were given 0.016 IU/0.1ml GH daily injection subcutaneously only for 30 days and T3 were given combination both of GH injection in same dose and swam for 30 minutes at a time 5

times a week for 30 days. After intervention for 30 days, blood was drawn in the periorbita area of rats about 1 ml per day. Then, serum IGF-1 concentration in ng/mL was tested with the ELISA technique (Enzym-Linked Immunosorbent Assay) in double replication using Rat IGF-1 Elisa Kit from Elabscience.

### Data analysis

The data were explored for normality using the Shapiro-Wilk test, homogeneity by the Levene test, and comparison test was analyzed by the One-Way ANOVA test. We also checked for post hoc analysis by LSD (least significant difference) on significance ( $p < 0.05$ ).

## RESULTS

### Serum IGF-1 levels in various treatments and control

The results showed normal distribution and homogeneity. The results of the significance analysis with the One Way Anova test are presented in Table 1 below. Table 1 showed that the average serum IGF-1 levels in the four groups after treatment were differed significantly ( $p < 0.05$ ). The Least Significant Difference-test (LSD) results are shown in Table 2.

Table 1. Mean difference of serum IGF-1 levels between groups

Treatment	n	IGF-1 level (ng/mL)	SD	p-value
Control (C)	6	242.03	46.08	0.000
Exercise (T1)	6	334.23	75.90	
GH (T2)	6	428.69	95.10	
GH and exercise (T3)	6	588.50	84.04	

p significance:  $p < 0.05$ , SD: Standard Deviation

Table 2. Comparison of serum IGF-1 levels after treatment between groups

Treatment	Mean difference (ng/mL)	p-value
C and T1	92.20	0.052
C and T2	186.67	0.000*
C and T3	364.48	0.000*
T1 and T2	94.46	0.047*
T1 and T3	254.27	0.000*
T2 and T3	159.81	0.002*

p\* significance:  $p < 0.05$ , C: Aquabidest only, T1: Exercise Only, T2: GH only, T3: GH and exercise

Table 2 showed that the comparison of C and T1 was not significantly different, but the other group was. It showed that group exercise only did not get a significant result when compared with group GH only or combination exercise and GH. It means higher serum IGF-1 levels in combination with GH and exercise reinforces the theory that exercise stimulates GH production physiologically and enhances GH action through increased circulating IGF-1 secretion.<sup>15</sup>

## DISCUSSION

The results showed that the GH group rats' serum IGF-1 levels were significantly higher than the control group (Table 1). It is in line with Kraemer et al. that the administration of GH may increase IGF-1 serum levels of old female mice to reach the level of IGF-1 at a young age.<sup>16</sup>

Various tissue-specific stimulatory factors can increase the production of IGF-1 in a specific tissue. As described, gonadotropin and sex hormone stimulate the production of IGF-1 in reproduction organs, such as the testis, ovarium, and uterus in humans.<sup>17</sup>

Growth hormone works physiologically both directly and indirectly through its receptors, stimulating the formation of IGF-1, a vital mediator of GH activity, produced in the liver and peripheral tissues. Analysis and interpretation of IGF-1 levels is a beneficial and essential diagnostic tool for follow-up in growth hormone deficiency (GHD) and GH-related disorders. Growth hormone administration generally aims to address clinical complaints with appropriate GH doses and within safe limits.<sup>2</sup>

The results of this study showed that the average serum IGF-1 levels in the exercise group were greater than in the control group, but the difference was not statistically significant (Table 2). However, results from previous studies on the effects of physical exercise on IGF-1 levels are not consistent on whether physical exercise increases IGF-1 levels. Serum IGF-1 levels elevated during resistive and endurance exercise.<sup>18</sup> Moreover, a report shows that physical exercise increases serum IGF-1 levels in transgenic mice with initially low serum IGF-1 levels.<sup>19</sup> Similarly, the study by Cetinkaya et al. in mice given aerobic exercise, also reported an increase in serum IGF-1 levels.<sup>20</sup> This is consistent with the study in older rats given continuous resistance training that reported an

increase in IGF-1 levels.<sup>21</sup>

Insulin-like Growth Factor-1 secretion due to physical exercise appears to be more correlated to the duration of the given physical exercise and the session of a single activity.<sup>22</sup> Acute physical exercise causes a temporary increase in serum IGF1 levels for the first 10 minutes before returning to baseline. Stein et al. also stated that the results from the different studies on the effects of physical exercise on IGF-1 levels seem to depend on the type and intensity of physical exercise.<sup>23</sup> Furthermore, it has been reported that continuous mild physical exercise raises serum IGF-1 levels while strength training lowers them.<sup>11,24</sup>

The physical exercise provided in this study is high-intensity training sub maximal type according to the FITT principle, such as 75% from maximum swim ability. This study used maximum swim duration from the mice based on preliminary research (40 minutes). It resulted in no significant increase in serum IGF-1 levels between the two groups (C and T1) due to the relatively short duration of physical exercise (30 days). Several studies found that IGF-1 levels are related to exercise length and intensity.<sup>23,25</sup>

Aside from physical activity, serum IGF-1 levels are influenced by nutrition, age, chronic disease, insulin, and growth hormone.<sup>26</sup> Physical exercise affects IGF-1 levels through mechanisms that are still incompletely understood.<sup>15</sup> It became a limitation of this study because there are numerous variables to investigate in order to determine the effect of physical exercise on IGF-1 secretions.

This study showed that the mean serum IGF-1 levels in the group of rats with a combination of GH injection and swimming (T3) were significantly higher compared to the mean serum IGF-1 levels of the control group (C), T1, and T2.

Physical activity is also a powerful physiological stimulant for GH secretion.<sup>16</sup> After 15 minutes, aerobic exercise stimulates GH secretion, with the peak level occurring after or near the end of the exercise. Furthermore, the GH response to exercise is affected by the intensity and duration of aerobic training, fitness level, sex, and age.<sup>27,28</sup> Following the study about the effect of aerobic exercise may increase GH levels in a group of middle-aged women and geriatrics. Exercise stimulates GH secretion less in older people than in younger people. Meanwhile, GH secretion in postmenopausal women is 5.7-7.3 times lower



than in premenopausal women.<sup>28</sup>

The neuroendocrine mechanism in the induction of GH by exercise is still obscure. It is because this naturally involves a complex mechanism, probably due to the input on the hypothalamus. This mechanism could involve the release of Growth hormone-releasing hormone (GHRH) and/or somatostatin withdrawal, as well as the natural release of a Growth hormone releasing peptide (GHRP)-like ligand (such as ghrelin), or some combination of these mechanisms.<sup>16</sup>

In addition, Exercise-induced GH secretion appeared to involve multiple neuroregulatory responses in the GH axis and varied in peripheral feedback signals. A number chemical messenger were played a role, but the typical pathway appeared to involve stimulation of GHRH secretion and/or inhibition of somatostatin release. Furthermore, sympathetic activity may be a crucial mediator of GH response to acute exercise via activation of central  $\alpha$ 2-adrenergic neurons. One study has reported that increased levels of adrenaline and noradrenaline plasma concentrations lead to an elevated GH concentration due to exercise.<sup>27</sup>

## CONCLUSION

This study concluded that serum IGF-1 levels are increased by GH administration and the combination of GH and exercise. Therefore, it will overcome the problems caused by a decline in GH and IGF-1 levels in old age.

Furthermore, higher serum IGF-1 levels obtained from the combination of GH administration and exercise compared to the levels from the use of GH and exercise alone reinforced the theory that exercise was a physiologic stimulus of GH secretion and increased its activity by increasing the secretion of IGF-1 in circulation.

## CONFLICT OF INTEREST

All authors declare no conflict of interest.

## ACKNOWLEDGEMENT

The authors express gratitude to Ketut Tunas, Department of Public Health, Faculty of Health Sciences and Technology, Dhyana Pura University, Bali, Indonesia. Furthermore, thanks to I Gede Wiranata, Pharmacology Department, Faculty of Medicine Universitas Udayana, Indonesia for sharing their pearls of wisdom during the course of this study.

## REFERENCES

1. Jones CM, Boelaert K. The Endocrinology of Ageing: A Mini-Review. *Gerontology*. 2015;61:291–300.
2. Martirosian A, Silverstein J. Diagnosis and management of growth hormone deficiency in adults. *Consultant*. 2022;62(6):e20–7.
3. Dalcik H, Dalcik C. Insulin-Like Growth Factor I (IGF-I); cytoprotective effects, idea of replacement therapy in the healthy elderly subjects. *Medical Research Archives*. 2020;8(6).
4. Li J, Chen Q, Wang J, Huang G, Ye H. Does growth hormone supplementation improve oocyte competence and IVF outcomes in patients with poor embryonic development? A randomized controlled trial. *BMC Pregnancy Childbirth*. 2020;20:310.
5. Stults-Kolehmainen MA, Sinha R. The effects of stress on physical activity and exercise. *Sports Medicine*. 2014;44(1):81–121.
6. Birzniece V. Exercise and the growth hormone–insulin-like growth factor axis. *Current Opinion Endocrine Metabolic Research*. 2019;9(5):1–7.
7. de Alcantara Borba D, da Silva Alves E, Rosa JPP, Facundo LA, Costa CMA, Silva AC, et al. Can IGF-1 Serum Levels Really be Changed by Acute Physical Exercise? A Systematic Review and Meta-Analysis. *Journal of Physical Activity and Health*. 2020; 17: 575–84.
8. Chia-Liang T, Chun-Hao W, Chien-Yu P, Fu-Chen C. The effects of long-term resistance exercise on the relationship between neurocognitive performance and GH, IGF-1, and homocysteine levels in the elderly. *Frontiers in Behavioural Neuroscience*. 2015;9(23):1–12.
9. Devin JL, Bolam KA, Jenkins DG, Skinner TL. The influence of exercise on the insulin-like growth factor axis in oncology: physiological basis, current and future perspectives. *Cancer Epidemiology, Biomarkers & Prevention*. 2016;25(2):239–49.
10. Jiang Q, Lou K, Hou L, Lu Y, Sun L, Tan SC, et al. The effect of resistance training on serum insulin-like growth factor 1(IGF-1): A systematic review and meta-analysis. *Complementary Therapies in Medicine*. 2020;50:102360.
11. Kim T, Chang JS, Kim H, Lee KH, Kong ID. Intense Walking Exercise Affects Serum IGF-1

- and IGFBP3. *Journal of Lifestyle Medicine*. 2015;5(1):21–5.
12. Wu J, Liu Y, Song Y, Wang L, Ai J, Li K. Aging conundrum: A perspective for ovarian aging. *Frontiers in Endocrinology (Lausanne)*. 2022;13:1664–2392.
  13. Hofstetter J, Suckow MA, Hickman D. *The Laboratory Rat Morphophysiology*. Second edi. Suckow MA, Weisbroth SH, Franklin CL, editors. London: Elsevier Inc.; 2006.
  14. Adhikari S, Tretriluxana J, Dev R, Eglitis E, Nistha S, Kerfeld CP. FITT-CORRECT: Updated dynamic and evidence-based principle of exercise prescription. *Journal of Novel Physiotherapy and Rehabilitation*. 2021;5:005–9.
  15. Sundari LPR, Arsani NLKA. Regular Physical Exercise Increase of Growth Hormone (GH) and Insulin-Like Growth Factor-1 (IGF-1) Activity in Elderly Improve the Aging Process and Quality of Life: A Mini Review. *Biomedical and Pharmacology Journal*. 2022;15(2).
  16. Kraemer WJ, Ratamess NA, Hymer WC, Nindl BC, Fragala MS. Growth Hormone(s), Testosterone, Insulin-Like Growth Factors, and Cortisol: Roles and Integration for Cellular Development and Growth With Exercise. *Frontiers in Endocrinology (Lausanne)*. 2020;11(33).
  17. Ipsa E, Cruzat VF, Kagize JN, Yovich JL, Keane KN. Growth Hormone and Insulin-Like Growth Factor Action in Reproductive Tissues. *Frontiers in Endocrinology*. 2019;10(777).
  18. Giovanni V, Giuseppe P, Maria V, Leo JH. ROLE of IGF-1 System in the Modulation of Longevity: Controversies and New Insights From a Centenarians' Perspective. *Frontiers in Endocrinology (Lausanne)*. 2019;10.
  19. Al-Samerria S, Radovick S. The Role of Insulin-like Growth Factor-1 (IGF-1) in the Control of Neuroendocrine Regulation of Growth. *Cells*. 2021;10(10):2664.
  20. Cetinkaya C, Sisman AR, Kiray M, Camsari UM, Gencoglu C, Baykara B, et al. Positive effects of aerobic exercise on learning and memory functioning, which correlate with hippocampal IGF-1 increase in adolescent rats. *Neuroscience Letters*. 2013;549:177–81.
  21. Cenyi W, Jiling L, Yuanyuan R, Jielun H, Baoming J, Guodong W, et al. A Preclinical Systematic Review of the Effects of Chronic Exercise on Autophagy-Related Proteins in Aging Skeletal Muscle. *Frontiers in Physiology*. 2022;13.
  22. Naderi A, Shaabani F, Esmaeili A, Salman Z, Borella E, Degens H. Effects of low and moderate acute resistance exercise on executive function in community-living older adults. *Sport, Exercise, and Performance Psychology*. 2019;8(1):106–22.
  23. Stein AM, Silva TMV, Coelho FGM. Physical exercise, IGF-1 and cognition A systematic review of experimental studies in the elderly. *Dementia e Neuropsychologia*. 2018;12(2):114–22.
  24. Arnarson A, Gudny-Geirsdottir O, Ramel A, Jonsson PV, Thorsdottir I. Insulin-Like Growth Factor-1 and Resistance Exercise in Community Dwelling Old Adults. *Journal of Nutrition Health and Aging*. 2015;19(8):856–60.
  25. Norling AM, Gerstenecker AT, Buford TW, Khan B, Oparil S, Lazar RM. The role of exercise in the reversal of IGF-1 deficiencies in microvascular rarefaction and hypertension. *GeroScience*. 2020;42(1):141–58.
  26. Gulick CN, Peddie MC, Jowett T, Hackney AC, Rehrer NJ. Exercise, Dietary Protein, and Combined Effect on IGF-1. *International Journal of Science and Research Methodology*. 2020;16(3):61–77.
  27. Danese E, Tarperi C, Luca SG, Guzzo A, Sanchis-Gomar F, Festa L, et al. Sympatho-adrenergic activation by endurance exercise: Effect on metanephrines spillover and its role in predicting athlete's performance. *OncoTarget*. 2018;9:15650–7.
  28. Gusatovic J, Gramkow MH, Hasselbalch SG, Frederiksen KS. Effects of aerobic exercise on event-related potentials related to cognitive performance: a systematic review. *PeerJ*. 2022;10:13604.

Quick jump to page content

- [Main Navigation](#)
- [Main Content](#)
- [Sidebar](#)
  
- [Register](#)
- [Login](#)

[JKKI : Jurnal Kedokteran dan Kesehatan Indonesia](#)

- [Home](#)
- [Current](#)
- [Archives](#)
- [Announcements](#)
- [About](#)

[SEARCH](#)





e-ISSN 2527-2950  
p-ISSN 2085-4145

## JURNAL KEDOKTERAN DAN KESEHATAN INDONESIA

### Indonesian Journal of Medicine and Health

Journal homepage: [www.journal.uii.ac.id/index.php/JKKI](http://www.journal.uii.ac.id/index.php/JKKI)

JKKI	Volume -	Number -	pp -	Jogjakarta	ISSN 2085-4145
------	----------	----------	------	------------	-------------------

Published by  
Faculty of Medicine  
Islamic University of Indonesia



## Journal Description

**JKKI: Jurnal Kedokteran dan Kesehatan Indonesia** is a peer-reviewed journal in the field of medical and health sciences. This journal is designed as a place of dissemination of information and scientific knowledge, which publishes three times a year at April, August and December. It publishes original article, article review, and case report. These comprise of biomedical sciences, clinical medicine, public health sciences, and medical science education.

This journal has been established since 2009 as *JKKI: Jurnal Kedokteran dan Kesehatan Indonesia* (Print ISSN 2085-4145, Online ISSN 2527-2950) managed by the Faculty of Medicine, [Universitas Islam Indonesia](#).

We are pleased to announce that the **JKKI: Jurnal Kedokteran dan Kesehatan Indonesia** (JKKI) has been **accredited** by Ministry of Research, Technology and Higher Education of the Republic Indonesia with Decree No. 164/E/KPT/2021. JKKI was accepted at the **S2** level in **SINTA** ([Science and Technology Index](#)). Articles published in the journal will be more visible and will most likely receive a higher citation.

- [Previous Issue](#)
- [Current Issue](#)
- [Upcoming Issue](#)

JKKI, Vol 14, No 1, (2023)

**Published:** April 5, 2023

[Technomedicine 5.0: The art of metaverse in healthcare](#)

[10.20885/JKKI.Vol14.Iss1.art1](#)

Dito Anurogo <sup>(1)</sup>

(1) 1) Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Makassar, Makassar, Indonesia, 2) International PhD Program for Cell Therapy and Regeneration Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan.

1-2

[Pdf](#)

Read Statistic: 325

[Analysis of hematological parameters in bladder cancer in early and advanced stages at Dr. Sardjito Hospital](#)

[10.20885/JKKI.Vol14.Iss1.art2](#)

*Hidayu Permata Hardi<sup>(1)</sup>, Indrawarman Soerohardjo<sup>(2)</sup>, Ahmad Zulfan Hendri<sup>(3)</sup>*

*(1) Division of Urology, Department of Surgery, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Dr. Sardjito Hospital, Sleman, Indonesia ,*

*(2) Division of Urology, Department of Surgery, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Dr. Sardjito Hospital, Sleman, Indonesia ,*

*(3) Division of Urology, Department of Surgery, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Dr. Sardjito Hospital, Sleman, Indonesia*

3-8

[Pdf](#)

Read Statistic: 407

[The effect of murottal Al Qur'an on the dysmenorrhea intensity among senior high school students](#)

[10.20885/JKKI.Vol14.Iss1.art3](#)

*Fathimah Az Zahra<sup>(1)</sup>, Tri Wahyuliati<sup>(2)</sup>*

*(1) Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Bantul, Indonesia ,*

*(2) Departement of Neurology, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Yogyakarta, Indonesia*

9-15

[Pdf](#)

Read Statistic: 304

[Analysis of C-reactive protein/albumin ratio as a predictor of mortality in sepsis patients](#)

[10.20885/JKKI.Vol14.Iss1.art4](#)

*Uswatun Hasanah<sup>(1)</sup>, Irdah Handayani<sup>(2)</sup>, Asvin Nurulita<sup>(3)</sup>*

*(1) Department of Clinical Pathology, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia ,*

*(2) Dr. Wahidin Sudirohusodo Province General Hospital Makassar, Makassar, Indonesia ,*

*(3) Clinical Pathology Medical Specialist Program, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia*

16-23

[Pdf](#)

Read Statistic: 263

[Analysis of demographic factors and anaemia of the incidence of spontaneous abortion](#)

[10.20885/JKKI.Vol14.Iss1.art5](#)

Farikha Dewi Larasati <sup>(1)</sup>, Sri Winarni <sup>(2)</sup>, Atik Mawarni <sup>(3)</sup>, Farid Agushyana <sup>(4)</sup>

(1) Department of Biostatistics and Population, Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,

(2) Department of Biostatistics and Population, Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,

(3) Department of Biostatistics and Population, Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia ,

(4) Department of Biostatistics and Population, Faculty of Public Health, Universitas Diponegoro, Semarang, Indonesia

24-30

[Pdf](#)

Read Statistic: 260

[Relationship between age, education and blood zinc levels, with cognitive function in menopausal women](#)

[10.20885/JKKI.Vol14.Iss1.art6](#)

Gea Pandhita <sup>(1)</sup>, Martiem Mawi <sup>(2)</sup>, Mieke Nuryani <sup>(3)</sup>

(1) 1)Department of Neurology, Faculty of Medicine, Universitas Muhammadiyah Prof. Dr. HAMKA, Jakarta, Indonesia, 2)Department of Epidemiology, Faculty of Medicine, Universitas Muhammadiyah Prof. Dr. HAMKA, Jakarta, Indonesia ,

(2) Department of Physiology, Faculty of Medicine, Universitas Muhammadiyah Prof. Dr. HAMKA, Jakarta, Indonesia ,

(3) Department of Occupational Health, Faculty of Medicine, Universitas Muhammadiyah Prof. Dr. HAMKA, Jakarta, Indonesia

31-36

[Pdf](#)

Read Statistic: 199

[Relationship between sociodemographic characteristics, knowledge, and behaviours to prevent COVID-19 among students of Manarul Huda Islamic Boarding School](#)

[10.20885/JKKI.Vol14.Iss1.art7](#)

Caecielia Makaginsar <sup>(1)</sup>, Meta Maulida Damayanti <sup>(2)</sup>, Yuniarti <sup>(3)</sup>, Siska Nia Irasanti <sup>(4)</sup>

(1) Department of Medical Education, Bioethics, Humanities and Islamic Inset, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia ,

(2) Pathology Anatomy Department, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia ,

(3) Anatomy Department, Faculty of Medicine Universitas Islam Bandung, Bandung Indonesia

(4) Public Health Department, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

37-46

[Pdf](#)

Read Statistic: 130

[Functional outcome in knee osteoarthritis after dextrose prolotherapy intervention: A severity-based pilot study](#)

[10.20885/JKKI.Vol14.Iss1.art8](#)

Yose Waluyo <sup>(1)</sup>, Sari Rajwani Artika <sup>(2)</sup>, Insani Nanda Wahyuni <sup>(3)</sup>, Endy Adnan <sup>(4)</sup>, Budu <sup>(5)</sup>, Agussalim Bukhari <sup>(6)</sup>

(1) Department of Physical Medicine and Rehabilitation, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia ,

(2) Department of Physical Medicine and Rehabilitation, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia ,

(3) Department of Physical Medicine and Rehabilitation, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia ,

(4) Rheumatology Subdivision, Department of Internal Medicine, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia ,

(5) Department of Ophthalmology, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia ,

(6) Department of Clinical Nutrition, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia

47-54

[Pdf](#)

Read Statistic: 147

[Growth hormone administration and exercise combination increase serum IGF-1 levels in perimenopausal wistar rats](#)

[10.20885/JKKI.Vol14.Iss1.art9](#)

Ni Luh Kadek Alit Arsani <sup>(1)</sup>, Luh Putu Ratna Sundari <sup>(2)</sup>

(1) Faculty of Medicine, Universitas Pendidikan Ganesha, Bali, Indonesia ,

(2) Department of Physiology, Faculty of Medicine, Universitas Udayana, Bali, Indonesia

55-60

[Pdf](#)

Read Statistic: 153

[Effects of high glucose, egg yolk intake, calorie restriction, and air pollution on rats' fat cell](#)

[10.20885/JKKI.Vol14.Iss1.art10](#)

Diniwati Mukhtar <sup>(1)</sup>, Aan Royhan <sup>(2)</sup>, Hasna Luthfiah Fitriani <sup>(3)</sup>, Melasari Marantika <sup>(4)</sup>, Dita Safira Salsabila <sup>(5)</sup>, Lena Fitriyana <sup>(6)</sup>, Febri Irwansyah <sup>(7)</sup>

(1) Faculty of Medicine, Yarsi University, Jakarta, Indonesia ,

(2) Faculty of Medicine, Yarsi University, Jakarta, Indonesia. ,

- (3) Faculty of Medicine, Yarsi University, Jakarta, Indonesia. ,  
(4) Faculty of Medicine, Yarsi University, Jakarta, Indonesia. ,  
(5) Faculty of Medicine, Yarsi University, Jakarta, Indonesia. ,  
(6) Faculty of Medicine, Yarsi University, Jakarta, Indonesia. ,  
(7) Faculty of Medicine, Yarsi University, Jakarta, Indonesia.

61-68

[Pdf](#)

Read Statistic: 182

### [Managing fractures in geriatrics: Current approaches and update](#)

[10.20885/JKKI.Vol14.Iss1.art11](#)

Tito Sumarwoto <sup>(1)</sup>, Dhimas Hartanto <sup>(2)</sup>, Pamudji Utomo <sup>(3)</sup>

- (1) Department of Orthopaedic & Traumatology, Faculty of Medicine, Universitas Sebelas Maret - Prof. Dr. R. Soeharso Orthopaedic Hospital, Surakarta, Indonesia ,  
(2) Department of Orthopaedic & Traumatology, Faculty of Medicine, Universitas Sebelas Maret - Prof. Dr. R. Soeharso Orthopaedic Hospital, Surakarta, Indonesia ,  
(3) Department of Orthopaedic & Traumatology, Faculty of Medicine, Universitas Sebelas Maret - Prof. Dr. R. Soeharso Orthopaedic Hospital, Surakarta, Indonesia

69-81

[Pdf](#)

Read Statistic: 191

### [The potential of sweet orange \(Citrus sinensis\) in cardiovascular health: A literature review](#)

[10.20885/JKKI.Vol14.Iss1.art12](#)

Gabriella Stephanie Minami <sup>(1)</sup>, Ellya Catharine Lumbantoruan <sup>(2)</sup>, Puteri <sup>(3)</sup>, Risa Nuraini <sup>(4)</sup>, Joan Carmen Harianto <sup>(5)</sup>, Andhi Fahrurroji <sup>(6)</sup>

- (1) Faculty of Medicine, Universitas Tanjungpura, Pontianak, Indonesia ,  
(2) Faculty of Medicine, Universitas Tanjungpura, Pontianak, Indonesia ,  
(3) Faculty of Medicine, Universitas Tanjungpura, Pontianak, Indonesia ,  
(4) Faculty of Medicine, Universitas Tanjungpura, Pontianak, Indonesia ,  
(5) Faculty of Medicine, Universitas Tanjungpura, Pontianak, Indonesia ,  
(6) Department of Pharmaceutics and Technology of Pharmacy, Faculty of Medicine, Universitas Tanjungpura, Pontianak, Indonesia

82-94

[Pdf](#)

Read Statistic: 364

### [Association between melanin and vitamin D: A systematic review](#)

[10.20885/JKKI.Vol14.Iss1.art13](#)



Andre Colin Hartono <sup>(1)</sup>, Veronika Maria Sidharta <sup>(2)</sup>, Yunisa Astiarani <sup>(3)</sup>, Regina Regina <sup>(4)</sup>

(1) Bachelor of Medicine Study Programme, School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia ,

(2) Department of Histology, School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia, Jakarta, Indonesia ,

(3) Department of Public Health and Nutrition, School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia, Jakarta, Indonesia ,

(4) Department of Dermatology and Venereology, School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia, Jakarta, Indonesia

95-103

[Pdf](#)

Read Statistic: 193

[Successful combination therapy of low-dose aspirin and unfractionated heparin in recurrent pregnancy loss: A case report](#)

[10.20885/JKKI.Vol14.Iss1.art14](#)

Adhi Pribadi <sup>(1)</sup>

(1) Division of Maternalfetal, Department of Obstetrics and Gynecology, Universitas Padjadjaran, Bandung, Indonesia

104-110

[Pdf](#)

Read Statistic: 140

[Successful systemic propranolol treatment for periocular capillary hemangioma in Indonesian infant girls: A case series](#)

[10.20885/JKKI.Vol14.Iss1.art15](#)

Neni Anggraini <sup>(1)</sup>, Mutmainah <sup>(2)</sup>

(1) Department of Ophthalmology, Faculty of Medicine, University of Indonesia/ Cipto Mangunkusumo General Hospital, Jakarta, Indonesia ,

(2) Department of Ophthalmology, Faculty of Medicine, University of Indonesia/ Cipto Mangunkusumo General Hospital, Jakarta, Indonesia

111-117

[Pdf](#)

Read Statistic: 125

[VIEW ALL ISSUES](#)



**Make Submission**

## ACCREDITATION

---



This journal has been accredited by the National Journal Accreditation (ARJUNA) Managed by the Ministry of Research and Technology/National Research and Innovation Agency (Kemenristek/BRIN), according to the Decree No. 164/E/KPT/2021

[Aims & scopes](#)

[Submission](#)

[Manuscript template](#)

[Publication ethics](#)

[Plagiarism](#)

[Article processing charges](#)

[Editorial team](#)

[Indexing](#)

[Statistics](#)

**Editorial Team**

---



**Novyan Lusiyana**  
**Editor in Chief**

Department of Parasitology, Faculty of Medicine, Universitas Islam Indonesia  
[0000-0002-3254-1974](tel:0000-0002-3254-1974)



**Miranti Dewi Pramaningtyas**  
**Managing Editor**

Department of Physiology, Faculty of Medicine, Universitas Islam Indonesia  
[0000-0003-1215-881X](tel:0000-0003-1215-881X)



**Asri Hendrawati**  
**Associate Editor**

Department of Biochemistry, Faculty of Medicine, Universitas Islam Indonesia  
[0000-0003-3084-1022](tel:0000-0003-3084-1022)



**Evy Sulistyoningrum**  
**Substantive Editor**

Department of Histology, Faculty of Medicine, Universitas Islam Indonesia  
[0000-0001-7487-5932](tel:0000-0001-7487-5932)

## WEB STATISTICS

---



000029468

[View My Stats](#)

## RECOMMENDED TOOLS

---



Published Volume By Year

2023

2022

2021

2020

2019

Keywords

COVID-19 antioxidant knowledge Covid-19 Indonesian risk factors Yogyakarta stroke prevention compliance BMI Body increase cast kidney renal

weight

Previous





Next





# JKKI : JURNAL KEDOKTERAN DAN KESEHATAN INDONESIA

## Address

Fakultas Kedokteran, Universitas Islam Indonesia  
Jl. Kaliurang Km 14,5 Yogyakarta, Indonesia

## Contact Info

Telp. (0274) 898444 ext. 2050  
Fax. (0274) 898580 ext. 2007  
[081238884726](tel:081238884726)

JKKI at <http://journal.uji.ac.id/JKKI/> is licensed under  a Creative Commons Attribution 4.0 International License.  
Mostcitedcrossrefblockplugin was unvalidated product, [Click here to support us](#)