

Journal of Ecophysiology and Occupational Health

 All

[Home](#)
[Current](#)
[Archives](#)
[Authors](#)
[Institutions](#)
[Editorial Board](#)
[Submit Article](#)
[Online First](#)

Editorial Board

Patrons

Dr. R. C. Dalela

Lucknow

Dr. Alok Dhawan

Lucknow

Dr. S. Ayyappan

New Delhi

Dr. Y. K. Gupta

New Delhi

Dr. H. N. Dutta

New Delh

Quick Links

Abstracting and Indexing

Instructions to Authors

Pre-submission checklist

Publication Fees

Journal Policies and Ethics

Submission Process

Information for Reviewers

Email Alerts

Editor-in-Chief

Dr. Krishna Gopal

Lucknow

Chief Editor

Prof. B Sharma

Allahabad

Editors

Dwijendra Upadhyay

Lucknow

Dr. A. B. Pant

Lucknow

Dr. Anil K. Agrawal

Lucknow

Advisory Editors

Dr. Virendra Misra

Dr. S. J. S. Flora

Archives

[\[Expand All\]](#)

[+ 2019](#)

[+ 2018](#)

[+ 2017](#)

[+ 2016](#)

[+ 2015](#)

[+ 2014](#)

[+ 2013](#)

[+ 2012](#)

[More...](#)



Lucknow

Dr. A. K. Singh

Bhimtal

Dr. H. C. Sharma

Solan

Dr. K. Vijaykumar

Gulbarga

Raebareli

Prof. C. P. M. Tripathi

Gorakhpur

Prof. A. K. Chaubey

Meerut

Prof. Ram Chandra

Lucknow

Technical Editor

Mr. Suyash Tripathi

Lucknow

Mr. Amit Kumar Dubey

Lucknow

Associate Editors

Prof. Frans D. Suyatna

Indonesia

Dr. Zakir A. Ansari

Prof. Divakar Dalela

Lucknow

Prof. R. S. Yadav



Goa

Dr. Madhu Tripathi

Lucknow

Dr. D. Kar Chaudhary

Lucknow

Dr. V. N. Mishra

Australia

Prof. S. V. S. Rana

Meerut

Prof. J. L. Bersillon

France

Prof. P. N. Saxena

Agra

Prof. Ramesh Chandra

Lucknow

Kolhapur

Prof. Rajavashisth Tripathi

USA

Dr. Rana P. Singh

Lucknow

Prof. C. D. Nwani

Nigeria

Prof. U. N. Dwivedi

Lucknow

Dr. Gu Zu-Wei

China

Prof. R. K. Sinha

Patna

Journal of Ecophysiology and Occupational Health

All

Search

Home

Current

Archives

Authors

Institutions

Editorial Board

Submit Article

Online First

Volume 19, Issue 1&2, June 2019

Table of Contents

Articles

Reviewing Influence of Information Asymmetry on Corporate Tax Avoidance

Manijeh Kord Manjiri, Mahdi Ahmadipanah


DOI: [10.18311/jeoh/2019/23075](https://doi.org/10.18311/jeoh/2019/23075), Pagination: 1-11

 [Full-Text PDF](#) 


Abstract Views: | Pdf Views:

An Investigation into the Effect of Patient Safety Indicators on Patient Satisfaction: A Case Study of

Quick Links

 [Abstracting and Indexing](#)

 [Instructions to Authors](#)

 [Pre-submission checklist](#)

 [Publication Fees](#)

 [Journal Policies and Ethics](#)

 [Submission Process](#)

 [Information for Reviewers](#)

 [Email Alerts](#)

An Investigation into the Effect of Patient Safety Indicators on Patient Satisfaction: A Case Study of Shooshtar Maternity Hospital in Shiraz

Ayeh Negahbani, Shaghayegh Vahdat

DOI: [10.18311/jeoh/2019/23479](https://doi.org/10.18311/jeoh/2019/23479), Pagination: 12-15

 [Full-Text PDF](#) 

Abstract Views: | Pdf Views:

A Clinical Evaluation of Daruharidra (*Berberis aristata* D.C.) in cases of Giardiasis

Syed Sadiq Abbas, Agha Parvez Masih, S. Shabihe Raza Baqri, Mohd Yusuf, Kumar Gaurav Bajpai

DOI: [10.18311/jeoh/0/23778](https://doi.org/10.18311/jeoh/0/23778), Pagination: 16-23

 [Full-Text PDF](#) 

Abstract Views: | Pdf Views:

Length Weight Relationship and Condition Factor of Cultured Rainbow Trout, *Oncorhynchus mykiss* (Walabum, 1792)

Ishtiyag Ahmad, Imtiaz Ahmed

DOI: [10.18311/jeoh/0/22304](https://doi.org/10.18311/jeoh/0/22304), Pagination: 24-27

 [Full-Text PDF](#) 

Abstract Views: | Pdf Views:

Systolic Blood Pressure and Physical Fitness in Security Personnel: A Cross Sectional Study

Nadya Gratia Juliawan, Listya Tresnanti Mirtha, Muchtarudin Mansyur

DOI: [10.18311/jeoh/0/22251](https://doi.org/10.18311/jeoh/0/22251), Pagination: 28-30

 [Full-Text PDF](#) 

Abstract Views: | Pdf Views:

Email Alerts


 CrossMark Policy

Archives

[\[Expand All\]](#)

 [2019](#)

 [2018](#)

 [2017](#)

 [2016](#)

 [2015](#)

 [2014](#)

 [2013](#)

 [2012](#)

[More...](#)

Current Issue

ATOM 1.0

RSS 2.0



Isometric was better than Isotonic Neck Stretching to Reduce Pain and Creatinine Kinease in Non-specific Neck Pain in Caring Students of Swadhyaseni Middle High School, Badung, Bali

Thomas Eko Purwata, I. Komang Sri Mahendra Putra, I. G. N. Purna Putra, I. Putu Eka Widyadharma

DOI: [10.18311/jeoh/0/23070](https://doi.org/10.18311/jeoh/0/23070), Pagination: 31-36

 Full-Text PDF 

Abstract Views: | Pdf Views:

An Analytical Framework for the Determinants of Infant Mortality based on 2005 – 06 NFHS Data in India

Anu Sirohi, Piyush Kant Rai

DOI: [10.18311/jeoh/0/23122](https://doi.org/10.18311/jeoh/0/23122), Pagination: 37-47

 Full-Text PDF 

Abstract Views: | Pdf Views:

Assessment of Socio-Demographic Factors with Knowledge and Attitiudes of Pakistani Married Women (15-49 Years) towards HIV/AIDS Transmission

Rukiya Tariq, Quratulain Bashir, Tanzeel Ul Rahman

DOI: [10.18311/jeoh/0/22518](https://doi.org/10.18311/jeoh/0/22518), Pagination: 48-55

 Full-Text PDF 

Abstract Views: | Pdf Views:

Effect of Vermicompost on the Growth Performance of Indian Major Carps in Fish Rearing Pond

Suresh Kumar, Sunita Godara

DOI: [10.18311/jeoh/2019/23378](https://doi.org/10.18311/jeoh/2019/23378), Pagination: 56-64

 Full-Text PDF 

Abstract Views: | Pdf Views:



Ovarian Impairment in Fish, *Channa punctata* (Bloch) on Exposure to Mercuric Chloride, Caldan 50 SP and their Mixture

Shalini Rai, Kamlesh K. Yadav, Sunil P. Trivedi

DOI: [10.18311/jeoh/2019/22787](https://doi.org/10.18311/jeoh/2019/22787), Pagination: 65-70

 Full-Text PDF 

Abstract Views: | Pdf Views:

Di (2-Ethylhexyl) Phthalate Induced Toxicological Effects on Reproductive System of Female Mice *Mus-Musculus*

Anjali Singh, Ravish Kumar, Jitendra Kumar Singh, Kumar Sanbhav Singh, Tanuja

DOI: [10.18311/jeoh/0/23333](https://doi.org/10.18311/jeoh/0/23333), Pagination: 71-75

 Full-Text PDF 

Abstract Views: | Pdf Views:

Erratum

Functional Differentiation of Neural Stem Cells into Neuronal Subtypes: A Biological Tool for Developmental Neurotoxicity Studies

Ankita Pandey

Pagination: 76-76

 Full-Text PDF 

Abstract Views: | Pdf Views:

Length-Weight Relationship and Condition Factor of *Schizothorax plagiostomus* found in River Jhelum from Kashmir Valley

Zubair Ahmad Sheikh

Journal of Ecophysiology and Occupational Health

All

Search

Home

Current

Archives

Authors

Institutions

Editorial Board


Submit Article


Online First

About the Journal



Journal of Ecophysiology and Occupational Health

 Editor-in-Chief : Dr. Krishna Gopal

 Online ISSN : 0974-0805

 Print ISSN : 0972-4397


 Frequency : Quarterly

 Published since : 2010

 Publisher(s) : Informatics Publishing Limited and The Academy of Environmental Biology, Lucknow

Subscription


Quick Links

 Abstracting and Indexing

 Instructions to Authors

 Pre-submission checklist

 Publication Fees

 Journal Policies and Ethics

 Submission Process

 Information for Reviewers

 Email Alerts

Journal of Ecophysiology and Occupational Health (JEOH) is a Quarterly publication. The Journal is open to every one who

has a scientific, technological, practical interest and inclination in the field of Science and Technology, Environmental Pollution, occupational health and toxicology.

The Journal is included in Web of Science BIOSIS Previews, EMBASE, The Journal is Indexed in U. S. National Library of Medicine, ICI and Annexure-II-Anna University, UGC Mandate.

Aims and Scope

Journal of Ecophysiology
and Occupational Health



SJR 2018
0.1



powered by scimagojr.com

[CrossMark Policy](#)

Archives

[\[Expand All\]](#)

[+ 2019](#)

[+ 2018](#)

[+ 2017](#)

[+ 2016](#)

[+ 2015](#)

[+ 2014](#)

[+ 2013](#)

[+ 2012](#)

[More...](#)

Current Issue

ATOM	1.0
RSS	2.0
RSS	1.0



Journal of Ecophysiology and Occupational Health

Country[India](#) -  [SJR Ranking of India](#)**Subject Area and Category**[Environmental Science](#)[Ecology](#)[Pollution](#)[Medicine](#)[Public Health, Environmental and Occupational Health](#)**Publisher**[Academy of Environmental Biology](#)**Publication type**[Journals](#)**ISSN**

09724397

Coverage

2003-2016 (cancelled)

Scope

The journal publishes in English language original research/review papers covering all areas of environmental sciences, environmental pollution and related disciplines. Other areas are

8

H Index

Scope

The journal publishes in English language original research/review papers covering all areas of environmental sciences, environmental pollution and related disciplines. Other areas are toxicology in relation to plants, animals and men including general toxicology, reproduction toxicology, metabolism and mechanism of action, safety assessment, clinical and veterinary toxicology, ecophysiology, biodiversity, risk assessment, risk management, cleaner technology, behavioral toxicology, experimental toxicology of pharmaceutical, surgical, food, agricultural and industrial products, impact assessment of pollution, responses to toxic agents and community.






[Homepage](#)




[Join the conversation about this journal](#)



Quartiles






- Cites / Doc. (4 years)
- Cites / Doc. (3 years)
- Cites / Doc. (2 years)





● Citable documents ● Non-citable documents +



● Cited documents ● Uncited documents +



**Journal of Ecophysiology
and Occupational Health**

Q4

Ecology

best quartile

SJR 2018

0.1

powered by scimagojr.com

← Show this widget in
your own website

Just copy the code below
and paste within your html
code:

```
<a href="https://www.scimagojr.c
```




Isometric was better than Isotonic Neck Stretching to Reduce Pain and Creatinine Kinease in Non-specific Neck Pain in Caring Students of Swadhyaseni Middle High School, Badung, Bali

Thomas Eko Purwata*, I. Komang Sri Mahendra Putra, I. G. N. Purna Putra and I. Putu Eka Widyadharma

Faculty of Medicine, Department of Neurology, Udayana University/Sanglah Hospital, Bali, Indonesia;
thomaseko2000@yahoo.com

Abstract

Neck pain is frequently complained by workers with static neck position. Complaints accompanied by discomfort around the neck muscles without any sensory disturbances or radiating pain, without specific causes such as history of neck trauma, neck malignancy or anatomical structural abnormalities. The samples were taken from carving students of Swadhyaya Middle School Badung. We evaluated 18 students divided into 2 groups: one with isometric neck stretches and another with isotonic neck stretches, 9 subjects each. The paired t-test found the number of decrease in NPRS in the isometric group was greater than isotonic group and the difference was statistically significant. The results of the unpaired student t-test obtained a mean decrease in NPRS in the isometric group greater than isotonic with a value of $P = 0.00$ ($p < 0.01$). Creatine kinase levels in the isometric group showed no significant difference, whereas creatine kinase levels in the isotonic group showed that the increase was not significantly different, the results of the independent student t test showed values not significantly different from the isometric and isotonic groups with the intention of $p = 0.284$ ($p > 0.01$). These results indicate isometric stretching is more effective in reducing the scale of NPRS pain in non-specific neck pain compared to isotonic stretches, isometric and isotonic stretching techniques are equally less effective in reducing creatine kinase levels.

Keywords: Creatine Kinase, Isometric, Isotonic, Nonspecific Neck Pain, Pain Scale

1. Background

Neck pain is a complaint that is often felt by workers, especially workers with a certain neck position in a long enough period of time resulting in excessive contraction of the neck muscles. Various types of work can lead to complaints of neck pain especially for workers with wrong body position that affects the position of the neck resulting in non-specific neck pain.

Approximately 54% of individuals in the world have complained of neck pain in a period of six months and this incidence continues to increase in prevalence around 6% to 22% and increases in the old age group of around 38%. The costs incurred by workers for treatment of neck pain are sufficiently high. Research showed that the prevalence of musculoskeletal pain in the neck in the community for one year was 40% and more prevalent in women, while the prevalence of neck pain in workers ranges from 6-76 % and mostly in women¹⁻³.

The research conducted by Nemade⁴ obtained musculoskeletal complaints in the form of neck pain in brick workers in India by 14% from 159 workers, while the research conducted by Sihawong et al.⁵ in 567 office workers provided stretching training in the treatment group totaling 285 people and without treatment of neck stretching in 282 people as the control group, resulted in a decrease in the incidence of neck pain in office workers who received neck stretch treatment^{4,5}.

Body posture and muscle contraction during work can cause trauma to various aspects of the musculoskeletal system, especially the vertebral column muscles. Forward head posture occurs due to increased dorsal spinal kyphosis, where the head is in front of the center of gravity so that the neck load in Broocrease. Dropping shoulders affect the cervical spine, and excessive hyperextension or hyperflexion in the neck also affect the incidence of neck pain in workers⁶.

Research conducted by Nejati et al.⁷ in Iran on 46 people without neck pain and 55 people with neck pain who worked in offices with computers found that head forward posture

correlated with the incidence of neck pain while shoulder posture had no effect⁷. Research conducted by Mufti on the craftsmen of songket weaving in Pandai Singkek⁸, West Sumatra regarding the study of workers' posture showed that all workers complained of neck pain (100%), followed by upper back pain⁸.

Research conducted by Palmer et al.⁹ in England, Scotland, and Wales in 12,907 respondents aged 16-64 years showed that people who worked with upper arms and shoulders more than one hour per day had a significant association with the onset of neck pain, while a prospective study conducted by Ariens et al. found that workers who worked in a sitting position with a static neck position when working was a risk factor for neck pain⁹.

Excessive muscle activity will cause damage to muscle cells. This condition is evidenced by high levels of Creatine Kinase (CK) in blood serum. After heavy activity the muscles need either active or passive recovery. Active recovery by mild stretches can reduce adverse metabolic results of muscles including reducing levels of creatine kinase and lactic acid in the muscles¹⁰.

Stretching is one way that is often done to reduce discomfort due to pain in the muscles of the neck and shoulders. Stretching provides benefits to strengthen the muscles of the neck and shoulders, increases flexibility and movement of the neck muscles, increases blood flow to the muscles, increases production energy, reduce pain and are relatively easy to do and do not require special time and place. Research conducted by Sowmya et al. in patients with non-specific chronic neck pain showed an isometric stretching technique as effective as dynamic or isotonic stretching to reduce non-specific neck pain¹¹. Gross et al. obtained moderate results in patients with non-specific neck pain who were given isometric stretches¹².

Neck pain can be caused by all components of the neck starting from the musculoskeletal system and nervous system. Neck pain in workers is generally more often caused by musculoskeletal disorders both due to tension and excessive contraction of muscles and ligaments around the neck and giving symptoms of pain and neck stiffness accompanied by muscular pain around the neck.

Neck pain experienced by workers often involves disorders of the musculoskeletal system, this neck pain according to the pathophysiological process including mechanical neck pain or axial neck pain or often called non-specific neck pain as there is no underlying disease or structural anatomic disorder. The symptoms that often accompany non-specific neck pain are stiffness in one side or both sides of the neck, and pain radiating to the head. Nonspecific neck pain is caused purely by the structure of the muscles or musculoskeletal system in

the neck and is often related to body posture or static neck position at work and excessive neck muscle workload when working for a certain period of time^{2,13}.

2. Objectives

To find out the more effective stretching technique between isometric and isotonic to reduce pain scale and creatine kinase levels in non-specific neck pain.

3. Materials and Methods

This research is an experimental study with a pretest-posttest control group design. Subjects was taken from students of Swadhyaya Middle Carving School Badung, Bali from November 2016 to December 2016 by simple random sampling method. Data collection was done by filling in the questionnaire by the subjects and by neurological clinical examinations.

Inclusion criteria were: students following carving art training activities; non-specific neck pain with NPRS 1-6 (mild - moderate); not taking or undergoing any pain relief therapy; filling in inform consent. The exclusion criteria were: neck trauma; history of neck surgery; severe neck pain; regular relaxation; radiculopathy; history of heart disease and hypertension.

The total sample size needed in this study was 18 people with details of 9 as isometric groups and 9 as isotonic groups.

The analysis and presentation of the data used were as follows:

- Descriptive analysis to give an idea of the characteristics of the data obtained in this study such as age, sex, weight, height, pain scale and creatin kinase levels pre- and post- treatment.
- The normality test on numerical data (NPRS) using the Shapiro Wilk test because the number of samples is less than 50.
- Bivariate analysis to compare the mean reduction in NPRS and creatine kinase before and after stretching. Numerical scale data in both groups tested using the Student paired t-test and unpaired t-test when the data was distributed normally and the Wilcoxon test if the data were not distributed normally. The level of significance was set as p value <0.01.
- If drop out > 20% then analysis of intention to threat is carried out. All data were analyzed using the SPSS 20.0 for Windows.

4. Results

4.1 Basic Characteristics of Research Subjects

Based on the sex category, all subjects, 9 in each group, were male (100%). The mean of the overall age category was 14.76 years. The average age of the isometric group is 14.44 years and the average age of the isotonic group is 15.11 years, the mean weight of the isometric group is 47.44 ± 1.74 and isotonic is 47.22 ± 1.64 while the mean height of the isometric group is 150.00 ± 2.42 and the isotonic group is 149.00 ± 3.20 . The initial mean NPRS in isometric group was 4.89 ± 0.601 and after 4 weeks following the isometric stretching exercise the scale of pain dropped to 1.89 ± 0.601 . The initial NPRS in the isotonic group was 4.67 ± 0.707 and the pain scale at the end of the exercise was 3.44 ± 0.527 . The initial CK level in the

isometric group was 170.11 ± 41.963 and at the end of the exercise was 156.00 ± 31.389 while the initial level of CK in the isotonic group was 170.67 ± 52.557 and at the end of the exercise was 177.78 ± 62.532 . The basic characteristics of the research subjects are presented in (Table 1).

4.2 Effects of Isometric and Isotonic Stretching on Decreasing the Scale of NPRS Pain

The effect of stretching on decreasing the NPRS pain scale in patients with non-specific neck pain was assessed by comparing the mean reduction in NPRS between isometric groups and isotonic groups after stretching exercises for 4 weeks. Normality tests on the NPRS reduction in both groups using the Shapiro Wilk test showed normal distribution. The analysis is presented in the following (Tables 2 and 3).

Table 1. Basic characteristics of subjects

Variable	isometric (n=9)	isotonic (n=9)
Mean age (years)	14.44±0,527	14,78±0,441
Sex (male)	9 (100%)	9 (100%)
Weight (kg)	47,44±1,74	47,22±1,64
Height (cm)	150,11±2,42	149,00±3,20
Initial (pre-treatment) NPRS	4,89±0,601	4,67±0,707
Final (post-treatment) NPRS	1,89±0,601	3,44±0,527
Initial (pre-treatment) CK (U/L)	170,11±41,963	170,67±52,557
Initial (pre-treatment) CK (U/L)	156,00±31,389	177,78±62,532

Table 2. Bivariate analysis of Student's paired t-test in mean decrease of NPRS

Group	Initial mean NPRS	Final mean NPRS	Percentage	p-value
Isometric	4,89±0,601	1,89±0,601	61,34%	0,000
Isotonic	4,67±0,707	3,44±0,527	26,39%	0,000

Table 3. Bivariate analysis of NPRS using unpaired t-test

Group	Mean NPRS decrease	p-value
Isometric	3,00±0,707	0,000
Isotonic	1,22±0,441	

4.3 Effects of Isometric and Isotonic Stretches on Decreasing Creatine Kinase Levels

The decrease in creatine kinase levels in the study subjects was calculated by comparing the mean creatine kinase levels

before and after stretching exercise. The normality test using the Shapiro Wilk test showed that the data were distributed normally. The analysis results are presented in the following (Tables 4 and 5).

Table 4. Bivariate analysis of Student's paired t-test of mean decrease of CK

Group	Initial mean CK	Final mean CK	Percentage	p-value
Isometric	170,11±41,963	156,00±31,389	8,29%	0,030
Isotonic	170,67±52,557	177,78±62,532	4.16%	0,709

Table 5. Bivariate analysis of unpaired t-test of decrease in CK level

Group	Mean decrease in CK level	p-value
Isometric	14,11±16,058	0,284
Isotonic	7,11±55,151	

5. Discussion

5.1 Basic Characteristics of Research Subjects

The subjects of this study were obtained from the students of Swadhyaya Middle School, Badung, Bali who experienced non-specific neck pain, selected by using the LEEDS questionnaire Assessment of Neuropathic Symptoms and Signs (LANNS) and simple randomization with coin toss on the students meeting inclusion criteria. A total of 18 students were divided into 9 students as treatment groups with isometric stretches and 9 students as a control group with isotonic stretches.

All subjects were male due to the school curriculum dictated that all male student in the Swadhyaya Middle School taking carving class while the female students taking the sewing class. The average age of the whole group in this study was 14.78 years, with the average age of isometric groups being 14.44 ± 0.527 and the age of the Isotonic group was 14.78 ± 0.441 with the lowest age of 14 years and the highest age of 15 years. The mean age of the isometric and isotonic groups did not differ statistically with a value of $p = 0.165$ ($p > 0.01$).

The mean weight of the isometric and isotonic groups was not significantly different with the value of $p = 0.784$ ($p > 0.01$) and the mean height of the isometric and isotonic groups also did not differ statistically with a value of $p = 0.419$ ($p > 0.01$). Data on the characteristics of the research subjects such as age, weight and height did not differ between isometric groups and isotonic groups because all subjects in this study were attending grade 3 and had never stayed in classes, participating in arts activities. carve regularly at school 3 times a week.

5.2 Effects of Isometric and Isotonic Stretching on Decreasing the scale of NPRS Pain

The mean NPRS in the isometric group at the start of stretching was 4.89 ± 0.601 and at the end of the exercise was 1.89 ± 0.601 . This result indicated a decrease in the mean value of 61.34% and this was statistically significant ($p = 0.000$). In the isotonic group, the initial mean of NPRS was 4.67 ± 0.707 and the final mean was 3.67 ± 0.707 . The decrease in the mean NPRS value in the isotonic group was 26.39% and statistically significant ($p = 0.000$). To determine the effectiveness between isometric and isotonic stretching, an independent student t-test was performed with the results of the mean reduction in the isometric group 3.00 ± 0.707 and the mean decrease in the

isotonic group 1.22 ± 0.441 ($p = 0.000$). This indicated that the isometric stretching technique was more effective in reducing the scale of NPRS pain in patients with non-specific neck pain compared to the isotonic stretching technique. The result of this study was consistent with the results of a study conducted by Gupta et al.¹⁴ who found a decrease in the pain scale of both isometric and isotonic groups in patients with acute non-specific neck pain; in the study the pain scale decreased in an average of 4 or around 72.

The isometric group showed better result compared to the isotonic group. This might be due to the mechanism of the isometric stretch to return muscle fibers, actin and myosin, to the normal position after overlapping between actin and myosin fibers due to excessive muscle contraction (cross link). The condition of cross links in muscle fibers will irritate the A delta and C fibers that is responsible for pain. The return of actin and myosin fibers to their normal position helps repair irregular muscle fibers and will cause the muscles to relax. There is an endogenous release of analgesic so that pain can be reduced or eliminated. In the isotonic group, the degree of improvement was not as high, possibly due to the muscle movements involving varying muscle lengths along with movement of the joints around the neck so metabolism is greater, and more energy is needed. This stretch can help return the position of muscle fiber fibers in a normal position, however the length of muscles changes so that normal conditions are achieved in only a short duration because the muscles contract again.

5.3 Effects of Isometric and Isotonic Stretches on Decreasing Creatine Kinase Levels

The initial mean creatine kinase level in the isometric group was 170.11 ± 41.963 and after 4 weeks stretching was $156.00 \pm 31,389$ with a decrease in creatine kinase level of 8.29%. This result was not statistically significant with a p value = 0.341. In the isotonic group, the initial mean creatine kinase level was 170.67 ± 52.557 and after stretching was 177.78 ± 62.532 ; the results interestingly showed an increase in the average CK level of 5.46%, even though it was not statistically significant ($p = 0.629$). To determine the effectiveness of isometric and isotonic stretching to reduce CK levels, an independent Student t-test was done and the CK reduction in isotonic groups was found to be an average $11 \pm 16,058$ and the mean increase in CK in the isotonic group was 7.11 ± 55.151

with a value of $p = 0.285$ ($p > 0.01$). It can be concluded that both isometric and isotonic stretches were equally ineffective in reducing CK levels in non-specific neck pain. This might be explained by the fact that CK level in the blood was influenced by genetics, sex, age and activity. Because the activity of the subject of research outside the school varied with different intensities so it was possible that the serum CK level differed essentially. Same reasoning may be applied to the increase in the average CK level in the isotonic group, in addition to the mechanism by which isotonic stretching requires greater energy and muscle contraction so that muscle metabolism and energy consumption increases, thus the elevated CK level. All subjects in this study had CK levels below 270 U / L (ages 13-17 years) which showed no muscle damage.

Research conducted by Clarkson et al.¹⁵ obtained an average increase in CK of 34% in the isometric group and 37.6% in the isotonic group. The research conducted by Agha-Alinejad et al.¹⁶ on 28 students who took a physical exercise program to determine the effect of isotonic and isometric stretches after certain activities had increased CK levels. Baird et al.¹⁰ obtained different results, namely a mild isometric stretching exercise program reduced muscle damage and CK levels compared to heavy physical exercise.

The weakness in this study is the small number of samples and the short duration of the study. Another disadvantage is that the subjects in this study were not quarantined so it was difficult to monitor activities carried out outside of school whereas CK levels in the blood were strongly influenced by the activity of the body muscles. Another disadvantage is that in this study the force or energy released during stretching could not be measured with certainty so that the force or energy released by each subject varied.

The strength of this study lies in the homogeneous sample of school children so that samples were easily followed up and bias could be avoided. Stretching techniques were trained by experienced physiotherapy instructors and stretching exercises were carried out in separate spaces between isometric and isotonic groups so as to minimize errors and mimic each other's movements in stretching. All subjects were not exercised or underwent exercise in school during the participation period, thus minimized the influence of other physical activities outside of the neck stretching activities.

6. Conclusion

Based on the results of this study, it can be concluded that isometric neck stretching is better than isotonic neck stretching to reduce pain scale in patients with non-specific neck pain and both isometric and isotonic stretches are equally ineffective in reducing creatine kinase in non-specific neck

pain. Stretching isometric techniques can be used as one of the non-pharmacological management modalities to reduce non-specific neck pain. However, further research should include a larger number of samples, involving female subjects, longer training duration and more specific biomarker measurements for inflammation such as interleukin and other muscle damage biomarkers such as myoglobin, lactate dehydrogenase (LDH) and aspartate aminotransferase (AST).

7. References

1. Ariens GA, van Mechelen W, Bongers PM, Bouter LM, van der Wal G. Psychosocial risk factors for neck pain: A systematic review. *American Journal of Industrial Medicine*. 2001; 39:180–93. [https://doi.org/10.1002/1097-0274\(200102\)39:2<180::AID-AJIM1005>3.0.CO;2-#](https://doi.org/10.1002/1097-0274(200102)39:2<180::AID-AJIM1005>3.0.CO;2-#)
2. Giannoula T, Roy R, Win D, Paul VN. Nonspecific neck pain and evidence based practice. *European scientific Journals*. 2013; 9:6.
3. Masum SB, Hage M, Haque M, Islam MS. Prevalence of neck pain and associated factors among office workers. *The American Journals of Innovative Research and Applied Sciences*. 2014; 1:4.
4. Nemade VV. Musculoskeletal disorders in brick client workers. Incidence and Effectiveness of Workers Education Programme. 2014; 3:3.
5. Sihawong R, Janwantanakul P, Sitthipornvorakul E, Pensri P. Exercise therapy for office workers with nonspecific neck pain: a systematic review. *Journal of Manipulative and Physiological Therapeutics*. 2011; 34:62–71. <https://doi.org/10.1016/j.jmpt.2010.11.005>. PMID:21237409
6. Brook P, Bogduk N. Evidence based management of acute musculoskeletal pain. The National Health and Medical Research Council, First Edition; 2003. p. 14.
7. Nejati P, Lotfian S, Moezy A, Nejati M. The study of correlation between forward head posture and neck pain in Iranian office workers. *International Journal of Occupational Medicine and Environmental Health*. 2015; 28:295–303. <https://doi.org/10.13075/ijomeh.1896.00352>
8. Mufti D, Suryani E, Sari N. Evaluation of work posture among Songket fabric craftsmen at Pandai Sikek. *Scientific Industrial Technique Journal*. 2013; 12:63–71.
9. Palmer KT, Walker-Bone K, Griffin MJ, Syddall H, Pannett B, Coggon D, Cooper C. Prevalence and occupational associations of neck pain in the British population. *Scandinavian Journal of Work, Environment & Health*. 2001; 27:49–56. <https://doi.org/10.5271/sjweh.586>. PMID:11266146
10. Baird MF, Graham SM, Baker JS, Bickerstaff GF. Creatine-kinase- and exercise-related muscle damage implications for muscle performance and recovery. *Journal of Nutrition and Metabolism*. 2012; 2012:13. <https://doi.org/10.1155/2012/960363>. PMID:22288008. PMCID:PMC3263635.

11. M.V S. Isometric neck exercises versus dynamic neck exercises in chronic neck pain; 2014.
12. Gross A, Kay TM, Paquin JP, Blanchette S, Lalonde P, Christe T, et al. Exercises for mechanical neck disorders. Cochrane Database of Systematic Reviews. 2015. <https://doi.org/10.1002/14651858.CD004250.pub5>
13. Binder A. The diagnosis and treatment of nonspecific neck pain and whiplash. *Europa Medicophysica*. 2007; 43:79–89. PMID:17369782
14. Gupta S, Jaiswal P, Chhabra D. A comparative study between postisometric relaxation and isometric exercises in non-specific neck pain. 2012.
15. Clarkson PM, Byrnes WC, McCormick KM, Turcotte LP, White JS. Muscle soreness and serum creatine kinase activity following isometric, eccentric, and concentric exercise. *International Journal of Sports Medicine*. 1986; 7:152–5. <https://doi.org/10.1055/s-2008-1025753>. PMID:3733311
16. Agha-Alinejad H, Shamsi M, Azarbayjan M, Rahimi M, Jafarabadi MA, Tofighi L, Mirani M. The effects of active recovery on serum IL-6, IL-8, IL-10 and CK concentrations after eccentric strenuous exercise in active female. *Iranian Journal of Endocrinology and Metabolism*. 2010; 11(5):553–60.