

The screenshot shows the website for Veterinary World, an open access and peer-reviewed journal. The page features a blue header with the journal's logo and name. Below the header, there is a navigation menu with links for Home, Editorial Board, For authors, FAQ, For reviewers, Archive, and Open access policy. The main content area is divided into two sections: Editor-in-Chief and Editorial board. The Editor-in-Chief section lists Anjum V. Sherasiya, an Ex-Veterinary Officer from Gujarat State, India, with her ORCID ID. The Editorial board section lists six members: Shambhunath Choudhary (USA), Suresh H. Basagoudanavar (India), R. G. Jani (India), G. N. Gongal (India), Md. Tanvir Rahman (Bangladesh), and Deepmala Agarwal (USA), each with their respective affiliations and ORCID IDs. The browser's taskbar at the bottom shows various open applications and the system clock.

Editor-in-Chief

Anjum V. Sherasiya - Ex-Veterinary Officer, Department of Animal Husbandry, Gujarat State, India

<https://orcid.org/0000-0002-1598-1820>

Editorial board

Shambhunath Choudhary - Veterinary Pathologist II, Pathology | Charles River Laboratories, 640 North Elizabeth Street, Spencerville, OH 45887, USA

Suresh H. Basagoudanavar - FMD Vaccine Research Laboratory, Indian Veterinary Research Institute, Bangalore- 560024, Karnataka, India
<https://orcid.org/0000-0001-7714-3120>

R. G. Jani - Ex-Coordinator of Wildlife Health, Western Region Centre, Indo-US Project, Department of Veterinary Medicine, Veterinary College, Anand Agricultural University, Anand - 388001, Gujarat, India

G. N. Gongal - WHO South -East Asia Regional Office, New Delhi -110002, India

Md. Tanvir Rahman Department of Microbiology and Hygiene, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh
<https://orcid.org/0000-0001-5432-480X>

Deepmala Agarwal - Cancer Prevention Laboratory, Pennington Biomedical Research Center, Baton Rouge, LA, USA
<https://orcid.org/0000-0002-9725-6111>

Fouad Kasim Mohammad - Department of Pharmacology & Toxicology, Vice President for Administrative & Financial Affairs, University of Mosul, P.O. Box 11136, Mosul, Iraq

Nicole Borel - Department of Pathology, Vetsuisse Faculty, University of Zurich, CH-8057 Zurich, Switzerland

B. A. Lubisi - Virology, MED Programme, ARC - Onderstepoort Veterinary Institute, No. 100 Old Soutpan Road, Onderstepoort, Tshwane, 0110, South Africa

Kumar Venkitanarayanan - Graduate Programs Chair, Honors and Pre-Vet Programs Advisor, Department of Animal Science, University of Connecticut, Storrs, CT 06269, U.S.A.

Kemin Xu - Department of Veterinary Medicine, University of Maryland, College Park College Park, MD, 20742, USA

Vassilis Papatsiros - Faculty of Veterinary Medicine, Department of Medicine (Porcine Medicine), University of Thessaly, Thessaly, Greece

K. P. Singh - School of Medicine and Dentistry, University of Rochester, Department of Environmental Medicine, Room: 4-6820, 601 Elmwood Avenue, Box-EHSC, Rochester, New York-14620, USA

Ashok K. Chockalingam - Division of Applied Regulatory Science, U.S. Food and Drug Administration, 10903, New Hampshire Avenue, Silver Spring, Maryland 20993, USA
<https://orcid.org/0000-0002-1360-8926>

Ashutosh Wadhwa - Center for Global Safe Water, Sanitation and Hygiene at Emory University, Hubert Department of Global Health, Rollins School of Public Health, Emory University, 1518 Clifton Rd. NE, Atlanta, GA 30322, USA

Luiz Otavio Pereira Carvalho - Laboratory of Immunomodulation and Protozoology, Oswaldo Cruz Institute, Ministry of health (Brazil), Pavilhao "108" - Sala: 09, Av. Brasil, 4365 - Manguinhos, Rio de Janeiro - RJ, CEP: 21040-360, Brazil
<https://orcid.org/0000-0002-3678-5175>

Mallikarjun Bidarimath - Department of Pathobiological Sciences, Louisiana State University, Louisiana Animal Disease Diagnostic Laboratory (LADDL), 1043 River Rd Rm 2024, Baton Rouge, LA 70803, USA
<https://orcid.org/0000-0002-9080-8737>

Chyer Kim - Virginia State University, Petersburg, VA, USA
<http://orcid.org/0000-0002-7481-6120>

Ionel D. Bondoc - Department of Public Health, Faculty of Veterinary Medicine Iasi, University of Agricultural Sciences and Veterinary Medicine Iasi, Romania
<https://orcid.org/0000-0002-5958-7649>

Filippo Giarratana - Department of Veterinary Medicine, University of Messina, Polo Universitario dell'Annunziata, 98168 Messina, Italy
<https://orcid.org/0000-0003-0892-4884>

Abdelaziz ED-DRA - Department of Biology, Faculty of Science, Moulay Ismail University, BP. 11201 Zitoune, Meknes, Morocco
<https://orcid.org/0000-0003-3273-1767>

Eduardo Jorge Boeri - Institute of Zoonosis Luis Pasteur, Buenos Aires, Argentina
<https://orcid.org/0000-0001-8535-0306>

Liliana Aguilar-Marcelino - CENID-PARASITOLOGIA VETERINARIA, Instituto Nacional de Investigaciones Forestales Agrícolas y Pecuarias: Jiutepec, Morelos, Mexico
<https://orcid.org/0000-0002-8944-5430>

Anut Chantiratikul - Department of Agricultural Technology, Faculty of Technology, Maharakham University, Muang, Maharakham Province 44150 Thailand
<https://orcid.org/0000-0002-8313-5802>

Panagiotis E Simitzis - Department of Animal Breeding and Husbandry, Faculty of Animal Science and Aquaculture, Agricultural University of Athens, 75 Iera Odos, 11855, Athens, Greece
<http://orcid.org/0000-0002-1450-4037>

Bartosz Kieronczyk - Poznan University of Life Sciences, Poznan, Greater Poland, Poland
<https://orcid.org/0000-0001-6006-117X>

Mario Manuel Dinis Ginja - Department of Veterinary Sciences, Center for Research and Agro-Environmental and Biological Technologies, University of Trás-os-Montes and Alto Douro, Portugal
<https://orcid.org/0000-0002-0464-7771>

Nuh Kilic - Department of Surgery, Faculty of Veterinary Medicine, Adnan Menderes University, Turkey
<https://orcid.org/0000-0001-8452-161X>

Hanna Markiewicz - Milk Examination Laboratory, Kazimierz Wielki University in Bydgoszcz, Poland
<https://orcid.org/0000-0001-8225-0481>

Kai Huang - University of Texas Medical Branch at Galveston, Galveston, TX, USA
<https://orcid.org/0000-0002-3373-5594>

N. De Briyne - Federation of Veterinarians of Europe, Brussels, Belgium
<https://orcid.org/0000-0002-2348-930X>

Hasan Meydan - Akdeniz University, Faculty of Agriculture, Antalya, Turkey
<https://orcid.org/0000-0003-4681-2525>

Suleyman Cilek - Kirikkale Universitesi, Kirikkale, kirikkale, Turkey
<https://orcid.org/0000-0002-2352-649X>

Rodrigo Alberto Jerez Ebensperger - University of Zaragoza, Spain

Joao Simoes - Universidade de Tras-os-Montes e Alto Douro, Vila Real, Portugal
<http://orcid.org/0000-0002-4997-3933>

Alberto Elmi - University Of Bologna, Ozzano dell'Emilia, Bologna, Italy
<https://orcid.org/0000-0002-7827-5034>

Parag Nigam - Department of Wildlife Health Management, Wildlife Institute of India, Dehradun, India

Ali Aygun - Selcuk Universitesi, Konya, Turkey
<https://orcid.org/0000-0002-0546-3034>

Karim El-Sabrou - Poultry Production Department, Alexandria University, Alexandria, Egypt
<https://orcid.org/0000-0003-2762-2363>

Gul Ahmad - Associate Professor of Biology (Tenured), Department of Natural Sciences, School of Arts & Sciences, Peru State College, Peru, Nebraska 68321, USA.

September 2020, Vol.13 No.9, Issue DOI-www.doi.org/10.14202/vetworld.2020.9

Research (Published online: 01-09-2020)

1. An immunohistochemical study of endocrine cells in the digestive tract of *Varanus salvator* (Reptile: Varanidae)

Mahfud Mahfud, Ernawati Ernawati, Nur R. Adawiyah Mahmud, Teguh Budipitojo and Hery Wijayanto

Veterinary World, 13(9): 1737-1742

Research (Published online: 02-09-2020)

2. Listeriosis in a peri-urban area: Cultural and molecular characterization of *Listeria monocytogenes* isolated from encephalitic goats

Nagendra Nath Barman, Anjan Jyoti Nath, Sharmita Doley, Shameem Ara Begum, Parikshit Kakati, Sailendra Kumar Das, Taibur Rahman, Dipak Bhuyan, Bhaben Chandra Baishya and Susanta Goswami

Veterinary World, 13(9): 1743-1749

Research (Published online: 02-09-2020)

3. Financial and economic analyses of the impact of cattle mastitis on the profitability of Egyptian dairy farms

M. F. Azooz, Safaa A. El-Wakeel and H. M. Yousef

Veterinary World, 13(9): 1750-1759

Research (Published online: 03-09-2020)

4. Topical effect of a specific spot-on treatment made of natural ingredients in rabbits (*Oryctolagus cuniculus*) with skin problems: A pilot study

Galia Sheinberg Waisburd, Alberto Martin Cordero, Camilo Romero Núñez, Laura Miranda Contreras, Rafael Heredia Cárdenas, and Linda G. Bautista Gómez

Veterinary World, 13(9): 1760-1763

Research (Published online: 03-09-2020)

5. *Candidatus* Rickettsia colombianensi in ticks from reptiles in Córdoba, Colombia

Jorge Miranda, Lina Violet-Lozano, Samia Barrera, Salim Mattar, Santiago Monsalve-Buriticá, Juan Rodas and Verónica Contreras

Veterinary World, 13(9): 1764-1770

Research (Published online: 04-09-2020)

6. Molecular characterization of antimicrobial resistance and enterobacterial repetitive intergenic consensus-PCR as a molecular typing tool for *Salmonella* spp. isolated from poultry and humans

María Paula Herrera-Sánchez, Roy Rodríguez-Hernández and Iang Schroniltgen Rondón-Barragán

Veterinary World, 13(9): 1771-1779

Research (Published online: 04-09-2020)

7. Prophylactic impact of nano-selenium on performance, carcasses quality, and tissues' selenium concentration using reversed-phase high-performance liquid chromatography during microbial challenge in broiler chickens

Essam S. Soliman, Fadwa F. Mahmoud, Mai A. Fadel and Rania T. Hamad

Veterinary World, 13(9): 1780-1797

Research (Published online: 05-09-2020)

8. Anti-feline immunodeficiency virus reverse transcriptase properties of some medicinal and edible mushrooms

Supaphorn Seetaha, Siriluk Ratanabunyong, Lueacha Tabtimmai, Kiattawee Choowongkomon, Jatuporn Rattanasrisomporn and Khuanjarat Choengpanya

Veterinary World, 13(9): 1798-1806

Research (Published online: 05-09-2020)

9. Phylogenetic group determination and plasmid virulence gene profiles of colistin-resistant *Escherichia coli* originated from the broiler meat supply chain in Bogor, Indonesia
Irma Rahayuningtyas, Agustin Indrawati, I Wayan Teguh Wibawan, Maria Fatima Palupi and Istiyaningsih Istiyaningsih
Veterinary World, 13(9): 1807-1814

Research (Published online: 07-09-2020)

10. Hepatoprotective activity of okra (*Abelmoschus esculentus* L.) in sodium nitrite-induced hepatotoxicity
Sri Puji Astuti Wahyuningsih, Elma Sakinatus Sajidah, Baiq Naili Dewi Atika, Dwi Winarni and Manikya Pramudya
Veterinary World, 13(9): 1815-1821

Review (Published online: 09-09-2020)

11. *Mycobacterium tuberculosis* complex in wildlife: Review of current applications of antemortem and postmortem diagnosis
Yusuf Madaki Lekko, Peck Toung Ooi, Sharina Omar, Mazlina Mazlan, Siti Zubaidah Ramanan, Sabri Jasni, Faez Firdaus Abdullah Jesse and Azlan Che-Amat
Veterinary World, 13(9): 1822-1836

Review (Published online: 10-09-2020)

12. Interaction between stress hormones and phagocytic cells and its effect on the health status of dairy cows: A review
Mohammed Naif Alhussien and Ajay Kumar Dang
Veterinary World, 13(9): 1837-1848

Research (Published online: 10-09-2020)

13. Relative white blood cell counts, heterophil-to-lymphocyte ratio, and discovery of blood parasites in wild dugong (*Dugong dugon*) at Lingayan Island, Toli-toli, Indonesia
Aryani Sismin Satyaningtjas, Agik Suprayogi, Huda Shalahudin Darusman, Ardiansyah Nurdin and Desrayni Hanadhita
Veterinary World, 13(9): 1849-1853

Research (Published online: 11-09-2020)

14. Association of bovine uterine involution disturbances with serum neuropeptide concentrations
Aida Abultdinova, Isatay Jakupov, Joachim Roth, Klaus Failing, Axel Wehrend and Marlene Sickinger
Veterinary World, 13(9): 1854-1857

Research (Published online: 11-09-2020)

15. Individual and combined anti-trypanosomal effects of arteether and diminazene aceturate in the treatment of experimental *Trypanosoma brucei brucei* infection in rats
Tobias Nnia Egbe-Nwiyi, Samson Enejo Abalaka, Nuhu Abdulazeez Sani, Oremeyi Zainab Tenuche and Idoko Sunday Idoko
Veterinary World, 13(9): 1858-1862

Research (Published online: 12-09-2020)

16. Leukocyte populations and their cell adhesion molecules expression in newborn dromedary camel calves

Muaadh M. Gaashan, Abdullah I. A. Al-Mubarak and Jamal Hussen

Veterinary World, 13(9): 1863-1869

Research (Published online: 15-09-2020)

17. Chronic unpredictable stress-induced inflammation and quantitative analysis of neurons of distinct brain regions in Wistar rat model of comorbid depression

Vandana Blossom, Megha Gokul, Nayanatara Arun Kumar, Rekha D. Kini, Shyamala Nayak and K. Bhagyalakshmi

Veterinary World, 13(9): 1870-1874

Research (Published online: 15-09-2020)

18. Identification and differentiation of *Campylobacter* isolated from chicken meat using real-time polymerase chain reaction and high resolution melting analysis of *hipO* and *glyA* genes

Ika Kartika Syarifah, Hadri Latif, Chaerul Basri and Puji Rahayu

Veterinary World, 13(9): 1875-1883

Research (Published online: 16-09-2020)

19. Molecular diagnosis and biochemical studies of tick-borne diseases (anaplasmosis and babesiosis) in Aberdeen Angus Cattle in New Valley, Egypt

Nani Nasreldin, Rania M. Ewida, Hatem Hamdon and Yasser F. Elnaker

Veterinary World, 13(9): 1884-1891

Research (Published online: 16-09-2020)

20. Prevalence and antibiotic resistance of *Campylobacter coli* isolated from broiler farms in the Marrakesh Safi region, Morocco

Raja Asmai, Bouchra Karraouan, Khadija Es-Soucratti, Houda En-Nassiri, Brahim Bouchrif, Hakim Karib and Réda Triqui

Veterinary World, 13(9): 1892-1897

Research (Published online: 17-09-2020)

21. First report of PCR-based detection of *Helicobacter* species DNA in *Camelus dromedarius* in Egypt

Ahmed Youssef, Ahmed Afifi, Ayman Hamed and Mohamed Enany

Veterinary World, 13(9): 1898-1901

Research (Published online: 18-09-2020)

22. Effects of hydrolyzed yeast supplementation on growth performance, immunity, antioxidant capacity, and microbial shedding in weaning pigs

Waewaree Boontiam, Chalong Wachirapakorn and Phreerapong Phaengphairee

Veterinary World, 13(9): 1902-1909

Meeting Report (Published online: 19-09-2020)

23. Development of a practical framework for sustainable surveillance and control of ticks and tick-borne diseases in Africa

Felix Nchu, Nkululeko Nyangiwe, Dennis Muhanguzi, Jahashi Nzalawahe, Yakob Petro Nagagi, George Msalya, Natala Audu Joseph, Esther Gwae Kimaro, Margaret Mollel, Violet Temba and Difo Voukang Harouna

Veterinary World, 13(9): 1910-1921

Research (Published online: 19-09-2020)

24. Novel probiotic lactic acid bacteria isolated from indigenous fermented foods from West Sumatera, Indonesia

Harnentis Harnentis, Yetti Marlida, Yuliaty Shafan Nur, Wizna Wizna, Melia Afnida Santi, Nadia Septiani, Frederick Adzitey and Nurul Huda

Veterinary World, 13(9): 1922-1927

Review (Published online: 21-09-2020)

25. Impact of COVID-19 and associated lockdown on livestock and poultry sectors in India
Jyotsnarani Biswal, Kennady Vijayalakshmy and Habibar Rahman

Veterinary World, 13(9): 1928-1933

Research (Published online: 22-09-2020)

26. Effect of hydrogen peroxide on the oxidative burst of neutrophils in pigs and ruminants

Francesco Mosca, Abigail R. Trachtman, Jasmine Hattab, Giuseppe Marruchella and Pietro G. Tiscar

Veterinary World, 13(9): 1934-1939

Research (Published online: 22-09-2020)

27. Isolation and molecular characterization of fowl aviadenovirus associated with inclusion body hepatitis from poultry in Banten and West Java, Indonesia

Otto Sahat Martua Silaen, Sri Murtini, Joko Pamungkas and Christian Marco Hadi Nugroho

Veterinary World, 13(9): 1940-1946

Research (Published online: 23-09-2020)

28. Prevalence and antibiotic susceptibility of *Mannheimia haemolytica* and *Pasteurella multocida* isolated from ovine respiratory infection: A study from Karnataka, Southern India

Swati Sahay, Krithiga Natesan, Awadhesh Prajapati, Triveni Kalleshmurthy, Bibek Ranjan Shome, Habibur Rahman and Rajeswari Shome

Veterinary World, 13(9): 1947-1954

Research (Published online: 23-09-2020)

29. Renal ultrasonographic shear-wave elastography and urinary procollagen type III amino-terminal propeptide in chronic kidney disease dogs

Chutimon Thanaboonnipat, Saikaew Sutayatram, Chollada Buranakarl and Nan Choisunirachon

Veterinary World, 13(9): 1955-1965

Research (Published online: 23-09-2020)

30. Evaluation of the immune status of birds and domestic and companion animals for the influenza A virus in Eastern Saudi Arabia
Abdelmohsen Abdualah Alnaeem, Abdulkareem Al-Shabeb and Maged Gomaa Hemida
Veterinary World, 13(9): 1966-1969

Research (Published online: 25-09-2020)

31. Effects of respiratory disease on Kele piglets lung microbiome, assessed through 16S rRNA sequencing
Jing Zhang, Kaizhi Shi, Jing Wang, Xiong Zhang, Chunping Zhao, Chunlin Du and Linxin Zhang
Veterinary World, 13(9): 1970-1981

Research (Published online: 25-09-2020)

32. Productivity, absence of a bull and endoparasitic nematodiosis in beef cattle farms in an upland area of East Java, Indonesia
Widi Nugroho, Siska Aditya, Rahadi Swastomo and Aulanni'am Aulanni'am
Veterinary World, 13(9): 1982-1987

Research (Published online: 25-09-2020)

33. Contamination factors associated with surviving bacteria in Thai commercial raw pet foods
Suppada Kananub, Nayika Pinniam, Sitthiporn Phothitheerabut and Praphaphan Krajanglikit
Veterinary World, 13(9): 1988-1991

Research (Published online: 26-09-2020)

34. Serological study on brucellosis in captive elephants (*Elephas maximus*) and stray dogs in North Thailand
Suchanit Ngamkala, Taweepoke Angkawanish, Weerapun Nokkaew and Nikorn Thongtip
Veterinary World, 13(9): 1992-1997

Review (Published online: 26-09-2020)

35. The COVID-19: Current understanding
Shweta Tripathi and Mayukh Mani Tripathi
Veterinary World, 13(9): 1998-2005

Research (Published online: 26-09-2020)

36. Influence of pre and postpartum alpha-tocopherol supplementation on milk yield, milk quality, and udder health of Jersey crossbred cows at tropical lower Gangetic region
Amit Kumar Singh, Champak Bhakat, Tripti Kumari, Dilip Kumar Mandal, Anupam Chatterjee, Muthupalani Karunakaran and Tapas Kumar Dutta
Veterinary World, 13(9): 2006-2011

Research (Published online: 28-09-2020)

37. Evaluation of the bacteriocin produced by strain 9 lactic acid bacteria isolate for biopreservation
I Dewa Made Sukrama, Juliana Franciska and I Wayan Suardana
Veterinary World, 13(9): 2012-2019

Research (Published online: 28-09-2020)

38. The prevalence and risk factors associated with *Leptospira* in donkeys in Ngaka Modiri Molema District, North West Province, South Africa

Kibambe Kiayima Daddy, Mulunda Mwanza, James Wabwire Oguttu and Lubanza Ngoma
Veterinary World, 13(9): 2020-2027

Research (Published online: 30-09-2020)

39. *Ocimum sanctum* Linn. ethanolic extract inhibits angiogenesis in human lung adenocarcinoma (A549) cells

Hevi Wihadmadyatami, Puspa Hening, Ulayatul Kustiati, Dwi Liliek Kusindarta, Teguh Triyono and Supriatno Supriatno

Veterinary World, 13(9): 2028-2032

Research (Published online: 30-09-2020)

40. Hematology and serum biochemistry of free-range brown-throated sloths in two urban areas in Brazil


Kissia Ferreira Pereira, Ita de Oliveira e Silva, Fernanda de Fátima Rodrigues Silva, Vinicius Herold Dornelas e Silva, Carla Soraia Soares de Castro and Vanner Boere

Veterinary World, 13(9): 2033-2038

The screenshot shows the Veterinary World journal website. The main heading is "Veterinary World" with a logo. On the right, the H-Index is prominently displayed as "23" with "H Index" written below it. On the left, there is a sidebar with the following details:


- Country:** India - [SIR Ranking of India](#)
- Subject Area and Category:** [Veterinary](#) / [Veterinary \(miscellaneous\)](#)
- Publisher:** [Veterinary World](#)
- Publication type:** Journals
- ISSN:** 22310916, 09728988
- Coverage:** 2008-2020
- Scope:** Veterinary World publishes high quality papers focusing on Veterinary and Animal Science. The fields of study are bacteriology, parasitology, pathology, virology, immunology, mycology, public health, biotechnology, meat science, fish diseases, nutrition, gynecology, genetics, wildlife, laboratory animals, animal models of human infections, prion diseases and epidemiology. Studies on zoonotic and emerging infections are highly appreciated. Review articles are highly appreciated. All articles published by Veterinary World are made freely and permanently accessible online. All articles to Veterinary World are posted online immediately as they are ready for publication.

At the bottom of the sidebar, there are links for [Homepage](#) and [How to publish in this journal](#). A cookie notice at the bottom of the page reads: "This website uses cookies to ensure you get the best experience on our website" with a "Got it!" button. The Windows taskbar at the bottom shows several open applications and the system clock indicating 21:45 on 30/09/2020.



This website uses cookies to ensure you get the best experience on our website. [Got it!](#)

Taskbar: Veterinary..., Montigo, Temp, GWK SIM..., Sukrama..., BAB I DEL..., X BAB II D..., BAB II DE..., Documen..., Documen..., Evaluatio..., 21:46 30/09/2020



Veterinary World ← Show this widget in your own website

Q2 Veterinary (miscellaneous) best quartile

SJR 2019 0.5

powered by scimagojr.com

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com/journalsearch.php?q=21100201717&tip=sid&clean=0">
```

research paper publication

This website uses cookies to ensure you get the best experience on our website. [Got it!](#)

Taskbar: Veterinary..., Montigo, Temp, GWK SIM..., Sukrama..., BAB I DEL..., X BAB II D..., BAB II DE..., Documen..., Documen..., Evaluatio..., 21:46 30/09/2020

URL: https://www.googleadservices.com/pagead/aclk?sa=L&ai=CeBQEEYy0X8fNj5u4rtoP_8GFIA:ChpTjWv_yyryPyCsCnTweQASCO2...kcpInCm8sig=AOD64_05ZLngOAdBhain3Qyg-m19UN8Cg&client=ca-pub-7636113250813806&nb=0&adurl=http://ijrar.org/

Evaluation of the bacteriocin produced by strain 9 lactic acid bacteria isolate for biopreservation

I Dewa Made Sukrama¹, Juliana Franciska² and I Wayan Suardana³ 

1. Department of Clinical Microbiology, Faculty of Medicine, Udayana University, Jl. PB. Sudirman Denpasar-Bali, 80234, Indonesia; 2. Department of Preventive Veterinary Medicine, Graduate Student of Veterinary Medicine, Faculty of Veterinary Medicine, Udayana University. Jl. PB. Sudirman Denpasar-Bali, 80232, Indonesia; 3. Department of Preventive Veterinary Medicine, Laboratory of Veterinary Public Health, Faculty of Veterinary Medicine, Udayana University, Jl. PB. Sudirman Denpasar-Bali, 80232, Indonesia.

Corresponding author: I Wayan Suardana, e-mail: wayan_suardana@unud.ac.id

Co-authors: IDMS: dewa_sukrama@unud.ac.id, JF: ulieffrancis@gmail.com

Received: 05-04-2020, **Accepted:** 03-08-2020, **Published online:** 28-09-2020

doi: www.doi.org/10.14202/vetworld.2020.2012-2019 **How to cite this article:** Sukrama IDM, Franciska J, Suardana IW (2020) Evaluation of the bacteriocin produced by strain 9 lactic acid bacteria isolate for biopreservation, *Veterinary World*, 13(9): 2012-2019.

Abstract

Aim: This study aimed to determine the effect of the bacteriocin produced by strain 9 lactic acid bacteria (LAB) isolate on the biopreservation of beef.

Materials and Methods: The strain 9 LAB isolate was identified conventionally by culturing with de Man, Rogosa, and Sharpe broth medium followed by Gram staining and catalase testing. The molecular confirmation of the isolate involved analyzing the 16S rRNA gene with specific primers, that is, B27F (5-AGAGTTTGATCCTGGCTCAG-3) and U1492R (5-GGTTACCTGTGTTACGACTT-3). Then, the isolate was centrifuged to evaluate the bacteriocin production, and the effect of the biopreservative activity in beef was evaluated by measuring the NH₃ produced with the Eber test and the organoleptic acceptance from expert panels.

Results: This study confirmed that the strain 9 LAB isolate was a strain of *Pediococcus pentosaceus*, and the bacteriocin product showed biopreservative potential. The biopreservative potential was characterized by a significant decline in the production of NH₃ and the panel's acceptance of the texture and tenderness of the beef, compared with the control, after 10 days of constant treatment.

Conclusion: This study highlighted the high biopreservative potency of pediocin produced by *P. pentosaceus* strain 9. This was noted by the production of NH₃ and the modifications in texture and tenderness.

Keywords: bacteriocin, beef, biopreservative, lactic acid bacteria.

Introduction

Lactic acid bacteria (LAB) are a group of Gram-positive, non-sporulating, anaerobic, or facultative aerobic rods or cocci. These organisms are known to produce lactic acid as a major product of carbohydrate fermentation [1]. Furthermore, these organisms are classified into four genera based on the mode of glucose fermentation, temperature range at which they grow, sugar utilization patterns, and cellular morphology. These genera include *Lactobacillus*, *Leuconostoc*, *Pediococcus*, and *Streptococcus*, although the recent use of molecular methods has identified more groups [2,3].

Based on their ability to inhibit the growth of various Gram-negative/-positive bacteria, LAB have thus been adopted as natural starters in food fermentations. These organisms also protect food from spoilage and pathogenic microorganisms through

the production of organic acids, hydrogen peroxide, diacetyl, and bacteriocins [4,5].

Specifically, bacteriocins are ribosomally synthesized small proteins known to inhibit spoilage and the growth of pathogenic bacteria in foods [6]. These characteristics have recently been of interest in research due to their possibility of replacing chemical preservation [7]. Furthermore, consumers are currently fascinated by the use of natural food preservation techniques [6,8].

Based on the structural, physicochemical, and molecular properties, bacteriocin peptides from LAB are subdivided into three major classes: Class I, which includes lantibiotics, is small in size (<5 kDa) and contains lanthionine and β -methylanthionine; Class II, which is also small (<10 kDa) and heat-stable, but non-lanthionine; and Class III, which is characterized by heat-sensitive large molecules [7].

Bacteriocin-producing species have been identified among all the genera of LAB, including *Lactobacillus*, *Carnobacterium*, *Pediococcus*, and *Enterococcus* [6,7]; a large number of these species have been isolated and characterized, and some serve as food preservatives. Some important examples include nisin, diplococcin, acidophilin, bulgarican, helveticins, lactacins, and plantaricins [9].

Copyright: Sukrama, et al. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

There have been several studies in Indonesia related to the use of LAB strains as probiotics due to the presence of inherent functional properties that contribute positively toward health. In 2012, Sujaya *et al.* [10] evaluated the application of exopolysaccharides produced by *Lactobacillus* spp. that were isolated from Sumbawa Mare's Milk as probiotics. Subsequently, Nuraida [11] reported on the effect of some probiotic strains on hypocholesterolemia, immune system stimulation, and the prevention of diarrhea in animals [11]. However, prior studies on the isolation, characterization, and application of bacteriocin are limited. According to a previous study conducted by Suardana in 2015, strain 9 LAB isolate was one of the LAB culture stocks isolated from the gastric juice of Bali cattle. Its activity was evidenced by the killing zones produced in *Staphylococcus aureus* bacterial indicator, at 0.77-1.26 cm, although a negative result was obtained in the treatment with *Escherichia coli*.

This study aimed to determine the effect of the bacteriocin produced by a strain 9 LAB isolate on the biopreservation of beef.

Materials and Methods

Ethical approval

The approval from the Institutional Animal Ethics Committee to carry out this study was not required as no invasive technique was used

Isolate cultivation

The strain 9 LAB isolate was grown in de Man, Rogosa, and Sharpe (MRS) broth medium (Oxoid CM0359.UK Ltd.) under anaerobic conditions with the addition of an Oxoid gas generating kit (Oxoid BR 38) in an anaerobic jar for further incubation at 37°C for 24 h. The isolate was then Gram-stained and underwent a catalase test with the addition of 10 µL of isolate and one drop of 10% H₂O₂, before performing molecular confirmation [12].

Study period and location

The study was conducted from July to September 2016 at the Laboratory of Veterinary Public Health, Faculty of Veterinary Medicine, Udayana University, Denpasar-Bali.

DNA isolation

The genomic DNA of strain 9 LAB was extracted using a DNA kit (Geneaid, GBB100. Biotech Ltd. Taiwan) according to the manufacturer's instructions. The culture was centrifuged at 16,000×g for 1 min. The supernatant was then discarded, followed by adding the pellet to 200 µL of Gram-positive buffer, which had previously been filled with lysozyme enzymes (0.8 mg/200 µL). The mixed culture was then incubated at 30°C for 30 min followed by the addition of 20 µL of K proteins and incubation at 60°C for 10 min. Subsequently, 200 µL of GB buffer was added before another round of incubation at 70°C for 10 min. Then, 200 µL of absolute ethanol

was introduced and centrifuged at 16,000×g for 2 min. The container tube was replaced, 400 µL of W1 buffer was added to the column, and it was centrifuged at 16,000×g for 30 s. The container tube was swapped with a new tube that had previously contained ethanol; 600 µL of wash buffer was added, and the tube was centrifuged at 16,000×g for 30 s. The column was centrifuged again for 3 min at 16,000×g until a dry column matrix was obtained. At the termination stage, 100 µL of elution buffer (TE buffer) was added and centrifuged at 16,000×g for 30 s. This was performed to dissolve the DNA and capture fragments ready for use [12].

16S rRNA gene analysis

The *16S rRNA* gene of the strain 9 LAB isolate was amplified using B27F universal primers arranged as 5-AGAGTTTGATCCTGGCTCAG-3 and U1492R as 5-GGTTACCTTGTTACGACTT-3, according to the previous procedure, with minimal modifications [12-14]. A total of 1 µL of genomic DNA were introduced to the polymerase chain reaction (PCR) tube, in addition to 25 µL of DreamTaq Green (Thermo Fisher Scientific, K1081.California US), 1 µL each of forward and reverse primers at a concentration of 20 pmol, and sterile water to adjust to a final volume of 40 µL. Therefore, the PCR reaction was programmed using single-step denaturation for 7 min, characterized by 35 cycles of exposure to a denaturation temperature of 94°C for 1 min, an annealing temperature of 56°C for 1 min, and an elongation stage (extension) at 72°C for 1.5 min. Subsequently, the PCR was evaluated using an additional extension temperature of 72°C for 5 min.

Electrophoresis with 1.5% agarose

The PCR products were analyzed by electrophoresis on 1.5% agarose gel (Sigma 5093-100G) using ethidium bromide staining (Merck, E1510. Germany). This involved the transfer of gel to the electrophoresis tank containing 1× Tris-Borate-EDTA (TBE) buffer, followed by the addition of 5 µL of PCR product and 1 µL of loading buffer, which were then mixed and added to the existing gel wells. Then, electrophoresis was performed at 100 volts for 30 min and viewed under ultraviolet light (transilluminator). The presence of a 1502 bp PCR product indicates a positive outcome [14].

Sequencing and phylogenetic analysis

The *16S rRNA* gene of the strain 9 LAB isolate was sequenced using a genetic analyzer (ABI Prism 3130 and 3130×1 Genetic Analyzer) at the 1st BASE DNA Sequencing, PT (Genetika Science, Jakarta). This procedure involved primers similar to those used for the PCR, and the outcomes were then manually edited to exclude the PCR primer binding sites, using MEGA version 5.2 software (The Pennsylvania State University) This modified result was automatically compared against the sequences of LAB available in databanks (www.ncbi.nlm.nih.gov) using the BLAST

program on MEGA version 5.2 software. Therefore, the data obtained were subsequently used in phylogenetic tree construction by means of the neighbor-joining algorithm method [15,16].

Isolation and purification of bacteriocin

Five microliters of the strain 9 LAB isolate were plated on the MRS broth medium (Oxoid CM0359. UK) under anaerobic conditions by adding the Oxoid gas generating kit (Oxoid BR 38. UK) in an anaerobic jar for further incubation at 37°C for 24 h. This mixture was then centrifuged at 12,000×g for 10 min at 4°C and filtered through a 0.22 µm sieve to separate the cell-free supernatants, which was considered the crude bacteriocin. Furthermore, 12 N NaOH was used to adjust the pH to 7.0, and ammonium sulfate (Fluka, Netherlands) was added to obtain 60% saturation, stirred overnight, and subjected to a round of centrifugation at 14,000×g for 1 h at 4°C. Subsequently, the pellets were dissolved with 20 mM potassium phosphate buffer, and the ready semipurified bacteriocin was collected for further evaluation [17,18].

Application of bacteriocin as a biopreservative

The application of bacteriocin as a biopreservative was evaluated by marinating beef in a bacteriocin solution with a concentration of 500 µL/mL (v/v). This study used 5 g beef samples that were 5×5×2 cm in size and were entirely marinated in 0.85% NaCl solution to align the initial microbial contamination. The samples were divided into three treatment groups as follows: Group I was the control, which was marinated without bacteriocin; Group II was marinated in 500 µL/mL (v/v) bacteriocin for a 5 min treatment; and Group III used similar materials for 10 min. Subsequently, all samples were wrapped in plastic clips and stored in the refrigerator at 5°C, and each treatment was replicated 3 times. Therefore, continuous operation was performed on days 0, 2, 4, 6, 8, and 10. The change in beef quality among the treatments in each observation was measured according to the meat quality parameters in this study.

Meat quality parameters

The meat quality parameters observed included spoilage, which was evaluated with the Eber test, and the texture and tenderness, which were evaluated by expert panels. The Eber test comprises a combination of 1 part 37% HCl (Merck 1.00317.2500.Germany) in three parts 96% alcohol (BD.SKU74000CPC) and 1 part 74.12 g/mol ether (Merck 1009301000.Germany). The results were declared negative in the absence of any white clouds around the meat, positive 1 (+) if a small quantity of white clouds formed (after 10 min), positive 2 (++) if quite a lot of white clouds formed (within 5-10 min), and positive 3 (+++) if large white clouds formed (within <5 min). Based on the formation of NH₄Cl with the Eber test, 0 is indicative of a negative (-) outcome, whereas 1 (+), 2 (++) and 3 (+++) denote positive outcomes.

Furthermore, the texture and tenderness were evaluated by five trained panelists (female students in their eighth semester) from the Faculty of Veterinary Medicine; the results were scored as 3, 2, and 1 for fresh, mushy, and watery meat, respectively. These values were then summed, and an average calculated, which was subsequently statistically analyzed with a one-way ANOVA, followed by Duncan's multiple range test. Next, a relationship between treatments and storage period was identified based on the formation of NH₄Cl gas and the texture and tenderness, and the results were further subjected to a regression correlation analysis using SPSS 18 software.

Results

Isolate cultivation

The strain 9 LAB isolate cultivation was characterized by luxuriant growth on the MRS broth medium in an anaerobic atmosphere, Gram-positive results, and a negative catalase test. The results of this study confirmed the strain 9 isolate to be an LAB, so we could proceed to the next analysis.

16S rRNA gene analysis

The 16S rRNA gene of the strain 9 LAB isolate was amplified successfully using the B27F and U1492R primers previously characterized by the clear 1502 bp PCR product. Therefore, the sequencing result was aligned against several LAB data from GenBank, which identified the bacteria as *Pediococcus pentosaceus*, of the group *P. pentosaceus* (AB682664) with a bootstrap value of 100 (Figure-1).

The assembly of the strain 9 LAB isolate with *P. pentosaceus* (AB682664) in one group was confirmed by the data in Table-1, which shows the similarity in the nucleotide sequences of both strains. However, the intrinsic characteristics are quite different from *Oenococcus oeni* (AB596939), with 164 different nucleotides per 1000 alignment. Based on this comparison concept, samples with nucleotide similarities >90% or differences between 1 and 1.5% (14-22 bp) are categorized as the same species [13,14,19].

Application of bacteriocin as a biopreservative

The Eber test is one of the methods used to determine the spoilage of meat and its derivatives and is shown by the production of NH₃, which is produced from the biochemical activity of detrimental microorganisms. The gas released is recognized by the formation of white clouds through a reaction between NH₃ and HCl around the test sample.

The Eber test showed the release of NH₃ from the 6th to the 10th day of observing the control group and the groups treated with bacteriocin and marinated for 5 and 10 min. The production of NH₃ was reduced in the treatment groups based on the relatively smaller amount of white clouds formed, and the production time was also longer. Furthermore, the data were converted into 0 for negative results (-), 1 for (+), 2 for (++) and 3 for (+++), after being transformed by $\sqrt{0.5}$.

Table-1: Pyelogram of strain 9 LAB isolate against other samples in GenBank. The phyelogram was constructed based on neighbor-joining algorithm.

	Lactic acid bacteria strain 9	Aerococcus viridans (LC487893)	Aerococcus viridans (AB680271)	Enterococcus faecalis (JX2421472)	Enterococcus faecalis (JX2421472)	Lactobacillus rhamnosus (D16552)	Lactococcus lactis (AB100796)	Oenococcus oeni (AB596939)	Pediococcus siamensis (AB258357)	Pediococcus pentosaceus (AB682664)	Streptococcus salivarius (AB608747)	Streptococcus pneumoniae (AB002522)	Vagococcus salmoninarum (X54272)	Vagococcus fluvialis (LC487872)
Lactic acid bacteria strain 9														
Aerococcus viridans (LC487893)	0.093													
Aerococcus viridans	0.092	0.001												
Enterococcus faecalis (JX2421472)	0.085	0.060	0.059											
Lactobacillus rhamnosus (D16552)	0.043	0.072	0.071	0.066										
Lactococcus lactis (AB100796)	0.134	0.133	0.131	0.118	0.137									
Leuconostoc lactis (A1970316)	0.132	0.136	0.134	0.120	0.130	0.156								
Oenococcus oeni (AB596939)	0.164	0.177	0.175	0.172	0.171	0.162	0.124							
Pediococcus siamensis (AB258357)	0.055	0.114	0.113	0.099	0.067	0.165	0.160	0.191						
Pediococcus pentosaceus (AB682664)	0.000	0.093	0.092	0.085	0.043	0.134	0.164	0.055						
Streptococcus salivarius (AB608747)	0.126	0.127	0.125	0.101	0.119	0.072	0.165	0.158	0.126					
Streptococcus pneumoniae (AB002522)	0.136	0.124	0.122	0.115	0.128	0.087	0.146	0.149	0.136	0.044				
Vagococcus salmoninarum (X54272)	0.092	0.066	0.064	0.046	0.080	0.114	0.146	0.188	0.092	0.112	0.122			
Vagococcus fluvialis (LC487872)	0.078	0.059	0.058	0.034	0.070	0.114	0.131	0.180	0.078	0.108	0.117	0.108	0.117	0.030




Figure-1: Phylogenetic tree constructed using the neighbor-joining algorithm of 16S rRNA gene nucleotides sequence. The number in each phylogram branch indicates a bootstrap value (%) by 1000-replication multiple, and scale shows two per 100 nucleotide substitutions.

Figure-2 shows that there was a significant difference ($p < 0.05$) in the amount of NH_4 formed in the group treated for 5 min compared with the control and with the group treated for a duration of 10 min. Furthermore, the regression analyses of the storage period for NH_4 gas formation for the control and 5 and 10 min bacteriocin treatments in Figure-3 showed regression curves of $Y = 0.056x$ (correlation coefficient $[R^2] = 0.863$); $Y = 0.013x$ ($R^2 = 0.865$); and $Y = -0.064x$ ($R^2 = 0.809$), respectively.

Figure-3 shows a strong correlation between the storage period and the formation of NH_4 gas. This is evidenced by the increase in the production of NH_4Cl by the control compared with the 5 and 10 min bacteriocin treatments. In addition, the graphics indicate a significant correlation between the production of NH_4Cl and the storage period.

Moreover, the organoleptic criteria of texture and tenderness were used to determine meat quality, and the result was congruent with the Eber test. These parameters are strongly associated with the amount of muscular connective tissues and have been selected as the most important qualitative attributes by the average consumer [20,21]. In addition, texture and tenderness, which were classified as fresh, mushy, and watery meat in this study, were closely associated with the spoilage process. Figure-4 shows the results of the panel tests on the effect of beef immersed in bacteriocin for 5 and 10 min compared with the control on the properties of beef's texture and tenderness.

Figure-4 indicates that there was a significant difference ($p < 0.05$) between the control experiment and the beef immersed in bacteriocin for 5 and 10 min. Figure-5 shows the regression correlation analysis, which displays a close correlation between the period of beef immersion and the storage phase. Furthermore, the results on texture and tenderness for




Figure-2: The properties of NH_4 produced after the reaction of NH_3 released during meat spoilage and the HCl substrate from Eber test between control and treatment groups. The table was construct base on Duncan's multiple range test, after conversion to $\sqrt{0.5}$.

the control and treatment with bacteriocin for 5 and 10 min were expressed with regression curves of $Y = 3$ ($R^2 = \text{N/A}$); $Y = -0.22X^2 + 0.151X + 2.892$ ($R^2 = 0.785$); and $Y = -0.022X^2 + 0.023X + 3.035$ ($R^2 = 0.929$), respectively.

Figure-5 shows higher scores for texture and tenderness in the beef treated with bacteriocin than the control, which showed a resilient decline up to day 10 of observation.

Discussion

The low carbohydrate content of beef (1% or less) indicates a higher tendency for spoilage by organisms that utilize free amino acid content as a source of energy. Furthermore, the process leads to the production of ammonia, hydrogen sulfide, indole, skatole, and amines [22], although storage at refrigerator temperatures significantly reduces the growth rate of all genera. However, some organisms are resistant in this regulated environment and have been identified in fresh meats. These include *Micrococcus*, *Streptococcus*, *Escherichia*, and *Klebsiella* which




Figure-3: Quadratic curves of NH_4 gas formation on the Eber test, resulting from the interaction between NH_3 released during spoilage and HCl substrate, for beef immersed in bacteriocin for 5 and 10 minutes, and the control.




Figure-4: The properties of beef's texture and tenderness between control and the treatments with bacteriocin. The table was construct base on Duncan's multiple range test after conversion of descriptive data.

may be eliminated in samples treated with antibiotics or other inhibitors [22].

The strain 9 LAB isolate and the bacteriocin produced in this research were confirmed to be *P. pentosaceus* and pediocin, respectively. Based on different sources, there has been a general increase in the application of bacteriocin-producing strains in the health and food industry [23]. This is evidenced by several related studies, including the study by Cortés *et al.* [24] which showed maximum bacteriocin production by *P. pentosaceus* at 37°C that occurred predominantly at the end of the exponential growth phase, followed by a stabilized yield rate, during the 24 h incubation period. Therefore, pediocin was confirmed to have potential use as a food biopreservative.

Other studies have identified pediocin A as a bacteriocin produced by *P. pentosaceus*; it is grouped as a Class II bacteriocin, with a molecular weight of 2.7 kDa. This product was generally sensitive to proteolytic enzymes and heat (10 min at 100°C). Meanwhile, pediocin F, which is produced by *Pediococcus acidilactici* and is placed in a similar category, has a weight of 4.5 kDa and is known to be sensitive toward proteolytic enzymes, is heat and organic solvent resistant, and is active under a wide range of pH levels [6]. Several forms of pediocin-producing *Pediococcus* spp. have also been isolated and

evaluated, and despite their structural similarities, the molecular weights always varied, alongside their spectrum of antimicrobial activity. Furthermore, the bactericidal activity against Gram-positive food spoiling and pathogenic bacteria increases the product importance in the class of biopreservatives [25].

Meat and its products have been considered excellent growth media for a variety of microflora (bacteria, yeasts, and molds), including some pathogens. This phenomenon makes the intestinal tract and the skin of animals a major source of microorganisms, although the microflora composition depends on various factors: (a) Preslaughter husbandry practices (free range vs. intensive rearing); (b) age of the animal at the time of slaughtering; (c) handling during slaughtering, evisceration, and processing; (d) temperature controls during slaughtering, processing, and distribution; (e) preservation methods; (f) type of packaging; and (g) handling and storage by the consumer [26]. In addition, the pediocin produced by the strain 9 LAB isolate was proven to be a biopreservative, which was indicated by its abilities to inhibit the formation of NH_4Cl in the Eber test on meat treated with pediocin. This assumption is supported by Lawrie's theory [21], which states that the process of decay caused by spoilage organisms in meat results from the use of free amino acids as an additional energy source. The reactions produce hydrogen, carbon dioxide, and ammonia. Furthermore, reduced NH_4 formation in beef treated with bacteriocin occurs due to reduced NH_3 release, which is a result of the antibacterial activity of pediocin killing many of the spoilage microorganisms. In general, pediocin inhibits bacteria through three mechanisms: (1) Creating work pores in the cytoplasm membrane of cell targets, (2) destroying the intracellular pH of the bacteria target, and (3) inhibiting the proton motive force for energy production in the bacterial cell. Pediocin was linked to cell targets through active sites such as the negative charges of phospholipid groups in the cell wall. The previous studies have shown a relationship between the amount of lipoteichoic acid in the cell wall of Gram-positive bacteria (G+) and increased pediocin activity against G+ bacteria. There




Figure-5: Quadratic curves of beef texture and tenderness between the sample immersed in bacteriocin and the control on storage.

is higher pediocin activity when the acid concentration is increased in the walls of bacterial cells [27,28]. This result agrees with the previous statement, in which pediocin was evaluated as low-molecular-weight (2.7-17 KDa) cationic molecules. Pediocin is composed of a hydrophilic N-terminal part, which contains the pediocin box (YGNGV) motif, and a hydrophobic or amphiphilic C-terminal variable part, often produced by some *Pediococcus* bacteria. Furthermore, pediocin has inhibition properties against several sensitive bacterial cells, resulting from the ability to act on the cytoplasmic membrane during pore formation and through the absorption of amino acids in the phospholipids of the target cell cytoplasmic membrane [29]. Its effectiveness of inhibiting food spoilage has also been identified against some species of *Enterococcus*, *Listeria*, *Leuconostoc*, and *Staphylococcus* [30].

Moreover, the previous reports have defined meat spoilage as a change in quality, leading to undesirable characteristics, making it unfit for consumption. This includes the development of off-flavors, off-odors, and often the formation of slime, resulting from the breakdown of valuable contents (fat, protein, and carbohydrates). In addition, microbial spoilage causes sour tasting, discoloration, gas production, pH changes, slime formation, degradation of structural components, and appearance modification [31]. The most important qualitative attributes of eating are texture and tenderness, which basically depend on three main components: (1) The degree of sarcomere contraction or "length," (2) the extent of structural myofibrillar protein integrity/degradation (proteolysis), and (3) the connective tissue content/composition [32]. Conversely, some strains of the genera *Pseudomonas* and *Achromobacter* grow at refrigerator temperatures and are often able to degrade gelatin, with poor decomposition activity against natural proteins, including casein and purified myofibrillar proteins. However, the physical changes produced tend to be more apparent compared with the chemical effects, as observed in terms of color, odor, flavor, and

tenderness [33]. Moreover, there is a high tendency to apply bacteriocin-producing LAB strains in biopreservation, evidenced by the higher texture and tenderness scores for meat immersed in pediocin than the control after 10 days. This outcome confirmed the spoilage inhibition characteristics of bacteriocin, which agrees with findings from a study by de Costa [34], who used LAB as a biological preservative on meat. Thus, using LAB are an important approach to maintain the microbiological quality and safety of meat and its products to extend food shelf-life and foster consumer acceptance. One such method involves using bacteriocin as a biopreservative, such as pediocin-producing *P. pentosaceus* strain 9.

Conclusion

The biopreservative potency of pediocin produced by *P. pentosaceus* strain 9 on beef is confirmed to be feasible. This outcome is evidenced by the production of NH_3 and the evaluation of texture and tenderness. There is a need for subsequent studies to perform a more complete test on the aspect of potency.

Authors' Contributions

IDMS, JF, and IWS created and designed the experiment. JF and IWS performed PCR. IDMS, JF, and IWS drafted and revised the manuscript. All authors have read and approved the final manuscript.

Acknowledgments

The authors are grateful to the Institute for Research and Community Services, Udayana University, for funding this study through the Udayana Invention Grant, in accordance with the Assignment Agreement Implementation Research No. 641-116/UN 14.2/PNL.01.03.00/2016. Date June 15, 2016.

Competing Interests

The authors declare that they have no competing interests.

Publisher's Note

Veterinary World remains neutral with regard to jurisdictional claims in published institutional affiliation.

References

- Hayek, S.A. and Ibrahim, S. (2013) Current limitations and challenges with lactic acid *Bacteria*: A review. *Food Nutr. Sci.*, 4(11): 73-87.
- Von Wright, A. and Axelsson, L. (2012) Lactic acid *Bacteria*: An introduction. In: Lahtinen, S., Ouwehand, A.C., Salminen, S. and von Wright, A., editors. *Lactic acid Bacteria: Microbiological and Functional Aspects*. Taylor & Francis Group LLC, CRC Press, Boca Raton. p1-16.
- Salminen, S. (2012) *Lactic acid Bacteria: Microbiological and Functional Aspects*. 4th ed. CRC Press, Taylor & Francis, Boca Raton. p777.
- Messens, W. and De, V.L. (2002) Inhibitory substances produced by lactobacilli isolated from sourdoughs--a review. *Int. J. Food Microbiol.*, 72(1-2): 31-43.
- De Vuyst, L. and Leroy, F. (2007) Bacteriocins from lactic acid *Bacteria*: Production, purification, and food applications. *J. Mol. Microbiol. Biotechnol.*, 13(4): 194-199.
- Parada, J.L., Caron, C.R., Medeiros, A.B.P. and Soccol, C.R. (2007) Bacteriocins from lactic acid *Bacteria*: Purification, properties and use as biopreservatives. *Braz. Arch. Biol. Technol.*, 50(3): 521-542.
- Perez, R.H., Zendo, T. and Sonomoto, K. (2014) Novel bacteriocins from lactic acid *Bacteria* (LAB): Various structures and applications. *Microb. Cell. Fact.*, 13(1): S3.
- Quinto, E.J., Jiménez, P., Caro, I., Tejero, J., Mateo, J. and Girbés, T. (2014) Probiotic lactic acid *Bacteria*: A review. *Food Nutr. Sci.*, 5(18): 1765-1775.
- Mokoena, M.P. (2017) Lactic acid bacteria and their bacteriocins: Classification, biosynthesis and applications against uropathogens: A mini-review. *Molecules*, 22(8): 1255.
- Sujaya, I.N., Aryantini, N.P.D., Nursini, N.W., Cakrawati, C.I.D., Juliasari, N.L.M., Dwipayanti, N.M.U. and Ramona, Y. (2012) Exopolysaccharides from *Lactobacillus* sp. isolated from Sumbawa mare's milk and its potential application as prebiotics. *J. Vet.*, 13(2): 136-144.
- Nuraida, L. (2015) A review: Health promoting lactic acid *Bacteria* in traditional Indonesian fermented foods. *Food Sci. Hum. Wellness*, 4(2): 47-55.
- Sukrama, I.D.M., Pinatih, K.J.P. and Suardana, I.W. (2017) Molecular analysis of lactic acid *Bacteria* isolate Sr2 from Bali cattle rumen. *J. Glob. Pharma Technol.*, 5(9): 44-49.
- Suardana, I.W. (2014) Erratum to analysis of nucleotide sequences of the 16S rRNA gene of novel *Escherichia coli* strains isolated from feces of human and Bali cattle. *J. Nucleic. Acids*, 2014(12): Article ID 412942.
- Suardana, I.W. (2014) Analysis of nucleotide sequences of the 16S rRNA gene of novel *Escherichia coli* strains isolated from feces of human and Bali cattle. *J. Nucleic. Acids.*, 2014(9): Article ID 475754.
- Saitou, N. and Nei, M. (1987) The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.*, 4(4): 406-425.
- Tamura, K., Dudley, J., Nei, M. and Kumar, S. (2007) MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol. Biol. Evol.*, 24(8): 1596-1599.
- Mandal, V., Sen, S.K. and Mandal, N.C. (2011) Isolation and characterization of pediocin NV 5 producing *Pediococcus acidilactici* LAB 5 from vacuum-packed fermented meat product. *Indian J. Microbiol.*, 51(1): 22-29.
- Muyanja, C.M., Narvhus, J.A., Treimo, J. and Langsrud, T. (2003) Isolation, characterisation and identification of lactic acid bacteria from Bushera: A Ugandan traditional fermented beverage. *Int. J. Food Microbiol.*, 80(3): 201-210.
- Janda, J.M. and Abbott, S.L. (2007) 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: Pluses, perils, and pitfalls. *J. Clin. Microbiol.*, 45(9): 2761-2764.
- Mennecke, B.E., Townsend, A.M., Hayes, D.J. and Lonergan, S.M. (2007) A study of the factors that influence consumer attitudes toward beef products using the conjoint market analysis tool. *J. Anim. Sci.*, 85(10): 2639-2659.
- Lawrie, R.A. (2017) *Lawrie's Meat Science*. 8th ed. Woodhead Publishing, USA. p730.
- Jay, J.M., Loessner, M.J. and Golden, D.A. (2005) *Modern Food Microbiology*. 7th ed. D.R. Heldman, Springer, USA.
- Balandin, S.V., Sheremeteva, E.V. and Ovchinnikova, T.V. (2019) Pediocin-like antimicrobial peptides of *Bacteria*. *Biochemistry (Mosc)*, 84(5): 464-478.
- Cortés, G., Suarez, H., Buitrago, G., Nero, L.A. and Todorov, S.D. (2018) Enhanced bacteriocin production by *Pediococcus pentosaceus* 147 in co-culture with *Lactobacillus plantarum* LE27 on cheese whey broth. *Front. Microbiol.*, 9(12): 2952.
- Papagianni, M. and Anastasiadou, S. (2009) Pediocins: The bacteriocins of pediococci. Sources, production, properties and applications. *Microb. Cell. Fact.*, 8(1): 3.
- Cervený, J., Meyer, J.D. and Hall, P.A. (2009) Microbiological spoilage of meat and poultry products in compendium of the microbiological spoilage, of foods and beverages. In: Sperber, W.H. and Doyle, M.P., editors. *Food Microbiology and Food Safety*. Springer Science and Business Media, NY, USA. p69-868.
- Perez, R.H., Perez, M.T.M. and Elegado, F.B. (2015) Bacteriocins from lactic acid *Bacteria*: A review of biosynthesis, mode of action, fermentative production, uses, and prospects. *Phil. Sci. Tech.*, 8(2): 61-67.
- Wang, J., Li, L., Zhao, X. and Zhou, Z. (2015) Partial characteristics and antimicrobial mode of pediocin produced by *Pediococcus acidilactici* PA003. *Ann. Microbiol.*, 65(3): 1753-1762.
- Niamah, A.K. (2018) Structure, mode of action and application of pediocin natural antimicrobial food preservative: A review. *Basrah. J. Agric. Sci.*, 31(1): 59-69.
- Mandal, V., Sen, S.K. and Mandal, N.C. (2010) Assessment of antibacterial activities of pediocin produced by *Pediococcus acidilactici* Lab 5. *J. Food Saf.*, 30(3): 635-651.
- Dave, D. and Ghaly, A.E. (2011) Meat spoilage mechanisms and preservation techniques: A critical review. *Am. J. Agric. Biol. Sci.*, 6(4): 486-510.
- Koohmaraie, M., Kent, M.P., Shackelford, S.D., Veiseth, E. and Wheeler, T.L. (2002) Meat tenderness and muscle growth: Is there any relationship? *Meat Sci.*, 62(3): 345-352.
- Aberle, E.D., Forrest, J.C., Gerrard, D.E. and Mills, E.W. (2001) *Principles of Meat Sciences*. 4th ed. Kendall/Hunt Publishing Company, Dubuque, Iowa. p354.
- da Costa, R.J., Voloski, F.L.S., Mondadori, R.G., Duval, E.H. and Fiorentini, A.M. (2019) Preservation of meat products with bacteriocins produced by lactic acid bacteria isolated from meat. *J. Food Qual.*, 2019(1): 12.
