

ANTIBACTERIAL ACTIVITIES OF *ANREDERA CORDIFOLIA* (TEN.) V. STEENIS LEAVES EXTRACTS AND FRACTIONS

NI PUTU EKA LELIQIA^{1,2*}, ELIN YULINAH SUKANDAR¹, IRDA FIDRIANNY³

¹Department of Pharmacology-Clinical Pharmacy, School of Pharmacy, Bandung Institute of Technology, Indonesia. ²Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Udayana University, Indonesia. ³Department of Pharmaceutical Biology, School of Pharmacy, Bandung Institute of Technology, Indonesia. Email: leliqia@gmail.com

Received: 22 July 2017, Revised and Accepted: 25 August 2017

ABSTRACT

Objective: This study aims to determine antibacterial activity of *Anredera cordifolia* leaves extracts and fractions.

Methods: Crude drug was extracted using two methods. First method was extraction by reflux using ethanol 96% and then fractionated by liquid-liquid extraction using n-hexane and ethyl acetate. Second method was gradually extraction by reflux using n-hexane, ethyl acetate and ethanol 96%, respectively. Phytochemical screening was applied to all extracts and fractions, followed by thin-layer chromatography using ursolic acid, oleanolic acid (OA), apigetrin, and rutin as reference substances. A two-fold serial microdilution method was used to determine minimum inhibitory concentration (MIC) against *Staphylococcus aureus* (ATCC 6538), methicillin-susceptible *S. aureus*, methicillin resistant *S. aureus* (MRSA), *Bacillus subtilis* (ATCC 6633), *Bacillus cereus* (ATCC 11778), *Pseudomonas aeruginosa* (ATCC 9027), *Escherichia coli* (ATCC 8939), *E. coli* H7 (O156), and ESBL *E. coli*. Bacteriostatic and bactericidal activities were determined using minimum bactericidal concentration/MIC ratio.

Results: The ethanolic extract, n-hexane and ethyl acetate fractions of *A. cordifolia* from the first method had antibacterial activity against *S. aureus*, MRSA, *B. subtilis*, and *B. cereus* (MIC 256-512 µg/ml). However, n-hexane and ethyl acetate extract from the second method had broad spectrum of antibacterial activity, which could inhibit the growth of *S. aureus*, MRSA, *B. subtilis*, *P. aeruginosa*, and *E. coli* (MIC 256-512 µg/ml). Extracts and fractions showed bacteriostatic and bactericidal activities, but n-hexane extract has most bactericidal activity. Furthermore, steroid/triterpenoid, ursolic, and OA were found in this extract.

Conclusion: The n-hexane extract from the second method showed the highest antibacterial activity.

Keywords: *Anredera cordifolia*, Extracts, Fractions, Antibacterial.

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2017.v10i12.21503>

INTRODUCTION

Bacterial infections remain as the main leading cause of death, particularly in developing countries [1]. Infection treatment with antibacterial has reduced the morbidity and improved patient's survival with bacterial infections. However, in many cases, the increasing prevalence of strains from common pathogenic bacteria resistance to widely available and affordable antimicrobials is dangerously eroding their effectiveness [2]. Therapeutic options for this case are extremely limited; therefore, it is needed to develop the new antibacterial agent. Antibacterial screening from the plant is an alternative to start the invention of new antibacterial.

Anredera cordifolia (Ten.) v. Steenis has been proven to own pharmacological activity such as gastro protector, antidiabetic, anti-obesity, antihyperlipidemic, vasodilator, and wound healer [3-8]. Ethanolic extract of *A. cordifolia* leaves was actively proven to inhibit the growth of some Gram-positive and Gram-negative bacteria such as *Bacillus subtilis*, *Bacillus cereus*, *Escherichia coli*, *Pseudomonas aeruginosa*, methicillin resistant coagulase negative staphylococci, and methicillin sensitive *Staphylococcus aureus* (MSSA) [9]. Aqueous extract of *A. cordifolia* leaves showed inhibition toward *B. subtilis*, *E. coli*, *S. aureus*, and *P. aeruginosa* growth [10].

Antibacterial activity from *A. cordifolia* leaves extract, and its fractions have not been reported by the previous researcher. Therefore, in this study, antibacterial activities *A. cordifolia* extract and its fractions in different solvents were determined toward some pathogen bacteria. This study could show active compounds of *A. cordifolia* leaves as an antibacterial agent.

MATERIALS AND METHODS

Materials

A. cordifolia dried leaves were collected from Lembang - West Java, Indonesia. Plant identification was confirmed by Herbarium Bandungense, Bandung Institute of Technology. Standard ursolic acid (UA), oleanolic acid (OA), apigetrin, rutin, and antibiotic amoxicillin were obtained from Sigma-Aldrich (USA). Mueller Hinton Broth and Mueller-Hinton Agar were obtained from Oxoid.

The tested microorganisms were obtained from School of Pharmacy, Bandung Institute of Technology, Indonesia, which included five of Gram-positive bacteria such as *S. aureus* (ATCC 6538), MSSA, methicillin resistant *S. aureus* (MRSA), *B. subtilis* (ATCC 6633), and *B. cereus* (ATCC 11778), and four of Gram-negative bacteria such as *P. aeruginosa* (ATCC 9027), *E. coli* (ATCC 8939), *E. coli* H7 (O156), and ESBL *E. coli*.

Extraction

First method of extraction: Crude drug of *A. cordifolia* leaves was extracted by reflux method using ethanol 96% as a solvent, then vaporated to obtain a thick extract (EE1). Ethanolic extract was added with boiling water and separated in separatory funnel by liquid-liquid extraction method using n-hexane and ethyl acetate solvents. This process produced three fractions at the end n-hexane (HF), ethyl acetate (EAF), and water fraction (WF). The second method of extraction was using gradual extraction. Crude drug was extracted by reflux using n-hexane, ethyl acetate, and ethanol 96% solvents, respectively, so there were three extracts: n-hexane extract (HE), ethyl acetate extract (EAE), and ethanol extract (EE2). EE1, HF, EAF, WF, HE, EAE, and EE2 were sample in this study.

Phytochemical screening

The following tests performed on extracts and fractions were to detect the presence of flavonoid, tannin, quinone, saponin, alkaloid, and steroid/triterpenoid, as detailed previously [10].

Detection of UA and OA

Extracts, fractions, UA, and OA in methanol were spotted on a thin-layer chromatography (TLC) plate. For pre-chromatographic derivatization, the plate was developed in a horizontal chamber with 1% iodine in chloroform at 1.2 cm distance. The plate was placed in darkness for 10 minutes and then dried in stream of warm air to remove the excess of iodine [11]. Then, the plate was developed with toluene ethyl acetate formic acid (35:15:1 v/v/v) as the mobile phase. After drying, the plate was sprayed using H_2SO_4 reagent and then heated. The visualized spots were documented under visible light.

Detection of apigetrin and rutin

Extracts, fractions, apigetrin, and rutin in methanol were spotted on a TLC plate. The plate was developed with ethyl acetate methanol water (7:1:1 v/v/v) as the mobile phase. After that, the dried plate was sprayed using citroborate reagent and then heated. Spots were documented in UV 1366 nm.

Determination of minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC)

Extracts and fractions were dissolved in 10% dimethyl sulfoxide to obtain stock solutions. Standard solution (amoxicillin) was dissolved in sterile water. A two-fold serial microdilution method was used to determine MIC of samples. The lowest concentration that showed no visible growth was regarded as the MIC. Cells from the wells will show no growth of sub-cultured on MHA agar plate to determine the inhibition is reversible or permanent. The MBC was determined as the lowest concentration at which no growth occurred on the plate. The procedure was performed according to the Clinical and Laboratory Standards Institute guidelines [12].

Statistical analysis

MIC data of extracts, fractions, and amoxicillin against the same bacteria were analysis using Kruskal-Wallis test and continued with Mann-Whitney for pairwise comparison. $P < 0.05$ was considered significant.

RESULTS

Phytochemical screening

The result of phytochemical screening of extracts and fractions could be seen in Table 1.

Detection of UA and OA

TLC profile of extracts and fractions toward the existence of UA (Rf 0.61) and OA (Rf 0.72) showed that EE1, HF, and HE contained both of those acids (Fig. 1).

Detection of apigetrin and rutin

TLC resulted that qualitatively analyze the existence of apigetrin (Rf 0.50) and rutin (Rf 0.25) could be seen in Fig. 2. Based on the




Fig. 1: Thin-layer chromatography profile of *Anredera cordifolia* leaves extract and fractions for detecting ursolic acid and oleanolic acid (after spraying with H_2SO_4 reagent, under visible light). Tracks: Ethanolic extract from 1st extraction (EE1), n-hexane fraction (HF), ethyl acetate fraction (EAF), water fraction (WF), and n-hexane extract (HE); ethyl acetate extract (EAE), ethanolic extract from 2nd extraction (EE2), ursolic acid (UA), and oleanolic acid (OA)




Fig. 2: Thin-layer chromatography profile of *Anredera cordifolia* leaves extract and fractions for detecting apigetrin and rutin (after spraying with citroborate reagent, under UV 1366 nm). Tracks: Ethanolic extract from 1st extraction (EE1), hexane fraction (EF), ethyl acetate fraction (EAF), water fraction (WF), and hexane extract (HE); ethyl acetate extract (EAE), ethanolic extract from 2nd extraction (EE2), apigetrin (API), and rutin (R)

Table 1: Phytochemical screening of *Anredera cordifolia* extracts and fractions

Sample	Flavonoid	Tannin	Quinone	Saponin	Alkaloid	Steroid/triterpenoid
First extraction method						
Ethanolic extract (EE1)	+	-	-	+	+	+
N-hexane fraction (HF)	-	-	-	-	-	+
Ethyl acetate fraction (EAF)	+	-	-	-	+	+
Water fraction (WF)	+	-	-	+	+	+
Second extraction method						
N-hexane extract (HE)	-	-	-	-	-	+
Ethyl acetate extract (EAE)	+	-	-	-	+	+
Ethanolic extract (EE2)	+	-	-	+	+	+

+: Detected, -: Not detected

Table 2: Antibacterial activities of *A. cordifolia* extracts and fractions

Bacterial species (µg/ml)	1 st extraction method				2 nd extraction method			Amoxicillin
	EE1	HF	EAF	WF	HE	EAE	EE2	
<i>S. aureus</i>								
MIC	512 ^a	256 ^{a*}	512 ^a	1024	256 ^{a*}	512 ^a	4096	0.5 ^{**}
MBC	>4096	>4096	>4096	>4096	256 ^{a*}	>4096	>4096	0.5
MSSA								
MIC	2048	2048	4096	>4096	1024	4096	4096	8
MBC	>4096	4096	>4096	>4096	2048	>4096	>4096	8
MRSA								
MIC	512 ^a	1024	1024	>4096	512 ^a	512 ^a	4096	32 ^{**}
MBC	>4096	2048	2048	>4096	1024	1024	>4096	32
<i>B. subtilis</i>								
MIC	2048	512 ^a	1024	>4096	512 ^a	512 ^a	4096	4 ^{**}
MBC	4096	4096	4096	>4096	1024	>4096	>4096	4
<i>B. cereus</i>								
MIC	>4096	512 ^a	>4096	>4096	1024	2048	4096	0.5 ^{**}
MBC	>4096	1024	>4096	>4096	2048	>4096	>4096	2
<i>P. aeruginosa</i>								
MIC	1024	>4096	>4096	>4096	256 ^{a*}	1024	4096	16 ^{**}
MBC	2048	>4096	>4096	>4096	512 ^a	4096	>4096	32
<i>E. coli</i>								
MIC	>4096	>4096	>4096	>4096	512 ^{a*}	4096	>4096	1 ^{**}
MBC	>4096	>4096	>4096	>4096	2048	>4096	>4096	1
<i>E. coli</i> H7 (O156)								
MIC	>4096	2048	>4096	>4096	>4096	4096	4096	32
MBC	>4096	4096	>4096	>4096	>4096	>4096	>4096	32
ESBL <i>E. coli</i>								
MIC	4096	1024	2048	>4096	2048	4096	>4096	>256
MBC	>4096	2048	4096	>4096	4096	>4096	>4096	>256

Experiment was conducted triplicate. EE1: Ethanolic extract of *A. cordifolia* leaves from 1st extraction method, HF: n-hexane fraction, EAF: Ethyl acetate fraction, WF: Water fraction, HE: n-hexane extract, EAE: Ethyl acetate extract, EE2: Ethanolic extract from 2nd extraction method, MIC: Minimum inhibitory concentration, MBC: Minimum bactericidal concentration, ^ahad antibacterial potential; ^{*}significant difference compared to other extracts and fractions of *A. cordifolia* which had antibacterial potential against the same bacteria (p<0.05); ^{**}Significant difference compared to extracts and fractions of *A. cordifolia* which had antibacterial potential against the same bacteria (p<0.05). *A. cordifolia*: *Anredera cordifolia*, MSSA: Methicillin sensitive *Staphylococcus aureus*, MRSA: Methicillin resistant *Staphylococcus aureus*, ESBL: Extended-spectrum beta-lactamases, *E. coli*: *Escherichia coli*, *P. aeruginosa*: *Pseudomonas aeruginosa*, *B. cereus*: *Bacillus cereus*, *B. subtilis*: *Bacillus subtilis*, *S. aureus*: *Staphylococcus aureus*

observation, EE1, EAF, WF, and EAE contained apigetrin and rutin. Meanwhile, EE2 only contained apigetrin.

Determination of MIC and MBC

Antibacterial activities of the sample were presented in Table 2. Extracts and fractions had antibacterial potential if their MIC value was <1024 µg/ml.

DISCUSSION

Biologically active compounds commonly occur in low concentration in plants. An extraction technique is able to obtain an extract with high yield and with minimum changes in functional properties of the required extract. Several studies have reported variations in the biological activities of extracts prepared using different extraction techniques [13-15]. Antibacterial activity test from 1st extraction method (ethanol extract and its fraction) showed that they could only inhibit Gram-positive bacteria. HF could inhibit three microorganisms, *S. aureus* (MIC 256 µg/ml), *B. subtilis*, and *B. cereus* (MIC 512 µg/ml). Meanwhile, EE1 could only inhibit the growth of two microorganisms, *S. aureus* and MRSA (MIC 512 µg/ml); EAF could only inhibit the growth of *S. aureus* (MIC 512 µg/ml). The WF had no antibacterial activity. Meanwhile from 2nd extraction showed broader antibacterial activity. HE could inhibit three Gram-positive bacteria, *S. aureus* (MIC 256 µg/ml), MRSA, and *B. subtilis* (MIC 512 µg/ml) and two Gram-negative bacteria, *P. aeruginosa* (MIC 256 µg/ml) and *E. coli* (MIC 512 µg/ml). Only HE could inhibit Gram-negative bacteria, and its activity was significantly different compared to the other extracts and fractions (p<0.05). EAE could inhibit *S. aureus*, MRSA, and *B. subtilis* (MIC 512 µg/ml). EE2 had no antibacterial activity. Antibacterial activities of HF and HE against *S. aureus* were significantly higher than other extracts and fractions of *A. cordifolia* (p<0.05).

Amoxicillin was used as standard in this research. Its activities against all bacteria were the highest and significantly different compared to extracts and fractions of *A. cordifolia*, (p<0.05). Amoxicillin is shown to be effective against a wide range of infections caused by wide range of Gram-positive and Gram-negative bacteria [16]. Amoxicillin is still being drug of choice within the class because it has better pharmacokinetic than other β-lactam antibiotics for the treatment of infections due to susceptible organisms [17].

Based on MBC to MIC ratio of the sample which had antibacterial potential (MIC <1024 µg/ml), some sample were known to have bacteriostatic and bactericidal activity. Bacteriostatic activity has been defined as a ratio of MBC to MIC >4 and bactericidal activity had ratio MBC to MIC <4 [18]. In this study, EE1 and EAE had bacteriostatic activity, while HF and HE had bactericidal activity except its activity toward *B. cereus* and *P. aeruginosa*. EAE only had bactericidal activity toward MRSA. HE had the highest antibacterial activity among all extracts and fractions of *A. cordifolia* leaves.

The chemical constituents in plants or extracts are known to be biologically active ingredients. Some chemical constituents are considered as secondary metabolites components. They are directly responsible for different activity. Saponin was found in AF and EE1 that was obtained from first and second extraction methods. Flavonoid and alkaloid were found in most samples, except in HF dan HE. All samples contained steroid/triterpenoid. Some research showed that flavonoid, saponin, alkaloid, and steroid/triterpenoid had antibacterial activity [19-23].

TLC result showed that EE1, HF, and HE (Fig. 1) contained ursolic acid and OA. Several studies had proven that ursolic acid could inhibit the growth of pathogen bacteria such as *B. cereus*, *S. aureus*, *P. aeruginosa*, *E.*

coli, and MRSA [24-26]. Apigenin and rutin that were available in EE1, EAF, and EAE (Fig. 2), were also had been proven as antibacterial in few studies. *Tripleurospermum disciforme* aerial part extracts that contained apigenin, demonstrated antibacterial activity against *S. aureus* [27]. Rutin could inhibit growth of *S. aureus*, *B. subtilis*, and MRSA [28-29].

CONCLUSION

The HE had the highest antibacterial activity. Phytochemical screening of the extract showed the presence of steroid/triterpenoid. Ursolic acid and OA that have been known to have antibacterial activity from previous research were found in this extract. In addition, HE had more bactericidal activity compared to others samples. This study could be used to determine the active compounds of *A. cordifolia* as antibacterial agent for the future research.

ACKNOWLEDGMENTS

This research was funded by Ministry of Research, Technology and Higher Education of The Republic of Indonesia of The Republic of Indonesia through Doctoral Dissertation Grant.

REFERENCES

- World Health Organization. Antimicrobial Resistance from World Health Organization. Available from: <http://www.who.int/mediacentre/factsheets/fs194/en>. [Last accessed on 2017 Apr 24].
- World Health Organization. WHO Model Prescribing Information: Drugs Used in Bacterial Infections. Available from: <http://www.apps.who.int/medicinedocs/en/d/Js5406e>; <http://www.who.int/mediacentre/factsheets/fs194/en>. [Last accessed on 2017 Apr 24].
- Hornig CT, Chao HR, Lee CF, Hsueh CW, Chen FA. Gastro protective effect of Madeira vine against ethanol-induced gastric mucosal lesion in rat. *Asian J Chem* 2012;24(2):765-8.
- Elya B, Handayani R, Sauriasari R, Azizahwati, Hasyiyati US, Permana IT, et al. Antidiabetic activity and phytochemical screening of extracts from Indonesian plants by inhibition of alpha amylase, alpha glucosidase and dipeptidyl peptidase IV. *Pak J Bio Sci* 2015;18(6):279-84.
- Lestari D, Sukandar EY, Fidrianny I. *Anredera cordifolia* leaves extract as antihyperlipidemia and endothelial fat content reducer in male Wistar rat. *Int J Pharm Clin Res* 2015;7(6):435-9.
- Sukandar EY, Ridwan A, Sukmawan YP. Vasodilatation effect of ethanolic extract of *Anredera cordifolia*, *Sonchus arvensis* L, and ursolic acid on isolated rabbit aortic and frog heart. *Int J Pharm Pharm Sci* 2016;8(2):145-9.
- Sukandar EY, Kurniati NF, Nurdianti AN. Antiobesity effect of ethanol extract of *Anredera cordifolia* (Ten.) Steenis leaves on obese male Wistar rats induced by high-carbohydrate diet. *Int J Pharm Pharm Sci* 2016;8(4):171-3.
- Yuliani SH, Fudholi A, Pramono S, Marchaban. The effect of formula to physical properties of wound healing gel of ethanolic extract of binahong (*Anredera cordifolia* (Ten.) Steenis). *Int J Pharm Sci Res* 2012;27(3):4254-9.
- Garmana AN, Sukandar EY, Fidrianny I. Activity of several plant extracts against drug-sensitive and drug-resistant microbes. *Procedia Chem* 2012;13:164-9.
- Alonso Paz E, Cerdeiras MP, Fernandez J, Ferreira F, Moyna P, Soubes M, et al. Screening of Uruguayan medicinal plants for antimicrobial activity. *J Ethnopharmacol* 1995;45(1):67-70.
- Wójciak-Kosior M. Separation and determination of closely related triterpenic acids by high performance thin-layer chromatography after iodine derivatization. *J Pharm Biomed Anal* 2007;45(2):337-40.
- CLSI. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically; Approved Standard. CLSI Document M07-A9. 9th ed., Vol. 32. Wayne, PA: Clinical and Laboratory Standards Institute; 2012. p. 18-9.
- Hayouni EA, Abedrabba M, Bouix M, Hamdi M. The effects of solvents and extraction method on the phenolic contents and biological activities *in vitro* of Tunisian *Quercus coccifera* L. and *Juniperus phoenicea* L. Fruit extracts. *Food Chem* 2007;105:1126-34.
- Abah SE, Egwari LO. Methods of extraction and antimicrobial susceptibility testing of plant extracts. *Afr J Basic Appl Sci* 2011;3(5):205-9.
- Abdelfadel MM, Khalaf HH, Sharoba AM, Assous MT. Effect of extraction methods on antioxidant and antimicrobial activities of some spices and herbs extracts. *J Food Technol Nutr Sci* 2016;1(1):1-14.
- Kaur SP, Rao R, Nanda S. Amoxicillin: A broad spectrum antibiotic. *Int J Pharm Pharm Sci* 2011;3(3):30-7.
- Shahhet L, Alraghban D, Chehna D. Improvement of the physicochemical properties of amoxicillin trihydrate powder by recrystallization at different pH values. *Int J Pharm Pharm Sci* 2011;3(3):1-9.
- Pankey GA, Sabath LD. Clinical relevance of bacteriostatic versus bactericidal mechanisms of action in the treatment of Gram-positive bacterial infections. *Clin Infect Dis* 2004;38(6):864-70.
- Cushnie TP, Lamb AJ. Antimicrobial activity of flavonoids. *Int J Antimicrob Agents* 2005;26(5):343-56.
- Moussaoui F, Zellaoui A, Segueni N, Touil A, Rhouati S. Flavonoid constituents from Algerian *Launaea resedifolia* (O.K.) and their antimicrobial activity. *Rec Nat Prod* 2010;4(1):91-5.
- Lunga PK, Qin XJ, Yang XW, Kuate JR, Du ZZ, Gatsing D. Antimicrobial steroidal saponin and oleanane-type triterpenoid saponins from *Paullinia pinnata*. *BMC Complement Altern Med* 2014;14:369.
- Maatalah MB, Bouzidi NK, Bellahouel S, Merah B, Fortas Z, Soulimani R, et al. Antimicrobial activity of the alkaloids and saponin extracts of *Anabasis articulata*. *J Biotechnol Pharm Res* 2012;3(3):54-7.
- Cushnie TP, Lamb AJ. Alkaloids: An overview of their antibacterial, antibiotic-enhancing and antivirulence activities. *Int J Antimicrob Agents* 2014;44(5):377-86.
- Fontanay S, Grare M, Mayer J, Finance C, Duval RE. Ursolic, oleanolic and betulinic acids: Antibacterial spectra and selectivity indexes. *J Ethnopharmacol* 2008;120(2):272-6.
- Kim SG, Kim MJ, Jin D, Park SN, Cho E, Freire MO, et al. Antimicrobial effect of ursolic acid and oleanolic acid against methicillin resistant *Staphylococcus aureus*. *Korean J Microbiol* 2012;48(3):212-5.
- do Nascimento PG, Lemos TL, Bizerra AM, Arriaga AM, Ferreira DA, Santiago GM, et al. Antibacterial and antioxidant activities of ursolic acid and derivatives. *Molecules* 2014;19(1):1317-27.
- Tofighia Z, Molazema M, Doostdarb B, Tabanb P, Shahverdic AR, Samadi N, et al. Antimicrobial activities of three medicinal plants and investigation of flavonoids of *Tripleurospermum disciforme*. *Iran J Pharm Res* 2015;14(1):225-31.
- Orhan DD, Ozcelik B, Ozgen S, Ergun F. Antibacterial, antifungal, and antiviral activities of some flavonoids. *Microbiol Res* 2010;165(6):496-504.
- Dubey S, Ganeshpurkar A, Bansal D, Dubey N. Experimental studies on bioactive potential of rutin. *Chron Young Sci* 2013;4:153-7.

Author details

[Print](#) [Email](#)

Leliqia, Ni Putu Eka

[Is this you? Claim profile ↗](#)

Institut Teknologi Bandung, Bandung, Indonesia

 Subject area: [Pharmacology, Toxicology and Pharmaceutics](#) [Medicine](#)
[View potential author matches](#)

Profile actions

[Edit author profile](#)
[Connect to ORCID](#)
[Alerts](#)
[Set citation alert](#)
[Set document alert](#)

Documents by author

3


Total citations

2 by 2 documents

h-index: [?](#)

1

Document and citation trends:


[3 Documents](#)
[Cited by 2 documents](#)
[5 co-authors](#)
[Author history](#)
[Topics](#)

Preview users can view an author's latest 10 documents.

[Set document alert](#)

Document title	Authors	Year	Source	Cited by
Overview of efficacy, safety and phytochemical study of anredera cordifolia (Ten.) steenis	Leliqia, N.P.E., Sukandar, E.Y., Fidrianny, I.	2017	Pharmacologyonline	0
View abstract View abstract Related documents				
Antibacterial activities of Anredera Cordifolia (Ten.) V. Steenis leaves extracts and fractions Open Access	Leliqia, N.P.E., Sukandar, E.Y., Fidrianny, I.	2017	Asian Journal of Pharmaceutical and Clinical Research	0
View abstract View abstract Related documents				
Safety of garlic (allium sativum) and turmeric (curcuma domestica) extract in comparison with simvastatin on improving lipid profile in dyslipidemia patients Open Access	Sukandar, E.Y., Sudjana, P., Sigit, J.I., Leliqia, N.P.E., Lestari, F.	2013	Journal of Medical Sciences 2 (Faisalabad)	2
View abstract View abstract Related documents				

Preview users can view an author's latest 10 documents.

[^ Top of page](#)

[About Scopus](#)

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

[Language](#)

[日本語に切り替える](#)

[切换到简体中文](#)

[切换到繁體中文](#)

[Русский язык](#)

[Customer Service](#)

[Help](#)

[Contact us](#)

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

 RELX



Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

Home

Journal Rankings

Country Rankings

Viz Tools

Help

About Us

Asian Journal of Pharmaceutical and Clinical Research

Country
Subject Area and Category

[India](#) -  [SJR Ranking of India](#)

[Medicine](#)
[Pharmacology \(medical\)](#)

[Pharmacology, Toxicology and Pharmaceutics](#)
[Pharmaceutical Science](#)
[Pharmacology](#)

Publisher

[Asian Journal of Pharmaceutical and Clinical Research](#)

Publication type

Journals

ISSN

09742441, 24553891

Coverage

2009-ongoing

Scope

AJPCR (Asian J Pharm Clin Res) is peer reviewed, Monthly (Onward Jan 2017) open access Journal. This journal publishes original research in the field of Pharmaceutical and Clinical Sciences. The Journal has been designed to cover all the fields of research, which has any correlation and impact on Pharmaceutical Science. It aims to publish all the original research in field of science so a correlation can be made between these researches. Knowledge gained by such researches can be exposed to all and it can be brought in real utilization as all the branches of science are correlated and will assist all the researchers to potentiate their research capabilities.



[Homepage](#)

[How to publish in this journal](#)

[Contact](#)











[Join the conversation about this journal](#)

26

H Index

Quartiles





4k |

4k |



Asian Journal of Pharmaceutical and Clinical...
Q3 Pharmaceutical Science
best quartile
SJR 2018 0.17
powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

Z **zaib** 7 months ago

Does Asian Journal of Pharmaceutical and Clinical Research is a fake journal?

reply

S **Sultan Alshahrani** 8 months ago

Does AJPCR have impact factor? Thanks

reply

M **Manoj Kumar Mudigubba** 9 months ago

Asian journal of pharmaceutical and clinical research is Elsevier indexed journal?

reply



Elena Corera 9 months ago

Dear Manoj,

thank you for your request, all the journals included in SJR are indexed in Scopus. Elsevier / Scopus is our data provider.

Best Regards,
SCImago Team

A

Amn 9 months ago

Dear Elena

Do you advice me to publish my paper in which journal:
Indian journal of pharmaceutical sciences or
Asian journal of pharmaceutical and clinical research?
I want a journal powered by scopus now and for the next year.
Thanks

reply

S

Sangmesh 11 months ago

What is the difference between Simago and scopus

reply



Elena Corera 11 months ago

Dear Sangmesh, Scopus is a bibliographic database of scientific journals, the most comprehensive in the world. SCImago Journal & Country Rank is a platform in which scientometric indicators of journals included in Scopus and countries are displayed. Best Regards, SCImago Team

A

Ashwani 11 months ago

Is Asian Journal of Pharmaceutical and Clinical Research is a scopus indexed journal?

reply



Elena Corera 11 months ago

Dear Ashwani, all the journals included in the SJR are indexed in Scopus. Elsevier / Scopus is our data provider. Best Regards, SCImago Team

T

Tamara Amelia 11 months ago

we are beginner and want to publish our observation. We are interesting to publish it in this journal. could you tell us the price to publish in your journal? Thank you

Best regards,
Tamara Amelia
Faculty of Pharmacy
Universitas Indonesia

reply

B

Bayu Ardiansah 10 months ago

It is currently 100 USD (2018)



Elena Corera 11 months ago

Dear Tamara, we suggest you locate the author's instructions on the journal's website. Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)

I'm not a robot reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2019. Data Source: Scopus®





Asian Journal of Pharmaceutical and Clinical Research

E-ISSN: 2455-3891

P-ISSN: 0974-2441

Home

About

Current

Archives

Editorial Board

Submissions

Instructions to Authors

Contact Us

Search

Welcome to AJPCR (Online ISSN: 2455-3891 | Print ISSN: 0974-2441)

The Asian Journal of Pharmaceutical and clinical research (AJPCR) welcomes high-quality science that seeks to elucidate the Pharmaceutical and clinical research. The mission of the AJPCR is to bring significant, enduring research to the scientific community in the field of Pharmaceutical sciences and clinical research. AJPCR believe in its responsibility to safeguard the research and publish by providing high-quality review process and maintaining firm standards on data presentation and deposition. AJPCR goal is to help scientists to disclose their findings in the most efficient and effective way possible by keeping review times short, The heart of AJPCR mission is the publication of original research in the form of Articles, which represent top content in the journal and are highlighted with additional coverage.

Why publish with AJPCR?

- Timely decisions: The standard time from submission to first decision is 20 days for regular articles
- Timely broadcasting: The average time from initial submission to publication for regular Research Articles is 40 days,
- Constructive reviews from renowned scientist and researcher at all editorial levels
- No page limits for regular Articles
- Indexed in Google Scholar, Elsevier Products (EBSCO and EMBASE), [SCImago\(SJR\)](#), [CNKI \(China Knowledge Resource Integrated Database\)](#), CAS, CASSI (American Chemical Society), ICAAP, Scientific commons, PSOAR, Open-J-Gate, Index Medicus for WHO South-East Asia (IMSEAR), OAI, LOCKKS, OCLC (World Digital Collection Gateway), UIUC.Open Access

Abstracting and Indexing

Google Scholar, Elsevier Products (EBSCO and EMBASE), [SCImago\(SJR\)](#), Index Copernicus (ICV: 92), [CNKI \(China Knowledge Resource Integrated Database\)](#), CAS, CASSI (American Chemical Society), ICAAP, Scientific commons, PSOAR, Open-J-Gate, Index Medicus for WHO South-East Asia (IMSEAR), OAI, LOCKKS, OCLC (World Digital Collection Gateway), UIUC.

UGC Listed Journal





Online ISSN: 2455-3891

Print ISSN: 0974-2441

ICV: 92



Journal Metrics 2018

Source Normalized Impact per Paper (SNIP): 0.655

SCImago Journal Rank (SJR): 0.17

Print ISSN: 0974-2441

Online ISSN: 2455-3891



Visitor No. 05651

[Our Journals](#) || [Open Access Policy](#) || [Publication & Peer Review Policy](#) || [Publication Ethics](#)

The publication is licensed under a Creative Commons License (CC BY). [View Legal Code](#)

Copyright © 2018 All Rights Reserved, **Innovare Academic Sciences** | Powered By **CyberDairy**



Asian Journal of Pharmaceutical and Clinical Research

E-ISSN: 2455-3891

P-ISSN: 0974-2441

Editorial Board

Editorial Board

AJPCR is committed to have dynamic and potential advisory-editorial board. Those established in the field can directly send their resume. New people are first needed to serve as referee before being considered member of advisory-editorial board. Email your resume to editor@ajpcr.com

Editor-in-Chief

- **Dr. Anurekha Jain**

Dept. of Pharmaceutical Sciences, Jyoti Mahila Vidyapeeth University, Jaipur, Rajasthan

Email: anurekhajain@jvwu.ae.in

Associate Editor

- **Dr. Nuray Ari**

Prof., Department of Pharmacology, Faculty of Pharmacy, Ankara University, 06100 Ankara, Turkey.

Email: ari@ankara.edu.tr

- **Dr. Neeraj Upmanyu**

Prof., Peoples Institute of Pharmacy & Research Center, Bhopal, MP, India. Email:

drneerajupmanyu@gmail.com

Assistant Editor

- **Dr. Omotoso Abayomi Ebenezer**

Prof., Department of Pharmaceutical & Medicinal Chemistry. Faculty of Pharmaceutical Sciences, University of Port Harcourt, Nigeria. Email: abatoseb2001@yahoo.com

- **Dr. Vimal Kumar Jain**

Principal, ITM School of Pharmacy, ITM Universe, Vadodara & Associate Dean, Pharmacy, GTU, Ahmedabad

Editorial Board Members

- **Dr. Vikas Sharma**
Shri Rawatpura Sarkar Institute of Pharmacy, Datiya, MP, India
- **Dr. Sadia shakeel**
Prof., Department of Pharmacy Practice, Dow College of Pharmacy, Dow University of Health Sciences, Karachi, Pakistan.
- **Dr. Rupesh Kumar Gautam**
Associate Prof., ADINA Institute of Pharmaceutical Sciences, Sagar, MP, India
- **Dr. Farhan Ahmed Siddiqui**
Faculty of Pharmacy, Federal Urdu University Arts, Science and Technology Karachi, Sindh, Pakistan
- **Dr. Javad Sharifi Rad**
Department of Pharmacognosy, Faculty of Pharmacy, Zabol University of Medical Sciences, P.O. Box 61615-585 Zabol, Iran
- **Dr. Rajesh Mohanraj**
Dept. of Pharmacology, CMHS, UAE
- **Dr. Sami Saqf El Hait**
Junior Executive - Quality Control At Jamjoom Pharmaceuticals Company Limited jeddah, Saudi Arabia
- **Md. Moklesur Rahman Sarker**
Faculty of Medicine, University of Malaya, Malaysia
- **Dr. Hao Wu**
Postdoctoral Fellow At Ngm Biopharmaceuticals, Inc, South San Francisco, CA 94080, USA
- **Dr. Madhu Bala**
Scientist 'F' and Joint Director, Institute of Nuclear Medicine and Allied Sciences (INMAS), India
- **Dr. Mohanraj Rathinavelu**
Department of Pharmacy Practice, Raghavendra Institute of Pharmaceutical Education & Research, Riper, India
- **Dr. Sandip Narayan Chakraborty**
Research Asst, Translational Molecular Pathology, Ut Md Anderson Cancer Center, Life Sciences Plaza, Houston, TX 77030
- **Dr. Tushar Treembak Shelke**
Head of Department of Pharmacology and Research Scholar, In Jspms Charak College of Pharmacy & Research, Pune, India
- **Dr. D. Nagsamy Venkatesh**
Associate Prof., Department of Pharmaceutics, JSS College of Pharmacy, Ooty, TN India
- **Dr. Subas Chandra Dinda**
Professor-cum-Director: School of Pharmaceutical Education & Research (SPER), Berhampur University, Berhampur, Orissa, India.
- **Dr. Kanagala Vijaya Sri**
Associate professor, Malla Reddy College of Pharmacy, Maisammaguda, Dhullapally, Secunderabad

- **Dr. Jagdale Swati Changdeo**
Professor and Head, Department of Pharmaceutics, MAEER's Maharashtra Institute of Pharmacy, S.No.124,MIT Campus,Kothrud,Pune-411038
- **Dr. Biplab Kumar Dey**
Principal, Department of Pharmacy, Assam down town University, Sankar Madhab Path, Panikhaiti 781026, Guwahati, Assam, India
- **Dr. Yogesh Pandurang Talekar**
Research Associate, National Toxicology Centre
- **Dr. Indranil Chanda**
Assistant Professor, Girijananda Chowdhury Institute of Pharmaceutical Science, Hathkhowapara, Azara Guwahati-17, Assam, India.

Editorial office

Asian Journal of Pharmaceutical and Clinical Research
B-11, In front of Beema Hospital, Nayi Awadi, Mandasaur 458001, MP, India
E-mail:editor@ajpcr.com



Online ISSN: 2455-3891

Print ISSN: 0974-2441

ICV: 92



Journal Metrics 2018

Source Normalized Impact per Paper (SNIP): 0.655

SCImago Journal Rank (SJR): 0.17

Print ISSN: 0974–2441

Online ISSN: 2455–3891



Visitor No. 05651

[Our Journals](#) || [Open Access Policy](#) || [Publication & Peer Review Policy](#) || [Publication Ethics](#)

The publication is licensed under a [Creative Commons License \(CC BY\)](#). [View Legal Code](#)

Copyright © 2018 All Rights Reserved, **Innovare Academic Sciences** | Powered By **CyberDairy**



[HOME](#) / [ARCHIVES](#) / Vol 10 Issue 12 December 2017

EDITORIAL

FUTURE MEDICAL DOCTORS ARE READY TO PRESCRIBE ANTIMICROBIALS SAFELY AND RATIONALLY

MAINUL HAQUE

Pages 1-2



[View PDF](#)



[Abstract](#)



[Download PDF](#)

CASE STUDY(S)

SEIZURES WITH ANTI-SEIZURE DRUG

KEERTHANA BRATTIYA, BHUVANESHWARI K

Pages 3-5



[View PDF](#)



[Abstract](#)



[Download PDF](#)

REVIEW ARTICLE(S)

TARGETING THE TARGET USING NANOPARTICLES - A REVIEW

ARAVINTHRAJKUMAR G, GAYATHRI R, VISHNUPRIYA V

Pages 6-10

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

UTILITIES AND LIMITATIONS OF CURRENT ANIMAL MODELS OF DEPRESSION

ANDY RAMJATTAN, PINTO PEREIRA LEXLEY M, SAMEER DHINGRA

Pages 11-16

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

A REVIEW ON SCREENING METHODS OF BREAST CANCER, CANCER BIOMARKERS AND PHYTOCONSTITUENTS AGAINST BREAST CANCER

JAYASHREE VENUGOPAL, PRIYANKA S, RESHMA A

Pages 17-21

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

A COMPREHENSIVE REVIEW ON BARLERIA PRIONITIS (L.)

KAMINI SINGH, DEEPIKA SHARMA, GUPTA RS

Pages 22-29

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

A COMPREHENSIVE REVIEW ON COMORBID DEPRESSION IN PATIENTS WITH EPILEPSY

KEZIYA ANN MAMMEN, SATHESH KUMAR S

Pages 30-35

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

GENETIC DISORDER ALZHEIMER

PRATIBHA RANI, KAMALDEEP SINGH, ANANIA ARJUNA, SAVITA DEVI

Pages 36-39

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

PEROXISOME PROLIFERATOR-ACTIVATED RECEPTOR-GAMMA, AN EMERGING POTENTIAL TARGET TO COMBAT METABOLIC DISORDER

PARTHA SARATHI BAIRY

Pages 40-44

[View PDF](#) [Abstract](#) [Download PDF](#)**RECENT REPORTS ON PYRAZOLE-BASED BIOACTIVE COMPOUNDS AS CANDIDATE FOR ANTICANCER AGENTS**

BAYU ARDIANSAH

Pages 45-51

[View PDF](#) [Abstract](#) [Download PDF](#)**ORIGINAL ARTICLE(S)****COMPARISON OF SURFACE RADIATION DOSE TO THE GONADS BY RADIOGRAPHIC EXAMINATION OF THE LUMBAR SPINE USING COMPUTED RADIOGRAPHY AND DIRECT DIGITAL RADIOGRAPHY**

PRIYANKA PRIYANKA, RAHUL P KOTIAN, NITIKA C PANAKKAL

Pages 52-55

[View PDF](#) [Abstract](#) [Download PDF](#)**COMPARISON OF HERBAL MOUTHWASH WITH COMMERCIALLY AVAILABLE 0.2% CHLORHEXIDINE AND 2% BETADINE MOUTHWASHES IN PATIENTS AFTER STAGE-1 IMPLANT SURGERY**

JEMBULINGAM SABARATHINAM, DEEPAK NALLASWAMY, DEEPAK NALLASWAMY, MURALIDHARAN NP, MURALIDHARAN NP

Pages 56-58

[View PDF](#) [Abstract](#) [Download PDF](#)**COMPARATIVE ASSESSMENT OF THERAPEUTIC POTENTIAL OF VACHADI AND HINGWADI GHTRITA IN PATIENTS OF DEPRESSION**

PAWAR MADHURI, WELE ASMITA

Pages 59-62

[View PDF](#) [Abstract](#) [Download PDF](#)

EFFECT OF ETHANOLIC EXTRACT OF CYANOTIS CRISTATA LEAVES APPLIED TOPICALLY ON WOUND HEALING IN WISTAR RATS

ANURAG PATHAK, SMITA SHENOY, SUSHIL KIRAN, AVINASH A, DEEPAK NAYAK, NAVEEN KUMAR TONSE, AMOD TILAK, MANJUNATH SHETTY

Pages 63-67

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

EFFICACY OF BOERHAAVIA DIFFUSA L. ON DISRUPTION OF GONADOTROPINS AND TESTOSTERONE IN FLUORIDE INTOXICATED MALE RATS

SHASHI AGGARWAL, IMTIAZA KHAN

Pages 68-73

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

SUPPLEMENTATION WITH 2:1 RATIO OF N-6:N-3 POLYUNSATURATED FATTY ACID IMPROVES LIVER STEATOSIS AND SERUM CYTOKINE LEVELS IN YOUNG OBESE BALINESE WOMEN: A RANDOMIZED CLINICAL TRIAL

I WAYAN WETA, MAHADEWA TJOKORDA GB, WAYAN P SUTIRTAYASA, ANAK AGUNG NGURAH SUBAWA, SAFARINA G MALIK, I PUTU EKA WIDYADHARMA

Pages 74-79

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

ANTI-ATHEROSCLEROTIC ACTIVITY OF ELAEOCARPUS GANITRUS ROXB. IN CHOLESTEROL-FED RABBITS

PRATIBHA KUMARI JAIN, PRIYANKA SHARMA, SURESH C JOSHI

Pages 80-84

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

STUDY ON APHRODISIAC ACTIVITY OF OLEA DIOICA ROXB. BARK, LEAF EXTRACTS, AND ITS PURE COMPOUND USING WISTAR ALBINO RATS

ASHWATHANARAYANA R, RAJA NAIKA

Pages 85-98

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

CYTOTOXIC LABDANE DITERPENOIDS FROM ANDROGRAPHIS PANICULATA (BURM.F.) NEES

MARIA CARMENS TAN, GLENN G OYONG, CHIEN CHANG SHEN, CONSOLACION Y RAGASA

Pages 99-104

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

THE COMPARISON OF THE EFFECT BETWEEN ALGINATE-BASED RAFT-FORMING LIQUID AND ALGINATE LIQUID ON GASTROESOPHAGEAL REFLUX DISEASE AND GASTRIC ULCER IN RATS

HAKIM BANGUN, ANAYANTI ARIANTO, RIRIN ASTYA, GONTAR A SIREGAR

Pages 105-110

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

AUTOMATIC FEATURE EXTRACTION FOR BREAST DENSITY SEGMENTATION AND CLASSIFICATION

ASWATHY K CHERIAN, POOVAMMAL E, MALATHY C

Pages 111-115

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

PERCEIVED POSTPARTUM STRESS AND COPING STRATEGIES AMONG POSTNATAL MOTHERS AT AIMS, KOCHI

RACHEL MARGRET MATHEW, ANJU PHILIP T, SREEJAMOL MG

Pages 116-119

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

A RAPID AND SENSITIVE LIQUID CHROMATOGRAPHY- MASS SPECTROMETRY/MASS SPECTROMETRY METHOD FOR ESTIMATION OF PIOGLITAZONE, KETO PIOGLITAZONE AND HYDROXY PIOGLITAZONE IN HUMAN PLASMA

REVATHI NAGA LAKSHMI PONNURI, PRAHLAD PRAGALLAPATI, RAVINDRA N, VENKATA BASAVESWARA

RAO MANDAVA

Pages 120-128

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

IN VITRO FREE RADICAL SCAVENGING ACTIVITY AND IN VIVO ANTICATARACT POTENTIAL OF HEMIDESMUS INDICUS ROOTS IN STREPTOZOTOCIN INDUCED DIABETIC RATS

PRABHAKAR TIRUMANI, RAJASHEKHAR AV

Pages 129-136

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

PHYSIOCHEMICAL ANALYSIS OF PRETREATED BIOMEDICAL WASTES

SREEREMYA S, RAJIV P

Pages 137-140

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

IS ACQUISITION OF METHICILLIN-RESISTANT STAPHYLOCOCCUS AUREUS - AN OCCUPATIONAL HAZARD FOR MEDICAL STUDENTS IN INDIA?

SHREYAS K, RADHAKRISHNA M, ASHWINI HEGDE, POOJA RAO

Pages 141-144

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

BACTERIAL CONTAMINATION OF HANDS OF MEDICAL INTERNS AND UNDERGRADUATE STUDENTS

SUNANDHA SENTHIL, ASHWINI HEGDE, VAMAN KULKARNI, RADHAKRISHNA M

Pages 145-149

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

VIRTUAL SCREENING, PHARMACOPHORE MODELING, AND QUANTITATIVE STRUCTURE ACTIVITY RELATIONSHIP STUDIES ON HISTAMINE 4 RECEPTOR

SHOBANA SUGUMAR

Pages 150-154

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

RAPID AND ECONOMICAL QUANTITATIVE DETERMINATION OF SEVERAL ANTIHYPERTENSIVE AGENTS IN PRESENCE OF HYDROCHLOROTHIAZIDE BY ISOCRATIC REVERSED-PHASE HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY IN THEIR PHARMACEUTICAL PREPARATIONS

PANCHUMARTHY RAVISANKAR, DEVALA RAO G, MD SHAHEEM SULTHANA, SUPRIYA K, MOUNIKA G, NAVEENA P

Pages 155-159

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

SCREENING OF TEN HERBAL EXTRACTS TO COMPARE AND INVESTIGATE EFFECT ON GLUCOSE LOWERING AND ANOREXIC ACTIVITY IN WISTAR RATS

DEEPTI KAUSHALKUMAR JANI, SUNITA GOSWAMI

Pages 160-165

[View PDF](#) [Abstract](#) [Download PDF](#)**ASSOCIATION OF LIPID PROFILES WITH 10-YEAR ATHEROSCLEROTIC CARDIOVASCULAR DISEASE RISK: STUDY AMONG SUBJECTS IN SLEMAN DISTRICT OF YOGYAKARTA INDONESIA**

RITA SUHADI, DITA MARIA VIRGINIA, CHRISTIANUS HERU SETIAWAN

Pages 166-170

[View PDF](#) [Abstract](#) [Download PDF](#)**INCREASED URINARY 8-OXO-7,8-DIHYDRO-2-DEOXYGUANOSINE EXCRETION IN A SAMPLE OF EGYPTIAN CHILDREN WITH BETA THALASSEMIA MAJOR: MARKER FOR LIPID PEROXIDATION-INDUCED DNA DAMAGE**

MONA A ELABD, DINA ABU ZEID, MARWA A ELHADY, MAGED A EL WAKEEL, GHADA M EL-KASSAS, RANIA N SABRY, EMAN AWADALLAH

Pages 171-174

[View PDF](#) [Abstract](#) [Download PDF](#)**ANTIBACTERIAL ACTIVITIES OF ANREDERA CORDIFOLIA (TEN.) V. STEENIS LEAVES EXTRACTS AND FRACTIONS**

NI PUTU EKA LELIQIA, ELIN YULINAH SUKANDAR, IRDA FIDRIANNY

Pages 175-178

[View PDF](#) [Abstract](#) [Download PDF](#)**PHARMACOLOGICAL EVALUATION OF ANTINEOPLASTIC ACTIVITY OF HYDROETHANOLIC EXTRACT OF UNRIPE FRUIT OF CARICA PAPAYA LINN USING ANIMAL MODEL**

PADMAM PRAVEENA, JAYAPALAN PRASANNAKUMARI JETHINLALKHOSH, VICTOR AROKIA DOSS

Pages 179-181

[View PDF](#) [Abstract](#) [Download PDF](#)

ANTICANCER ACTIVITY OF GARCINIA MORELLA CHLOROFORM FRACTION AND ITS ACTIVE COMPOUND GARCINOL ON NEUROBLASTOMA

BHASWATI CHOUDHURY, RAGHURAM KANDIMALLA, RUPJYOTI BHARALI, JIBON KOTOKY

Pages 182-185

[View PDF](#) [Abstract](#) [Download PDF](#)**IN SILICO STUDIES FOR VARIOUS ANTIBACTERIAL BENZIL AND ITS SUBSTITUTED ANALOGS**

NITHYA G, SUDHA R, BRINDHA DEVI P, CHARLES C KANAKAM

Pages 186-189

[View PDF](#) [Abstract](#) [Download PDF](#)**PHYTOCHEMICAL SCREENING AND ANTIDIABETIC ACTIVITY OF DIFFERENT LEAF EXTRACTS FROM LOTUS (NELUMBO NUCIFERA GAERTN.) IN STREPTOZOTOCIN INDUCED MICE**

RUTH MAYANA RUMANTI, MARLINE NAINGGOLAN, URIP HARAHAAP

Pages 190-192

[View PDF](#) [Abstract](#) [Download PDF](#)**ANTIBACTERIAL ACTIVITY OF THREE PLANT EXTRACTS AGAINST MULTIDRUG RESISTANCE PSEUDOMONAS AERUGINOSA**

YASSIR HAIDER ALMAWLAH, HADEEL ALAA, SANAA O ALJELAWI

Pages 193-197

[View PDF](#) [Abstract](#) [Download PDF](#)**QUANTITATIVE PHYTOCHEMICAL ANALYSIS AND ANTIOXIDANT ACTIVITIES OF SOME CITRUS FRUITS OF SOUTH INDIA**

PALLAVI M, RAMESH CK, KRISHNA V, SAMEERA PARVEEN, NANJUNDA SWAMY L

Pages 198-205

[View PDF](#) [Abstract](#) [Download PDF](#)**BINARY QUANTITATIVE STRUCTURE-ACTIVITY RELATIONSHIP ANALYSIS IN RETROSPECTIVE STRUCTURE-BASED VIRTUAL SCREENING CAMPAIGNS TARGETING ESTROGEN RECEPTOR ALPHA**

ENADE PERDANA ISTYASTONO, NUNUNG YUNIARTI, MAYWAN HARIONO, SRI HARTATI YULIANI,

FLORENTINUS DIKA OCTA RISWANTO

Pages 206-211

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

MOLECULAR DOCKING, MOLECULAR DYNAMICS, AND IN SILICO TOXICITY PREDICTION STUDIES OF COUMARIN, N-OXALYLGLYCINE, ORGANOSELENIUM, ORGANOSULFUR, AND PYRIDINE DERIVATIVES AS HISTONE LYSINE DEMETHYLASE INHIBITORS

FAUZAN ZEIN MUTTAQIN, TAUFIK MUHAMMAD FAKIH, HUBBI NASHRULLAH MUHAMMAD

Pages 212-215

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

EVALUATION OF RESPONSE TO ANTIHYPERTENSIVE DRUGS IN PREECLAMPSIA AND GESTATIONAL HYPERTENSION CASES WITH DIFFERENT ACE GENE INSERTION/DELETION GENOTYPES

MOHD ISHAQ, ARSHIYA AFREEN, FATMA ALI, JUHI AZIZ, NIDA MAKEEN, NUSRATH FATHIMA, P PADMAJA, MOHD MOHIUDDIN

Pages 216-219

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

A PROSPECTIVE STUDY ON GERIATRIC PRESCRIBING PATTERN AND MEDICATION ADHERENCE IN A TERTIARY CARE HOSPITAL

SEEMA ANJUM M, CHINCHU JACOB, CHINCHU JACOB, AKHIL BENCHAMIN, AKHIL BENCHAMIN, PRUDENCE A RODRIGUES, PRUDENCE A RODRIGUES

Pages 220-225

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

VANILLIN EXTRACTED FROM PROSO MILLET AND BARNYARD MILLET INDUCE APOPTOSIS IN HT-29 AND MCF-7 CELL LINE THROUGH MITOCHONDRIA MEDIATED PATHWAY

DEEPA PRIYA RAMADOSS, NAGESWARAN SIVALINGAM

Pages 226-229

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

EFFECTIVENESS OF LOW-COST CUSTOMIZED SADDLE PILLOW ON PAIN IN POST-EPISIOTOMY PATIENTS

NAIR SHALINI KRISHNAN, ANBUPRIYA M, SIVAKUMAR VPR

Pages 230-233

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

KNOWLEDGE AND SELFCARE PRACTICES AMONG DIABETICS

SHEEBA S, SNEHA AK, VEENA BIJU

Pages 234-237

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

ELECTRONIC INFORMED CONSENT MODEL DEVELOPMENT FOR IMPLEMENTATION IN CLINICAL TRIALS IN UKRAINE

DOBROVA VY, ZUPANETS KÐŽ, KOLODYEZNA TY, YURIYTIMCHENKO TIMCHENKO

Pages 238-241

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

SYNTHESIS OF 2-HIDROXYXANTHONE FROM XANTHONE AS A BASIC MATERIAL FOR NEW ANTIMALARIAL DRUGS

AMANATIE AMANATIE, JUMINA JUMINA, MUSTOFA MUSTOFA, HANAFI M, LA ODE KADIDAE, SAHIDIN I

Pages 242-246

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

ISOLATION AND IDENTIFICATION OF PHYTOSTEROLS FROM BIGNONIA VENUSTA (L.)

MANOJ SHARMA, MUKESH KUMAR KHICHAR, AGRAWAL RD

Pages 247-251

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

DEVELOPMENT AND VALIDATION OF HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY WITH DIODE ARRAY DETECTOR METHOD TO ANALYSIS OF KAEMPFEROL MARKER FROM EXTRACTS OF POINCIANELLA PYRAMIDALIS (TUL) L.P QUEIROZ

AGNA HELIADE OLIVEIRA, VALMIR GOMESDE SOUZA, FÁBIO SANTOSDE SOUZA, RUI OLIVEIRA MACEDO

Pages 252-259

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

CHEMOPROTECTIVE EFFECT OF EDIBLE GASTROPOD, XANCUS PYRUM AND ITS USEFULNESS IN THE AMELIORATION OF CISPLATIN-INDUCED TOXICITY

BINDHU JAYAPRAKASH, EMILIN RENITTA, EMILIN RENITTA

Pages 260-265

[View PDF](#) [Abstract](#) [Download PDF](#)

STOOL FORM SCALE AS AN INDICATOR OF KLUTUK BANANA (MUSA BALBISIANA COLLA) FRUIT EXTRACTS INHIBITION EFFECT AGAINST SHIGELLA DYSENTERIAE ATCC 13313 IN VIVO

SRI AGUNG FITRI KUSUMA, PUSPAGITA WARDHANI, ELLIN FEBRINA

Pages 266-268

[View PDF](#) [Abstract](#) [Download PDF](#)

COMBINATION OF SAMBILOTO (ANDROGRAPHIS PANICULATA (BURM.F.) NEES) EXTRACT AND SPIRULINA (ARTHROSPIRA PLATENSIS GOMONT) TO PREVENT ANEMIA IN MICE INFECTED WITH PLASMODIUM BERGHEI ANKA

KUSMARDI KUSMARDI, BERNA ELYA, TRI WAHYUNI, RAFIKA INDAH PARAMITA

Pages 269-273

[View PDF](#) [Abstract](#) [Download PDF](#)

DETERMINATION OF BIOLOGICAL ACTIVITIES OF THREE MARINE ALGAE COLLECTED FROM VISAKHAPATNAM COAST

SATYA LAKSHMI S

Pages 274-279

[View PDF](#) [Abstract](#) [Download PDF](#)

CARDIOPROTECTIVE EFFECT OF PHYLLANTHUS FRATERNUS LEAVES EXTRACT AGAINST CYCLOPHOSPHAMIDE - INDUCED MYOCARDIAL INJURY IN RATS

RAKESH SINGH MOIRANGTHEM, NGANGOM GUNINDRO, DIPDEBA SINGH TAKHELLAMBAM, SUCHETA DEVI KHURAIJAM, MEENA N, RITA S

Pages 280-285

[View PDF](#) [Abstract](#) [Download PDF](#)

FACTORS INFLUENCING COMPLIANCE TO THERAPEUTIC REGIMEN AMONG PATIENTS WITH HYPERTENSION

DHANYA R DHARAN, MOLY KT

Pages 286-289

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

ANTIOXIDANT PROFILE AND PHYTOCHEMICAL CONTENT OF DIFFERENT PARTS OF SUPER RED DRAGON FRUIT (HYLOCEREUS COSTARICENSIS) COLLECTED FROM WEST JAVA-INDONESIA

IRDA FIDRIANNY, NADIA ILHAM, RIKA HARTATI

Pages 290-294

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

A NEW ELECTROCHEMICAL SENSOR BASED ON MODIFIED CARBON NANOTUBE-GRAPHITE MIXTURE PASTE ELECTRODE FOR VOLTAMMETRIC DETERMINATION OF RESORCINOL

MANJUNATHA JG

Pages 295-300

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

EVALUATION OF ANTIHYPERGLYCAEMIC EFFECT OF BUTEA MONOSPERMA LEAF EXTRACT ON ADRENALIN INDUCED AND HIGH GLUCOSE FEED ANIMAL MODEL

BISWARANJAN RAY, SANTOSH KUMAR MAHAPATRA, PRADIPTA KUMAR BEHERA, ASHOK K PANIGRAHI

Pages 301-304

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

ANTIOXIDANT AND ANTICANCER POTENTIALS OF BIOFABRICATED SILVER NANOPARTICLES

MERINA PAUL DAS, L JEYANTHI REBECCA

Pages 305-308

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

THE EFFECT OF MAHKOTA DEWA (PHALERIA MACROCARPA) (SCHEFF.) FRUIT PERICARP EXTRACT ON INOS IN MICE COLON INTERMITTENTLY-INDUCED BY DEXTRAN SODIUM SULFATE

KUSMARDI KUSMARDI, DILLA SHAVERA, ARI ESTUNINGTYAS, ARYO TEDJO, BAMBANG

PRIYOSOERYANTO

Pages 309-312

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

A COMPARATIVE STUDY OF EFFICACY AND SAFETY AMONG METFORMIN WITH SITAGLIPTIN, METFORMIN WITH VOGLIBOSE, AND METFORMIN WITH GLIMEPIRIDE IN PATIENTS WITH TYPE 2 DIABETES MELLITUS

KALA P, JAMUNA RANI R, KUMAR JS

Pages 313-316

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

PHYTOCHEMICAL INVESTIGATION AND ANTIMICROBIAL PROPERTIES OF DIOSCOREA BULBIFERA TUBER

DAHIYA P

Pages 317-319

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

EFFECTIVITY ANALYSIS OF NEUROPROTECTOR (VITAMIN B COMPLEX AND MECOBALAMIN) AS NEUROPATHIC PAIN SUPPORTIVE THERAPY IN ELDERLY WITH TYPE 2 DIABETES MELLITUS

MADE KRISNAADI JAYA, NIMADE OKA DWICANDRA

Pages 320-323

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

ASSESSMENT OF KNOWLEDGE AND PRACTICE TOWARD DISPOSAL OF EXPIRED DRUGS AMONG MEDICAL PROFESSIONALS IN CHENNAI - A CROSS-SECTIONAL PROSPECTIVE STUDY

UMAMAGESWARI ARUNACHALAM, GURU SCINDIA MV, PRIESTLY VIVEKKUMAR S, GEETHA R

Pages 324-327

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

ISOLATION OF SOME POTENTIAL PHYTOCOMPOUNDS FROM ADHATODA VASICA THROUGH GAS CHROMATOGRAPHY-MASS SPECTROSCOPY ANALYSIS

AZIZ MOHAMMAD KHAN, SEEMA BHADAURIA

Pages 328-332

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

A SIMPLE REVERSE PHASE-HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY METHOD DEVELOPMENT AND VALIDATION OF VALSARTAN IN BULK AND IT'S TABLET DOSAGE FORM

RAMREDDY GODELA, SHERISHA BHAVANI

Pages 333-337

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

ANALYSIS AND RESULTS: CONFIRMATORY FACTOR ANALYSIS THE MALAY VERSION OF DREEM INVENTORY WITH MEDICAL STUDENTS OF UNISZA, KUALA TERENGGANU, MALAYSIA

MAINUL HAQUE, MUHAMAD SAIFUL BAHRI YUSOFF, MD. ANWARUL AZIM MAJUMDER, ZAINAL ZULKIFLI, FARAH HANANI BINTI MOHD NASIR

Pages 338-344

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

IN SILICO STUDY OF ARYL EUGENOL DERIVATIVES AS ANTI-COLORECTAL CANCER BY INDUCING OF APOPTOSIS

FADILAH FADILAH, ARRY YANUAR, ADE ARSIANTI, RETNOSARI ANDRAJATI, ERNI HERNAWATI PURWANINGSIH

Pages 345-349

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

PREVALENCE OF HYPERTENSION AMONG ADULT POPULATION IN SLUMS OF WEST DELHI

SUPRIYA DWIVEDI, ZAOZIANLUNGLIU GONMEI, GURUDAYAL SINGH TOTEJA, NIDHI SRIVASTAVA, NAVAL KISHORE VIKRAM, SPRIHA RAO, CHETRAM MEENA

Pages 350-352

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

IMMUNE PROTECTION OF SALVINIA MOLESTA D.S. MITCHELL IN FRESHWATER CRAB OZIOTELPHUSA SENEX SENEX BACTERIALLY CHALLENGED WITH AEROMONAS HYDROPHILA

NITHYA TG, JAYANTHI J, RAGUNATHAN MG, DEVAKUMAR D

Pages 353-356

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

IDENTIFICATION OF AMPC β -LACTAMASE-PRODUCING CLINICAL ISOLATES OF ESCHERICHIA COLI

TANUSHREE BARUA GUPTA, MALINI SHARIFF, THUKRAL SS, S.S THUKRAL

Pages 357-361

[View PDF](#) [Abstract](#) [Download PDF](#)

DECREASEMENT OF LYSOPHOSPHATIDYLCHOLINE LEVEL, NF-KB EXPRESSION, INTIMA MEDIA THICKNESS AND IMPROVEMENT OF INSULIN RESISTANCE BY DARAPLADIB TREATMENT: IN VIVO STUDIES OF TYPE 2 DIABETES MELLITUS SPRAGUE-DAWLEY RAT MODEL

TITIN ANDRI WIHASTUTI, DINDA ZAHRA PUTRI ANDIYANI, SRI ANDARINI, TEUKU HERIANSYAH

Pages 362-365

[View PDF](#) [Abstract](#) [Download PDF](#)

PREPARATION AND STABILITY ASSESSMENT OF PERILLA FRUTESCENS SEED OIL POWDER

CHAIYAVAT CHAIYASUT, BHAGAVATHI SUNDARAM SIVAMARUTHI, NETNAPA MAKHAMRUEANG, PERIYANAINA KESIKA, SASITHORN SIRILUN, SARTJIN PEERAJAN

Pages 366-369

[View PDF](#) [Abstract](#) [Download PDF](#)

THE POTENCY OF INDONESIA'S POMEGRANATE PEEL ETHANOL EXTRACT (PUNICA GRANATUM LINN.) AS ANTI-INFLAMMATORY AGENT IN MICE COLON INDUCED BY DEXTRAN SODIUM SULFATE: FOCUS ON CYCLOOXYGENASE-2 AND INOS EXPRESSIONS

KUSMARDI KUSMARDI, DONY HERMANTO, ARI ESTUNINGYAS, ARYO TEDJO, BAMBANG P PRIOSOERYANTO

Pages 370-375

[View PDF](#) [Abstract](#) [Download PDF](#)

FORMULATION AND EVALUATION OF NOVEL ANTIBACTERIAL AND ANTI-INFLAMMATORY CREAM CONTAINING MUNTINGIA CALABURA LEAVES EXTRACT

MAHENDRAN SEKAR, NUR SYAHIRA ABDUL JALIL

Pages 376-379

[View PDF](#) [Abstract](#) [Download PDF](#)

TASTE ALTERATION AND QUALITY OF LIFE OF PATIENTS RECEIVING CHEMOTHERAPY

ANOOJA ANTONY, SHEELA PAVITHRAN

Pages 380-383

[View PDF](#) [Abstract](#) [Download PDF](#)**COMPARATIVE STUDY ON ANTIMICROBIAL ACTIVITY OF SEaweEDS**

SASIKALA C, GEETHA RAMANI D

Pages 384-386

[View PDF](#) [Abstract](#) [Download PDF](#)**ANTIOXIDANT POTENTIALS AND SIMULTANEOUS ESTIMATION OF QUERCETIN, RUTIN, AND GALLIC ACID IN CURCUMA SPECIES**

SOWMYA PRIYA M, SANGILIMUTHU A, KARPAGAVALLI M, RAMKUMAR M, JOSPHIN NIRMALA A

Pages 387-391

[View PDF](#) [Abstract](#) [Download PDF](#)**LUPEOL VALIDATION AND QUANTIFICATION IN HETEROPOGON CONTORTUS (L.) BEAUV. (SPEAR GRASS) THROUGH HIGH-PERFORMANCE THIN-LAYER CHROMATOGRAPHY**

NAVJOT KAUR, RAGHBIR CHAND GUPTA

Pages 392-395

[View PDF](#) [Abstract](#) [Download PDF](#)**STAPHYLOCOCCUS AUREUS INFECTIONS AND THEIR ANTIBIOTIC SUSCEPTIBILITY PROFILE AT A TERTIARY CARE HOSPITAL**

ANTHONY N OFULAH, SHASHIDHAR VISHWANATH, SHASHIDHAR VISHWANATH, BARNINI BANERJEE, SHASHIDHAR VISHWANATH, KIRAN CHAWLA, BARNINI BANERJEE, BARNINI BANERJEE, KIRAN CHAWLA, KIRAN CHAWLA

Pages 396-399

[View PDF](#) [Abstract](#) [Download PDF](#)**PHYTOCHEMICAL AND ANTIOXIDANT ACTIVITY OF CAMELLIA MISTLETOE (KORTHALSELLA JAPONICA) EXTRACTS**

MIN YOUNG KIM

Pages 400-403

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

COMPARISON OF ANTI-DIABETIC AND ANTI-OXIDANT ACTIVITY OF WILD AND CULTIVATED VARIETY OF RAUWOLFIA SERPENTINA

SAVEENA CHAUHAN, AMRINDER KAUR, MANISH VYAS, GOPAL L KHATIK

Pages 404-406

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

ESTIMATION OF CARDIAC SURGERY RESULTS

ALEXEY LEONIDOVICH CHARYSHKIN, ALEXANDER NIKOLAEVICH YUDIN

Pages 407-410

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

INVESTIGATION THE ASSOCIATION BETWEEN MTHFR GENE POLYMORPHISM AND HOMOCYSTEINE IN IRANIAN PREGNANT WOMEN

ELHAM SADAT SONBOLESTAN, HOSSEIN SAZEGAR, NOOSHA ZIA-JAHROMI, FARZANEH MOHAMADI FARSANI

Pages 411-414

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

SCREENING OF ANTIOXIDANT PROPERTY IN MEDICINAL PLANTS BELONGING TO THE FAMILY APOCYNACEAE ZAHRA

ZAHRA IQBAL, MOHAMMED SHARIQ IQBAL, KUMKUM MISHRA

Pages 415-418

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

BERBERINE HYDROCHLORIDE COULD PROVE TO BE A PROMISING BULLET AGAINST CLOSTRIDIUM DIFFICILE INFECTION: A PRELIMINARY STUDY FROM SOUTH INDIA

RITUPARNA CHAKRABORTY, RICHARD LOBO, MUKHYAPRANA M PRABHU, SHASHIKIRAN UMAKANTH, GOUTAM CHOWDHURY, ASISH K MUKHOPADHYAY, THANDAVARAYAN RAMAMURTHY, MAMATHA BALLAL

Pages 419-424

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

DETERMINATION OF 5H-BENZO[2,3][1,4]OXAZEPINO[5,6-B]INDOLES IN RAT PLASMA BY REVERSED-PHASE HIGH-PERFORMANCE LIQUID CHROMATOGRAPHIC-ULTRAVIOLET METHOD: APPLICATION TO PHARMACOKINETIC STUDIES

ASHOK K SINGH, VINIT RAJ, AMIT RAI, AMIT K KESHARI, PRANESH KUMAR, SUDIPTA SAHA

Pages 425-430

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

LEPTIN AND ADIPONECTIN LEVELS OF ASTHMATIC IRAQI CHILDREN IN HILLA PROVINCE

AFRAH AL NAZARMAMoori, MUFEEDJALIL EWADH, SUHAYR ESSA ALQAYSI

Pages 431-434

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

A CLINICAL STUDY TO EVALUATE THE EFFICACY OF SAINDHAVADI TAILA MATRA VASTI IN THE MANAGEMENT OF AMAVATA W.S.R. RHEUMATOID ARTHRITIS

ARCHANA NEGI, PUNITA PANDEY, BHUPINDER SINGH

Pages 435-439

 [View PDF](#)  [Abstract](#)  [Download PDF](#)

ANTICANCER POTENTIAL OF Î±-MANGOSTIN

MUCHTARIDI MUCHTARIDI, CINDY APRILLIANIE WIJAYA

Pages 440-445

 [View PDF](#)  [Abstract](#)  [Download PDF](#)



Online ISSN: 2455-3891

Print ISSN: 0974-2441

ICV: 92



Journal Metrics 2018

Source Normalized Impact per Paper (SNIP): 0.655

SCImago Journal Rank (SJR): 0.17

Print ISSN: 0974-2441

Online ISSN: 2455-3891



Visitor No. 05651

[Our Journals](#) || [Open Access Policy](#) || [Publication & Peer Review Policy](#) || [Publication Ethics](#)

The publication is licensed under a Creative Commons License (CC BY). [View Legal Code](#)

Copyright © 2018 All Rights Reserved, **Innovare Academic Sciences** | Powered By **CyberDairy**

ANTIBACTERIAL ACTIVITIES OF ANREDERA CORDIFOLIA (TEN.) STEENIS LEAVES EXTRACTS AND FRACTIONS

by Ni Putu Eka Leliqia

Submission date: 01-Jul-2019 01:55PM (UTC+0700)

Submission ID: 1148360125

File name: DERA_CORDIFOLIA_TEN._V._STEENIS_LEAVES_EXTRACT_AND_FRACTIONS.pdf (705.27K)

Word count: 3433

Character count: 19481

ANTIBACTERIAL ACTIVITIES OF ANREDERA CORDIFOLIA (TEN.) V. STEENIS LEAVES EXTRACTS AND FRACTIONSNI PUTU EKA LELIQIA^{1,2*}, ELIN YULINAH SUKANDAR¹, IRDA FIDRIANNY³

¹Department of Pharmacology-Clinical Pharmacy, School of Pharmacy, Bandung Institute of Technology, Indonesia. ²Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Udayana University, Indonesia. ³Department of Pharmaceutical Biology, School of Pharmacy, Bandung Institute of Technology, Indonesia. Email: leliqia@gmail.com

Received: 22 July 2017, Revised and Accepted: 25 August 2017

ABSTRACT**Objective:** This study aims to determine antibacterial activity of *Anredera cordifolia* leaves extracts and fractions.**Methods:** Crude drug was extracted using two methods. First method was extraction by reflux using ethanol 96% and then fractionated by liquid-liquid extraction using n-hexane and ethyl acetate. Second method was gradually extraction by reflux using n-hexane, ethyl acetate and ethanol 96%, respectively. Phytochemical screening was applied to all extracts and fractions, followed by thin-layer chromatography using ursolic acid, oleanolic acid (OA), apigetrin, and rutin as reference substances. A two-fold serial microdilution method was used to determine minimum inhibitory concentration (MIC) against *Staphylococcus aureus* (ATCC 6538), methicillin-susceptible *S. aureus*, methicillin resistant *S. aureus* (MRSA), *Bacillus subtilis* (ATCC 6633), *Bacillus cereus* (ATCC 11778), *Pseudomonas aeruginosa* (ATCC 9027), *Escherichia coli* (ATCC 8939), *E. coli* H7 (O156), and ESBL *E. coli*. Bacteriostatic and bactericidal activities were determined using minimum bactericidal concentration/MIC ratio.**Results:** The ethanolic extract, n-hexane and ethyl acetate fractions of *A. cordifolia* from the first method had antibacterial activity against *S. aureus*, MRSA, *B. subtilis*, and *B. cereus* (MIC 256-512 µg/ml). However, n-hexane and ethyl acetate extract from the second method had broad spectrum of antibacterial activity, which could inhibit the growth of *S. aureus*, MRSA, *B. subtilis*, *P. aeruginosa*, and *E. coli* (MIC 256-512 µg/ml). Extracts and fractions showed bacteriostatic and bactericidal activities, but n-hexane extract has most bactericidal activity. Furthermore, steroid/triterpenoid, ursolic, and OA were found in this extract.**Conclusion:** The n-hexane extract from the second method showed the highest antibacterial activity.**Keywords:** *Anredera cordifolia*, Extracts, Fractions, Antibacterial.

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2017.v10i12.21503>

INTRODUCTION

Bacterial infections remain as the main leading cause of death, particularly in developing countries [1]. Infection treatment with antibacterial has reduced the morbidity and improved patient's survival with bacterial infections. However, in many cases, the increasing prevalence of strains from common pathogenic bacteria resistance to widely available and affordable antimicrobials is dangerously eroding their effectiveness [2]. Therapeutic options for this case are extremely limited; therefore, it is needed to develop the new antibacterial agent. Antibacterial screening from the plant is an alternative to start the invention of new antibacterial.

Anredera cordifolia (Ten.) v. Steenis has been proven to own pharmacological activity such as gastro protector, antidiabetic, anti-obesity, antihyperlipidemic, vasodilator, and wound healer [3-8]. Ethanolic extract of *A. cordifolia* leaves was actively proven to inhibit the growth of some Gram-positive and Gram-negative bacteria such as *Bacillus subtilis*, *Bacillus cereus*, *Escherichia coli*, *Pseudomonas aeruginosa*, methicillin resistant coagulase negative staphylococci, and methicillin sensitive *Staphylococcus aureus* (MSSA) [9]. Aqueous extract of *A. cordifolia* leaves showed inhibition toward *B. subtilis*, *E. coli*, *S. aureus*, and *P. aeruginosa* growth [10].

Antibacterial activity from *A. cordifolia* leaves extract, and its fractions have not been reported by the previous researcher. Therefore, in this study, antibacterial activities *A. cordifolia* extract and its fractions in different solvents were determined toward some pathogen bacteria. This study could show active compounds of *A. cordifolia* leaves as an antibacterial agent.

MATERIALS AND METHODS**Materials**

A. cordifolia dried leaves were collected from Lembang - West Java, Indonesia. Plant identification was confirmed by Herbarium Bandungense, Bandung Institute of Technology. Standard ursolic acid, oleanolic acid (OA), apigetrin, rutin, and antibiotic amoxicillin were obtained from Sigma-Aldrich (USA). Mueller Hinton Broth and Mueller-Hinton Agar were obtained from Oxoid.

The tested microorganisms were obtained from School of Pharmacy, Bandung Institute of Technology, Indonesia, which included five of Gram-positive bacteria such as *S. aureus* (ATCC 6538), MSSA, methicillin resistant *S. aureus* (MRSA), *B. subtilis* (ATCC 6633), and *B. cereus* (ATCC 11778), and four of Gram-negative bacteria such as *P. aeruginosa* (ATCC 9027), *E. coli* (ATCC 8939), *E. coli* H7 (O156), and ESBL *E. coli*.

Extraction

First method of extraction: Crude drug of *A. cordifolia* leaves was extracted by reflux method using ethanol 96% as a solvent, then vaporated to obtain a thick extract (EE1). Ethanolic extract was added with boiling water and separated in separatory funnel by liquid-liquid extraction method using n-hexane and ethyl acetate. This process produced three fractions at the end n-hexane (HF), ethyl acetate (EAF), and water fraction (WF). The second method of extraction was using gradual extraction. Crude drug was extracted by reflux using n-hexane, ethyl acetate, and ethanol 96% solvents, respectively, so there were three extracts: n-hexane extract (HE), ethyl acetate extract (EAE), and ethanol extract (EE2). EE1, HF, EAF, WF, HE, EAE, and EE2 were sample in this study.

Phytochemical screening

The following tests performed on extracts and fractions were to detect the presence of flavonoid, tannin, quinone, saponin, alkaloid, and steroid/triterpenoid, as detailed previously [10].

Detection of UA and OA

Extracts, fractions, UA, and OA in methanol were spotted on a thin-layer chromatography (TLC) plate. For pre-chromatographic derivatization, the plate was developed in a horizontal chamber with 1% iodine in chloroform at 1.2 cm distance. The plate was placed in darkness for 10 minutes and then dried in stream of warm air to remove the excess of iodine [11]. Then, the plate was developed with toluene ethyl acetate formic acid (35:15:1 v/v/v) as the mobile phase. After drying, the plate was sprayed using H_2SO_4 reagent and then heated. The visualized spots were documented under visible light.

Detection of apigetrin and rutin

Extracts, fractions, apigetrin, and rutin in methanol were spotted on TLC plate. The plate was developed with ethyl acetate methanol water (7:1:1 v/v/v) as the mobile phase. After that, the dried plate was sprayed using citroborate reagent and then heated. Spots were documented in UV 366 nm.

Determination of minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC)

Extracts and fractions were dissolved in 10% dimethyl sulfoxide to obtain stock solutions. Standard solution (amoxicillin) was dissolved in sterile water. A two-fold serial microdilution method was used to determine MIC of samples. The lowest concentration that showed no visible growth was regarded as the MIC. Cells from the wells will show no growth of sub-cultured on MHA agar plate to determine the inhibition is reversible or permanent. The MBC was determined as the lowest concentration at which no growth occurred on the plate. The procedure was performed according to the Clinical and Laboratory Standards Institute guidelines [12].

Statistical analysis

MIC data of extracts, fractions, and amoxicillin against the same bacteria were analysis using Kruskal-Wallis test and continued with Mann-Whitney for pairwise comparison. $P < 0.05$ was considered significant.

RESULTS

Phytochemical screening

The result of phytochemical screening of extracts and fractions could be seen in Table 1.

Detection of UA and OA

TLC profile of extracts and fractions toward the existence of UA (Rf 0.61) and OA (Rf 0.72) showed that EE1, HF, and HE contained both of those acids (Fig. 1).

Detection of apigetrin and rutin

TLC resulted that qualitatively analyze the existence of apigetrin (Rf 0.50) and rutin (Rf 0.25) could be seen in Fig. 2. Based on the




Fig. 1: Thin-layer chromatography profile of *Anredera cordifolia* leaves extract and fractions for detecting ursolic acid and oleanolic acid (after spraying with H_2SO_4 reagent, under visible light). Tracks: Ethanolic extract from 1st extraction (EE1), n-hexane fraction (HF), ethyl acetate fraction (EAF), water fraction (WF), and n-hexane extract (HE); ethyl acetate extract (EAE), ethanolic extract from 2nd extraction (EE2), ursolic acid (UA), and oleanolic acid (OA)




Fig. 2: Thin-layer chromatography profile of *Anredera cordifolia* leaves extract and fractions for detecting apigetrin and rutin (after spraying with citroborate reagent, under UV 366 nm). Tracks: Ethanolic extract from 1st extraction (EE1), hexane fraction (HF), ethyl acetate fraction (EAF), water fraction (WF), and hexane extract (HE); ethyl acetate extract (EAE), ethanolic extract from 2nd extraction (EE2), apigetrin (API), and rutin (R)

Table 1: Phytochemical screening of *Anredera cordifolia* extracts and fractions

Sample	Flavonoid	Tannin	Quinone	Saponin	Alkaloid	Steroid/triterpenoid
First extraction method						
Ethanolic extract (EE1)	+	-	-	+	+	+
N-hexane fraction (HF)	-	-	-	-	-	+
Ethyl acetate fraction (EAF)	+	-	-	-	+	+
Water fraction (WF)	+	-	-	+	+	+
Second extraction method						
N-hexane extract (HE)	-	-	-	-	-	+
Ethyl acetate extract (EAE)	+	-	-	-	+	+
Ethanolic extract (EE2)	+	-	-	+	+	+

+: Detected, -: Not detected

Table 2: Antibacterial activities of *A. cordifolia* extracts and fractions

Bacterial species ($\mu\text{g/ml}$)	1 st extraction method				2 nd extraction method			Amoxicillin
	EE1	HF	EAF	WF	HE	EAE	EE2	
<i>S. aureus</i>								
MIC	512 ^a	256 ^{**}	512 ^a	1024	256 ^{**}	512 ^a	4096	0.5 ^{**}
MBC	>4096	>4096	>4096	>4096	256 ^{**}	>4096	>4096	0.5
MSSA								
MIC	2048	2048	4096	>4096	1024	4096	4096	8
MBC	>4096	4096	>4096	>4096	2048	>4096	>4096	8
MRSA								
MIC	512 ^a	1024	1024	>4096	512 ^a	512 ^a	4096	32 ^{**}
MBC	>4096	2048	2048	>4096	1024	1024	>4096	32
<i>B. subtilis</i>								
MIC	2048	512 ^a	1024	>4096	512 ^a	512 ^a	4096	4 ^{**}
MBC	4096	4096	4096	>4096	1024	>4096	>4096	4
<i>B. cereus</i>								
MIC	>4096	512 ^a	>4096	>4096	1024	2048	4096	0.5 ^{**}
MBC	>4096	1024	>4096	>4096	2048	>4096	>4096	2
<i>P. aeruginosa</i>								
MIC	1024	>4096	>4096	>4096	256 ^{**}	1024	4096	16 ^{**}
MBC	2048	>4096	>4096	>4096	512 ^a	4096	>4096	32
<i>E. coli</i>								
MIC	>4096	>4096	>4096	>4096	512 ^{**}	4096	>4096	1 ^{**}
MBC	>4096	>4096	>4096	>4096	2048	>4096	>4096	1
E 33 H7 (O156)								
MIC	>4096	2048	>4096	>4096	>4096	4096	4096	32
MBC	>4096	4096	>4096	>4096	>4096	>4096	>4096	32
ESBL <i>E. coli</i>								
MIC	4096	1024	2048	>4096	2048	4096	>4096	>256
MBC	>4096	2048	4096	>4096	4096	>4096	>4096	>256

Experiment was conducted triplicate. EE1: Ethanolic extract of *A. cordifolia* leaves from 1st extraction method, HF: hexane fraction, EAF: Ethyl acetate fraction, WF: Water fraction, HE: n-hexane extract, EAE: Ethyl acetate extract, EE2: Ethanolic extract from 2nd extraction method, MIC: Minimum inhibitory concentration, MBC: Minimum bactericidal concentration, had antibacterial potential; *significant difference compared to other extracts and fractions of *A. cordifolia* which had antibacterial potential against the same bacteria ($p < 0.05$); **Significant difference compared to other extracts and fractions of *A. cordifolia* which had antibacterial potential against the same bacteria ($p < 0.05$). *A. cordifolia*: *Anredera cordifolia*, MSSA: Methicillin sensitive *Staphylococcus aureus*, MRSA: Methicillin resistant *Staphylococcus aureus*, ESBL: Extended-spectrum beta-lactamases, *E. coli*: *Escherichia coli*, *P. aeruginosa*: *Pseudomonas aeruginosa*, *B. cereus*: *Bacillus cereus*, *B. subtilis*: *Bacillus subtilis*, *S. aureus*: *Staphylococcus aureus*

observation, EE1, EAF, WF, and EAE contained apigenin and rutin. Meanwhile, EE2 only contained apigenin.

Determination of MIC and MBC

Antibacterial activities of the sample were presented in Table 2. Extracts and fractions had antibacterial potential if their MIC value was <1024 $\mu\text{g/ml}$.

DISCUSSION

Biologically active compounds commonly occur in low concentration in plants. An extraction technique is able to obtain an extract with high yield and with minimum changes in functional properties of the required extract. Several studies have reported variations in the biological activities of extracts prepared using different extraction techniques [13-15]. Antibacterial activity test from 1st extraction method (ethanol extract and its fraction) showed that they could only inhibit Gram-positive bacteria. HF could inhibit three microorganisms, *S. aureus* (MIC 256 $\mu\text{g/ml}$), *B. subtilis*, and *B. cereus* (MIC 512 $\mu\text{g/ml}$). Meanwhile, EE1 could only inhibit the growth of two microorganisms, *S. aureus* and MRSA (MIC 512 $\mu\text{g/ml}$); EAF could only inhibit the growth of *S. aureus* (MIC 512 $\mu\text{g/ml}$). The WF had no antibacterial activity. Meanwhile from 2nd extraction showed broader antibacterial activity. HE could inhibit three Gram-positive bacteria, *S. aureus* (MIC 256 $\mu\text{g/ml}$), MRSA, and *B. subtilis* (MIC 512 $\mu\text{g/ml}$) and two Gram-negative bacteria, *P. aeruginosa* (MIC 256 $\mu\text{g/ml}$) and *E. coli* (MIC 512 $\mu\text{g/ml}$). Only HE could inhibit Gram-negative bacteria, and its activity was significantly different compared to the other extracts and fractions ($p < 0.05$). EAE could inhibit *S. aureus*, MRSA, and *B. subtilis* (MIC 512 $\mu\text{g/ml}$). EE2 had no antibacterial activity. Antibacterial activities of HF and HE against *S. aureus* were significantly higher than other extracts and fractions of *A. cordifolia* ($p < 0.05$).

Amoxicillin was used as standard in this research. Its activities against all bacteria were the highest and significantly different compared to extracts and fractions of *A. cordifolia*, ($p < 0.05$). Amoxicillin is shown to be effective against a wide range of infections, used by wide range of Gram-positive and Gram-negative bacteria [16]. Amoxicillin is still being drug of choice within the class because it has better pharmacokinetic than other β -lactam antibiotics for the treatment of infections due to susceptible organisms [17].

Based on MBC to MIC ratio of the sample which had antibacterial potential (MIC <1024 $\mu\text{g/ml}$), some sample were known to have bacteriostatic and bactericidal activity. Bacteriostatic activity has been defined as a ratio of MBC to MIC >4 and bactericidal activity had ratio MBC to MIC <4 [18]. In this study, EE1 and EAE had bacteriostatic activity, while HF and HE had bactericidal activity except its activity toward *B. cereus* and *P. aeruginosa*. EAE only had bactericidal activity toward MRSA. HE had the highest antibacterial activity among all extracts and fractions of *A. cordifolia* leaves.

The chemical constituents in plants or extracts are known to be biologically active ingredients. Some chemical constituents are considered as secondary metabolites components. They are directly responsible for different activity. Saponin was found in AF and EE1 that was obtained from first and second extraction methods. Flavonoid and alkaloid were found in most samples, except in HF dan HE. All samples contained steroid/triterpenoid. Some research showed that flavonoid, saponin, alkaloid, and steroid/triterpenoid had antibacterial activity [19-23].

TLC result showed that EE1, HF, and HE (Fig. 1) contained ursolic acid and OA. Several studies had proven that ursolic acid could inhibit the growth of pathogen bacteria such as *B. cereus*, *S. aureus*, *P. aeruginosa*, *E.*

coli, and MRSA [24-26]. Apigetrin and rutin that were available in EE1, EAF, and EAE (Fig. 2), were also had been proven as antibacterial in few studies. *Tripleurospermum disciforme* aerial part extracts that contained apigetrin, demonstrated antibacterial activity against *S. aureus* [27]. Rutin could inhibit growth of *S. aureus*, *B. subtilis*, and MRSA [28-29].

CONCLUSION

The HE had the highest antibacterial activity. Phytochemical screening of the extract showed the presence of steroid/triterpenoid. Ursolic acid and OA that have been known to have antibacterial activity from previous research were found in this extract. In addition, HE had more bactericidal activity compared to others samples. This study could be used to determine the active compounds of *A. cordifolia* as antibacterial agent for the future research.

ACKNOWLEDGMENTS

This research was funded by Ministry of Research, Technology and Higher Education of The Republic of Indonesia of The Republic of Indonesia through Doctoral Dissertation Grant.

REFERENCES

- World Health Organization. Antimicrobial Resistance from World Health Organization. Available from: <http://www.who.int/mediacentre/factsheets/fs194/en>. [Last accessed on 2017 Apr 24].
- World Health Organization. WHO Model Prescribing Information: Drugs Used in Bacterial Infections. Available from: <http://www.apps.who.int/medicinedocs/en/d/Js5406e>; <http://www.who.int/mediacentre/factsheets/fs194/en>. [Last accessed on 2017 Apr 24].
- Homg CT, Chao HR, Lee CF, Hsueh CW, Chen FA. Gastro protective effect of Madeira vine against ethanol-induced gastric mucosal lesion in rat. *Asian J Chem* 2012;24(2):765-8.
- Elya B, Handayani R, Sauriansa R, Azizahwati, Hasyati US, Permana IT, et al. Antidiabetic activity and phytochemical screening of extracts from Indonesian plants by inhibition of alpha amylase, alpha glucosidase and dipeptidyl peptidase IV. *Pak J Bio Sci* 2015;18(6):279-84.
- Lestari D, Sukandar EY, Fidrianny I. *Anredera cordifolia* leaves extract as antihyperlipidemia and endothelial fat content reducer in male Wistar rat. *Int J Pharm Clin Res* 2015;7(6):435-9.
- Sukandar EY, Ridwan A, Sukmawan YP. Vasodilatation effect of ethanolic extract of *Anredera cordifolia*, *Sonchus arvensis* L., and ursolic acid on isolated rabbit aortic and frog heart. *Int J Pharm Pharm Sci* 2016;8(2):145-9.
- Sukandar EY, Kurniati NF, Nurdianti AN. Antiobesity effect of ethanol extract of *Anredera cordifolia* (Ten.) Steenis leaves on obese male Wistar rats induced by high-carbohydrate diet. *Int J Pharm Pharm Sci* 2016;8(4):171-3.
- Yuliani SH, Fudholi A, Pramono S, Marchaban. The effect of formula to physical properties of wound healing gel of ethanolic extract of binahong (*Anredera cordifolia* (Ten.) Steenis). *Int J Pharm Sci Res* 2012;27(3):4254-9.
- Garmana AN, Sukandar EY, Fidrianny I. Activity of several plant extracts against drug-sensitive and drug-resistant microbes. *Procedia Chem* 2012;13:164-9.
- Alonso Paz E, Cerdeiras MP, Fernandez J, Ferreira F, Moyna P, Soubes M, et al. Screening of Uruguayan medicinal plants for antimicrobial activity. *J Ethnopharmacol* 1995;45(1):67-70.
- Wójciak-Kosior M. Separation and determination of closely related triterpenic acids by high performance thin-layer chromatography after iodine derivatization. *J Pharm Biomed Anal* 2007;45(2):337-40.
- CLSI. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically; Approved Standard. CLSI Document M07-A9. 9th ed., Vol. 32. Wayne, PA: Clinical and Laboratory Standards Institute; 2012. p. 18-9.
- Hayouni EA, Abedrabba M, Bouix M, Hamdi M. The effects of solvents and extraction method on the phenolic contents and biological activities *in vitro* of Tunisian *Quercus coccifera* L. and *Juniperus phoenicea* L. Fruit extracts. *Food Chem* 2007;105:1126-34.
- Abah SE, Egwari LO. Methods of extraction and antimicrobial susceptibility testing of plant extracts. *Afr J Basic Appl Sci* 2011;3(5):205-9.
- Abdelfadel MM, Khalaf HH, Sharoba AM, Assous MT. Effect of extraction methods on antioxidant and antimicrobial activities of some spices and herbs extracts. *J Food Technol Nutr Sci* 2016;1(1):1-14.
- Kaur SP, Rao R, Nanda S. Amoxicillin: A broad spectrum antibiotic. *Int J Pharm Pharm Sci* 2011;3(3):30-7.
- Shahhet L, Alraghban D, Chehna D. Improvement of the physicochemical properties of amoxicillin trihydrate powder by recrystallization at different pH values. *Int J Pharm Pharm Sci* 2011;3(3):1-9.
- Pankey GA, Sabath LD. Clinical relevance of bacteriostatic versus bactericidal mechanisms of action in the treatment of Gram-positive bacterial infections. *Clin Infect Dis* 2004;38(6):864-70.
- Cushnie TP, Lamb AJ. Antimicrobial activity of flavonoids. *Int J Antimicrob Agents* 2005;26(5):343-56.
- Moussaoui F, Zellagui A, Segueni N, Touil A, Rhouati S. Flavonoid constituents from Algerian *Launaea resedifolia* (O.K.) and their antimicrobial activity. *Rec Nat Prod* 2010;4(1):91-5.
- Lunga PK, Qin XJ, Yang XW, Kuatie JR, Du ZZ, Gatsing D. Antimicrobial steroidal saponin and oleanane-type triterpenoid saponins from *Paulinia pinnata*. *BMC Complement Altern Med* 2014;14:369.
- Maatalah MB, Bouzidi NK, Bellahouel S, Merah B, Fortas Z, Soufmani R, et al. Antimicrobial activity of the alkaloids and saponin extracts of *Anabasis articulata*. *J Biotechnol Pharm Res* 2012;3(3):54-7.
- Cushnie TP, Cushnie B, Lamb AJ. Alkaloids: An overview of their antibacterial, antibiotic-enhancing and antivirulence activities. *Int J Antimicrob Agents* 2014;44(5):377-86.
- Fontanay S, Grare M, Mayer J, Finance C, Duval RE. Ursolic, oleanolic and betulinic acids: Antibacterial spectra and selectivity indexes. *J Ethnopharmacol* 2008;120(2):272-6.
- Kim SG, Kim MJ, Jin D, Park SN, Cho E, Freire MO, et al. Antimicrobial effect of ursolic acid and oleanolic acid against methicillin resistant *Staphylococcus aureus*. *Korean J Microbiol* 2012;48(3):212-5.
- do Nascimento PG, Lemos TL, Bizerra AM, Arriga AM, Ferreira DA, Santiago GM, et al. Antibacterial and antioxidant activities of ursolic acid and derivatives. *Molecules* 2014;19(1):1317-27.
- Tofighia Z, Molazema M, Doostdarb B, Tabanb P, Shahverdic AR, Samadi N, et al. Antimicrobial activities of three medicinal plants and investigation of flavonoids of *Tripleurospermum disciforme*. *Iran J Pharm Res* 2015;14(1):225-31.
- Orhan DD, Ozcelik B, Ozgen S, Ergun F. Antibacterial, antifungal, and antiviral activities of some flavonoids. *Microbiol Res* 2010;165(6):496-504.
- Dubey S, Ganeshpurkar A, Bansal D, Dubey N. Experimental studies on bioactive potential of rutin. *Chron Young Sci* 2013;4:153-7.

ANTIBACTERIAL ACTIVITIES OF ANREDERA CORDIFOLIA (TEN.) STEENIS LEAVES EXTRACTS AND FRACTIONS

ORIGINALITY REPORT

18%

SIMILARITY INDEX

10%

INTERNET SOURCES

12%

PUBLICATIONS

13%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Matan, N.. "Durability of rubberwood (*Hevea brasiliensis*) treated with peppermint oil, eucalyptus oil, and their main components", *International Biodeterioration & Biodegradation*, 200907
Publication 2%
- 2 www.i-scholar.in
Internet Source 1%
- 3 Submitted to UIN Maulana Malik Ibrahim Malang
Student Paper 1%
- 4 P. Mučaji, M. Nagy. "Contribution to the TLC separation of ursolic and oleanolic acid mixture", *Acta Facultatis Pharmaceuticae Universitatis Comenianae*, 2011
Publication 1%
- 5 www.biomedcentral.com
Internet Source 1%

6

Internet Source

1%

7

Ashok K. Singh, B. Gordon, D. Hewetson, K. Granley, M. Ashraf, U. Mishra, D. Dombrovskis. "Screening of steroids in horse urine and plasma by using electron impact and chemical ionization gas chromatography-mass spectrometry", Journal of Chromatography A, 1989

Publication

1%

8

Submitted to CVC Nigeria Consortium

Student Paper

1%

9

dashboardffup.info

Internet Source

1%

10

Submitted to Higher Education Commission
Pakistan

Student Paper

1%

11

hal.univ-lorraine.fr

Internet Source

1%

12

pbc.gda.pl

Internet Source

1%

13

www.nijophasr.com

Internet Source

1%

14

Megdiche-Ksouri, Wided, Najla Trabelsi,
Khaoula Mkadmini, Soumaya Bourgou, Amira

1%

Noumi, Mejdi Snoussi, Rahma Barbria, Olfa Tebourbi, and Riadh Ksouri. "Artemisia campestris phenolic compounds have antioxidant and antimicrobial activity", Industrial Crops and Products, 2015.

Publication

15

www.ijddr.in

Internet Source

<1%

16

academic.oup.com

Internet Source

<1%

17

Submitted to Kingston University

Student Paper

<1%

18

Submitted to Nanyang Technological University, Singapore

Student Paper

<1%

19

HANEN NAJJAA. "PHYTOCHEMICAL SCREENING AND ANTIBACTERIAL PROPERTIES OF ALLIUM ROSEUM L., A WILD EDIBLE SPECIES IN NORTH AFRICA : WILD ALLIUM ROSEUM", Journal of Food Biochemistry, 06/2011

Publication

<1%

20

tci-thaijo.org

Internet Source

<1%

21

www.ijper.org

Internet Source

<1%

22

Renmin Liu, Lingyi Kong, Aifeng Li, Ailing Sun.
" Preparative Isolation and Purification of
Saponin and Flavone Glycoside Compounds
from by High-Speed Countercurrent
Chromatography ", Journal of Liquid
Chromatography & Related Technologies, 2007

Publication

<1%

23

Gamage, G.R.. "Effectiveness of antimicrobial
coated oriented polypropylene/polyethylene
films in sprout packaging", Food Research
International, 200908

Publication

<1%

24

Muthu K. Shanmugam, Xiaoyun Dai, Alan
Prem Kumar, Benny K.H. Tan, Gautam Sethi,
Anupam Bishayee. "Ursolic acid in cancer
prevention and treatment: Molecular targets,
pharmacokinetics and clinical studies",
Biochemical Pharmacology, 2013

Publication

<1%

25

Adeshina, G, C-L Okeke, N Onwuegbuchulam,
and J Ehinmidu. "Preliminary studies on
antimicrobial activities of ethanolic extracts of
Ficus sycomorus Linn. and Ficus platyphylla
Del. (Moraceae)", International Journal of
Biological and Chemical Sciences, 2010.

Publication

<1%

26

macha.itc.griffith.edu.au

Internet Source

<1%

27 orca.cf.ac.uk Internet Source <1%

28 Aydin, S.. "The effects of thyme volatiles on the induction of DNA damage by the heterocyclic amine IQ and mitomycin C", Mut.Res.-Genetic Toxicology and Environmental Mutagenesis, 20050307 Publication <1%

29 linknovate.com Internet Source <1%

30 Submitted to University of Mauritius Student Paper <1%

31 Submitted to National University of Singapore Student Paper <1%

32 Submitted to RMIT University Student Paper <1%

33 Jacek Cybulski. "Long-Alkyl-Chain Quaternary Ammonium Lactate Based Ionic Liquids", Chemistry - A European Journal, 10/20/2008 Publication <1%

34 Submitted to University of Strathclyde Student Paper <1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On