ISSN 2550-1283 VOLUME 1 NUMBER 1 - FEBRUARY 2017











PUBLISHED BY INSTITUTE FOR RESEARCH AND COMMUNITY SERVICES, UDAYANA UNIVERSITY, BALI, INDONESIA

### ISSN 2550-1283

## JOURNAL OF VETERINARY AND ANIMAL SCIENCE

Editorial Team

*Editor-in-Chief* I Wayan Suardana (Udayana University, Indonesia)

Co Editors in Chief I Nyoman Suarsana (Udayana University, Indonesia) I Nyoman Suartha (Udayana University, Indonesia) Ni Ketut Suwiti (Udayana University, Indonesia) I Gusti Ayu Agung Suartini (Udayana University, Indonesia) Aida L.T. Rompis (Udayana University, Indonesia)

Editorial Boards

A.A. Ayu Mirah Adi	I Ketut Puja (Indonesia)
(Udayana University, Indonesia)	(Udayana University, Indonesia)
Aris Haryanto	Komang G. Wiryawan
(Gadjah Mada University, Indonesia)	(Bogor Agriculture University, Indonesia)
Christian Bauer	Michael Haryadi Wibowo
(Justus Liebig University, Germany)	(Gadjah Mada University, Indonesia)
Cristina W. Cunha	Mirnawati Sudarwanto
(USDA-Washington State University, USA)	(Bogor Agriculture University, Indonesia)
Dewa Ketut Harya Putra	Nadeeka Wawegama
Udayana University, Indonesia)	(The University of Melbourne, Australia)
Dyah Ayu Widiasih	Oemar Akineden
(Gadjah Mada University, Indonesia)	(Justus Liebig University, Germany)
I G N Kade Mahardika	Saleha Abd. Azis
(Udayana University, Indonesia)	(Universiti of Putra Malaysia, Malaysia)
I Gede Mahardika	Wasmen Manalu
(Udayana University, Indonesia)	(Bogor Agriculture University, Indonesia)
I Made Damriyasa	Yasunobu Matsumoto
(Udayana University, Indonesia)	(The University of Tokyo, Jepan)
I Nyoman Sadra Dharmawan	Duangporn Pichpol
(Udayana University, Indonesia)	(Chiang Mai University, Thailand)

Technical Editors I G M Krisna Erawan (Udayana University, Indonesia) I N Kerta Besung (Udayana University, Indonesia)

Published by Institute for Research and Community Services Udayana University

Secretariat

Institute for Research and Community Services Udayana University Bukit Jimbaran, Badung, Bali, Indonesia Phone: +62-361- 704622 / 703367 Email: jvas@unud.ac.id



## Journal of Veterinary and Animal Science

Journal of Veterinary and Animal Science is a peer-reviewed journal which devoted to the advancement and dissemination of scientific knowledge concerning veterinary and animal science which includes research findings, case report, experimental design, and their application for the treatment of diseases in birds, wild and domestic animals. This journal published in English being distributed worldwide. It covers all the scientific and technological aspects of veterinary medicine in general, anatomy, physiology, biochemistry, pharmacology, microbiology, pathology, public health, parasitology, infectious diseases, clinical sciences, biotechnology, alternative veterinary medicine and other biomedical fields. In the field of animal science, the journal receives original manuscripts covering breeding and genetics, reproduction and physiology, nutrition, feed sciences, animal products, biotechnology, behavior, livestock farming system, sosio-economic, and policy.

## Journal of Veterinary Medicine and Animal Science Vol. 1 Vol. 1, February 2017

## **Table of Contents**

### **Original Article**

Antimicrobial Susceptibility Patterns and DNA Plasmid Profiles of	01-05
Escherichia Coli 0157:H7 Isolated From Feces of Chickens	
I Wayan Suardana, Iwan Harjono Utama, Michael Haryadi Wibowo	
Patological Changes in Liver and Gall Bladder of Bali Cattle Infected by	06-10
Fasciolosis	
I Made Kardena, Ida Bagus Oka Winaya, Anak Agung Ayu Mirah Adi, I Ketut	
Berata, Ida Bagus Windia Adnyana, I Made Sukada, Putu Agus Trisna Kusuma	
Antara	
Detection of Haemophilus Parasuis from The Respiratory Tract of Pigs	11-14
I Gusti Ketut Suarjana, I Nengah Kerta Besung, Ketut Tono PG	
The Characteristic of Egg Drop Syndrome Virus of Medan Isolate	15-19
Gusti Ayu Yuniati Kencana, I Nyoman Suartha, Arini Nurhandayani, Syamsidar	
Phenotypic Identification of Lactobacillus from Breast Milk Based on Their	20-23
Ability to Ferment Sugars	
Ni Made Indri Hapsari Arihantana, Ni Nyoman Puspawati	
The Performance of Balinese Gilt Reared Intensively and	24-28
Supplemented with Probiotic	
K. Budaarsa, A. W. Puger, T. I. Putri , I D.G.A. Udayana	
Herbal Extract as An Antibacterial Against Gram Positive Bacteria Causing	29-31
Dermatitis Complex	
I Nyoman Suartha, I Gusti Ketut Suarjana, Luh Made Sudimartini, I Made	
Merdana, I Made Dira Swantara	
Tracking Antigen Expression of Toxoplasma gondii in Mice by	32-35
Immuno (Cyto) Histochemistry	
Ida Ayu Pasti Apsari, Ida Bagus Oka Winaya, Tjokorda Sari Nindhia. Ida Bagus	
Ngurah Swacita	
Identification of Growth Hormone Gene of Bali Cattle with Qualitative	36-39
Superior in Bali Province	
Ni Ketut Suwiti, I Wayan Suardana, Ni Luh Watiniasih, I Nengah Kerta	
Besung, Chandra Yowani	

### **Case Report**

Macroconidia of Dermatophytes Fungi on Direct Microscopic Examination40-42Putu Ayu Sisyawati Putriningsih, I Putu Gede Yudhi Arjentinia40-42

# Detection of Haemophilus Parasuis from The Respiratory Tract of Pigs

#### I Gusti Ketut Suarjana<sup>\*</sup>, I Nengah Kerta Besung<sup>1</sup>, Ketut Tono PG

Faculty of Veterinary Medicine, Udayana University Jl. PB Sudirman Denpasar 80361.Phone: 0361223791 \*Corresponding author: <u>kt suarjana@unud.ac.id</u>

**Abstract.** Study in order to detect the presence of Haemophilus parasuis in pig farms in Tabanan, Badung, and Gianyar Bali have been undertaken. A total of 197samples including nasal and tracheal swabs, organs and synovial fluids were collected from 69, 28 and 100 animals from pig farms in Tabanan, Badung and Gianyar regency, respectively. Samples were collected mainly from animals which showing clinical signs of coughing, asphyxiate, swollen joints and death animals. Isolation and identification of Haemophilus spp was performed following the methods described by Priadi et.al. (2004) and Nedbalkova et al. (2006) with slightly modification. Antimicrobial sensitivity test was performed using the disc diffusion method by Kirby-Bauer.Haemophilus parasuis was detected from one sample which shown resistant to streptomycin only and sensitive to ampicillin, doxycycline and kanamycin, respectively.

#### Keywords: Hamemophilus spp, pigs, respiratory tract, sensitivity test

#### I. INTRODUCTION

Besides cows, buffaloes, sheep, and goats, pig is one of meat production animal. Majority of the pig farming in Bali is raised for meat consumption and also for ceremonies purposes. Apart from Bali pig farming can be found in other areas such as North Sumatra, Papua, East Nusa Tenggara and Moluccas. Some obstacles in pig farming including reduction in body weight, and infectious diseases. Infection due to Haemophilus spp particularly H. parasuis rarely reported in Indonesia, especially in Bali no report of disease due to this bacteria. This bacteria inhabits in the upper respiratory tract of pig. Several strains of Haemophilus have been reported to cause diseases which typically produce edematous of the serous such as fibrous polyserositis, polyarthritis, meningitis, and pneumonia [1][2][3]. Disease caused by Haemophilus spp. can affect pigs at any age but mostly early post weaning animals with clinical signs of fever up 40-41°C, difficulties in breathing, coughing, swollen joints, lameness and paralysis. The animals died within 2-5 days whereas recover animals suffer arthritis chronically, coughing and bronchitis [4][5]. Upon pathological examination meningitis, serositis, pleuritic,

bronchopneumonia, pericarditis and arthritis can be observed as sole or multiple manifestation [6].There are numbers of cases in the pig farms in Bali in which the animals showed clinical signs such as fever, respiratory distress (coughing, abdominal breathing), swollen joints and lameness of hind legs similar to the clinical signs of the disease caused by Haemophilus parasuis. This study aimed to observe the prevalence of Haemophilus spp in pig farms and its sensitivity to the antibiotics normally used in the pig farms in Bali.

#### II. RESEARCH METHOD

#### Samples

Samples including nasal swabs, tracheal swabs, organ and synovial fluids were collected from 197 pigs which showed signs such as coughing, asphyxiate, and swollen joints from the three regency namely: Tabanan, Badung and Gianyar, respectively. All samples were collected aseptically and kept in Stuart transport media.

#### Isolation and Identification of Bacteria

Isolation of Haemophilus spp was performed by culturing samples on to Sheep Blood Agar (SBA) close to Staphylococcus aureusstreak culture as a source of NAD (feeder culture). Plate culture were then kept in a plastic container with AnaerogenTM (OxoidR) for CO2 7-15% aerated, then incubated at 37°C for 24-48 hours following the methods by Priadi et al. (2004) and Nedbalkova et al. (2006) [7][8]. Identification was performed by the conventional methods which is Gram staining, oxidase and catalase test, and biochemical test using **API®** 20E<sup>™</sup>(Biomerieux). Characteristics of Haemophilus sp isolate were tiny colonies on blood agar close to the Staph. aureus streak culture, Gram negative rods, hemolytic, where hemolytic, catalase and oxidase varied. Isolates showing those characters were further identified by indole test, urease and fermentation of carbohydrate (glucose, arabinose, lactose, mannitol, xylose, galactose, mannose, raffinose, sorbitol and inositol [9]. Further confirmation was performed using the API® 20E<sup>™</sup> (Biomerieux).

#### Antimicrobial Sensitivity Test

Sensitivity test was performed following the disc diffusion method by Kirby-Bauer with slightly modification [10]. The antibiotic paper disc used were ampicillin, kanamycin, streptomycin, doxycycline and trimethoprim sulfamethoxazole. Escherichia coli ATCC 8739 was used as a control. One to two colonies of bacteria were put into 3 ml bouillon, incubated at 37°C for 2-6 hours then the culture turbidity was adjusted to McFarland 0.5. Following this, the culture were spread on to the surface of Mueller Hinton Agar plate and antibiotic paper discs were put on the surface of it and incubated at 35°C for 18-24 hours. The diameter of inhibition zone of each antibiotic was then recorded.

#### III. RESULTS AND ANALYSIS

#### Isolation and Primary Test of Haemophilus sp isolate

Five (2.5%) of the 197 samples were suspected Haemophilus sp based on their morphological characteristics on SBA. The five samples comes from 3 pigs from Tabanan (2 tracheal swabs, BIIT50 and BIIT53 and one lung sample BIIT1) and 2 pigs (two nasal swabs, BIBd1 and BIBd2) from Badung, respectively. Characteristics of the isolate were transparent tiny nonhemolytic satellite colonies growing near the Staph. aureus streak as shown on Figure 1. All five isolates were Gram positive rods, catalase positive, however the isolates from Tabanan (BIIT50, BIIT53, and BIIT1) were oxidase negative whereas those two (BIBd1 and BIBd2) from Badung were oxidase positive.



Fig. 1. Suspected colonies of Haemophilus sp (b) growing close to (a) Staphylococcus aureus streak colonies.

#### Biochemically Identification of Haemophilus sp.

Based on the biochemical confirmation test using API®20E<sup>TM</sup> two isolates BIIT1 and BIBd2 were identified as Enterobacter cloacae, BIIT53 as Proteus mirabilis, whereas BIIT50 and BIBd1 did not belong to a specific bacterial species. However, when comparing these two isolates physical and biochemical characteristics with H. parasuis, H. pleuropneumoniae, H. haemolyticus and H. influenza, one isolate, BIIT50 showed close characteristic to H. parasuis (see Table 1).Generally the occurrence of Glasser's disease in pig farms with characteristics syndrome is low. One case of the disease was reported by Priadi et al., (2006) occurred in pig farm in Pulau Bulan where it affected 10 weeks old post weaning pigs. The animals had pneumonia and died, lung and synovial fluids were sampled and Haemophilus parasuis serotype 12 was isolated.

TABLE 1.

PHYSIOLOGICAL AND BIOCHEMICAL CHARACTERISTICS OF TWO ISOLATES (BIIT50 AND BIBD1) IN COMPARISON TO HAEMOPHILUS PARASUIS, HAEMOPHILUS PLEUROPNEUMONIA, HAEMOPHILUS HAEMOLYTICUS, ANDHAEMOPHILUS INFLUENZA.

H.para Characteristics H.pleur H.hae H.inf BIB BIIT50 criteria opneum molytic luenz d1 suis onia us ae Growth factor + + + + + + Catalase + + + Oxidase + CO<sub>2</sub>(7-15%) + + Indole d d Hemolytic + Growth onMacCon keyAgar Sucrose + + Lactose d d + ONPG d +ADH d LDC \_ ? ODC ? d  $H_2S$ d Urease d Glucose MAN + Rhamnose Melibiose d Arabinose d Mannitol Sorbitol Inositol d Sucrose +Maltose \*Adapted fromMacedoet al., 2014; d = >90% negative

Similarly, in other country the prevalence of Glasser's disease in pig farms appeared to be low. Smart et al., (1989) reported a prevalence of 1.0% in Southern Ontario following 17,332 pigs reported sick [11]. Apparently Haemophilus sp is a fastidious fragile organism which is very sensitive to changing in humidity and temperature. Therefore it is very hard to isolate the bacteria. Moreover often the disease is in combination with other diseases in the respiratory tract of the pig [12].

#### Antimicrobial Sensitivity Test

Results of the sensitivity of the isolate towards several antibiotics tested is shown in Table 2 and Figure 2. As shown in Table 2 isolate BIIT50 which is suspected as H. parasuis appeared to be resistant to streptomycin only, whereas H. parasuis reported by Priadi et al. (2004) Nedbalcova et al. (2006) showed resistance to streptomycin, ampicillin and doxycycline. The differences in antibiotic sensitivity pattern is influenced by the improper and long term use of antibiotics which resulted in resistance strain [13]. Our isolate showed sensitive to ampicillin, kanamycin and doxycycline, respectively. There are two ways of bacterial resistance to antibiotics; by chromosomal and extra-chromosomal. Chromosomal resistance is genetically, occurs spontaneous and very fast, whereas extrachromosomal resistance occurs by the role of R-factor of bacterial plasmid [14][15].

TABLE 2.

DIAMETER OF INHIBITION ZONE OF ESCHERICHIA COLI ATCC 8739, BIIT50 AND BIBD1 TOWARDS AMPICILLIN, KANAMYCIN, DOXYCYCLINE, STREPTOMYCIN, AND TRIMETHOPRIM-SUL FAMETHOY AZOL F

SULFAMETHOXAZOLE.					
	Diameter of Inhibition Zone (mm)				(mm)
Isolates	Ampici	Kanam	Doxyc	Strepto	Trimethoprim-
	llin	ycin	ycline	mycin	Sulfamethoxazole
	(Amp)	(K)	(D)	(St)	(Sr)
Escheric	0(R)	21(S)	16(S)	16(S)	24(S)
hia coli					
ATCC87					
39					
BIIT50	20(S)	24(S)	30(S)	10(R)	15(I)
BIBd1	8(R)	19(S)	25(S)	13(I)	0(R)

R = Resistant; I = Intermediate; S =Sensitive;



Fig. 2. Antibiotic Sensitivity Test Plates of Escherichia coli ATCC 8739 (a); Isolate BIBd1 (b); and Isolate BIIT50 (c), Showing Diameter of Zone Inhibition on Mueller Hinton Agar.

#### IV. CONCLUSION

In conclusion we managed to isolate H. parasuis (one isolate) from pigs, where the isolate is resistant to streptomycin only and sensitive to ampicillin, kanamycin and doxycycline, respectively.

#### ACKNOWLEDGMENT

The authors would like to express their appreciation to The Ministry of Research & Technology and Higher Education through Udayana University Institute for Research and Community Service for supporting this project under Study Program Superior Grant 2015.

#### References

- Lawrence, P., Russell Bey. (2015). Map-Based Comparative Genomic Analysis of Virulent Haemophilus parasuis Serovar 4 and 5. Genom. 3: 59-71.
- [2] Bello-Ortí, B., Deslandes, V., Tremblay, Y.D., Labrie, J., Howell, K.J., Tucker, A.W., Maskell, D.J., Aragon, V., Jacques, M. (2014). Biofilm Formation by Virulent And Non-Virulent Strains of Haemophilus parasuis. Vete. Res. 45:104
- [3] Oliveira, P.S, Pérez-Simó, Aragón, V., Segalés J., Bensaid, A. (2011). Immunogenicity and Protection Against Haemophilus parasuis Infection after Vaccination with Recombinant Virulence Associated Trimetric Auto-transporters (VtaA). J. Vet. Sci. 14 (1): 111-116.
- [4] Macedo, N., Rovira, N., Torremorell, M., (2015).
  Haemophilus parasuis: Infection, Immunity and Enrofloxacin. Vet. Res. 46(128): 1-8.
- [5] Oliveira, S., (2015). Haemophilus Parasuis. Extention. http:// articles.extension. org/pages /27270 /haemophilus-parasuis.
- [6] Bochev, I. (2007). Porcine Respiratory Disease Complex (PRDC): A Review. I. Etiology, Epidemiology, Clinical Forms and Pathoanatomical Features. Bulg. J. Vet. Med. 10(3):131-146.
- [7] Priadi, A., Natalia, L., Poernomo, S. (2004).
  PenyakitGlasser's pada Babi di Pulau Batam, Propinsi Riau. BalaiPenelitianVeteriner, PO Box 151, Bogor 16114. JITV. 9(4).
- [8] Nedbalcova, K., Stran, P., Jaglic, Z., Ondriasova, R., Kucerova, Z. (2006). Haemophilusparasuis and Glasser's Disease in Pigs : a Review. Vet. Med. 51(5): 168-179.
- [9] Howell, K.J., Peters, S,E,, Wang, J., Garcia, J.H., Weinert, L.A., Luan, S.L., Chaudhuri, R.R., Angen,O., Aragon,V., Williamson, S.M., Parkhill, J., Langford, P.R., Rycroft,A.N., Wren, B.W., Maskell,D.J., W. Tucker,A.W., (2015). Development of a Multiplex PCR Assay for Rapid Molecular Serotyping of Haemophilus parasuis. J. Clin. Microbiol.53(12): 3812-3821.

- [10] Bagul, U.S., Sivakumar, S.M., (2016). Antibiotic Susceptibility Testing: A Review On Current Practices. Int. J. Phar. 6(3): 11-17.
- [11] Smart, N.L., Miniats, O.P, Rosendal S., Friendship, R.M. (1989). Glassers Disease and Prevalence of Subclinical Infection with Haemophilus parasuis in Swine in Southern Ontario. Can Vet. J. 30: 339-343.
- [12] Oliveira, S., Pijoan, C., Morrison, R. (2004). Evaluation of Haemophilus parasuis Control in Nursery Using Vacination and Controlled Exposure. J. Swine Health Prod.12(3): 123-128.
- [13] Rajurkar, G., Roy, A., Yadav. M.M., (2010). Antimicrobial Sensitivity Pattern of Haemophilus paragallinarum Isolated from Suspected Cases of

Infectious Coryza in Poultry. Vet. World, 3(4): 177-181.

- [14] Jain, A., Kumar, P., and Awasthi, S., (2005). High Nasopharyngeal Carriage of Drug Resistant Streptococcus pneumoniae and Haemophilus influenzae in North Indian School Children. Tro.l Med. Int. Health. 10(3): 234–239.
- [15] Veloo, A. C. M., Seme, K., Raangs, E., Rurenga, P., Singadji, Z., Wekema-Mulder, G., & van Winkelhoff, A. J. (2012). Antibiotic Susceptibility Profiles of Oral Pathogens. Int. J.Antimicrob.l Agents, 40(5): 450-454.

## Detection of Haemophilus Parasuis from The Respiratory Tract of Pigs

ORIGIN	ALITY REPORT	
% SIMILA	8 % 17 % 12 % 8 NTERNET SOURCES PUBLICATIONS STUDENT P	APERS
PRIMAR	Y SOURCES	
1	grjournals.com Internet Source	%1
2	www3.vf.uni-lj.si Internet Source	%1
3	KANG, Ikjae, Hwi Won SEO, Changhoon PARK, Yeonsu OH, Jeehoon LEE, Ok Heui YOU, Sung-Hoon KIM, Marcelo GOTTSCHALK, and Chanhee CHAE. "Digoxigenin-Labeled In Situ Hybridization for the Detection of Streptococcus suis DNA in Polyserositis and a Comparison with Biotinylated In Situ Hybridization", Journal of Veterinary Medical Science, 2014. Publication	% <b>1</b>
4	digital.library.adelaide.edu.au	%1
5	www.veterinaryresearch.org	%1
6	Sourav, S. Patricia, A. Sharma, S. Kanun. "Detection of pneumolysin and autolysin	%1

genes among antibiotic resistant Streptococcus pneumoniae in ", Indian Journal of Medical Microbiology, Jan-March 2010 Issue

Publication

7	medcraveonline.com Internet Source	%1
8	molecularbacteriology.com	%1
9	Huang, X., Y. Li, Y. Fu, Y. Ji, K. Lian, H. Zheng, J. Wei, X. Cai, and Q. Zhu. "Cross- Protective Efficacy of Recombinant Transferrin-Binding Protein A of Haemophilus parasuis in Guinea Pigs", Clinical and Vaccine Immunology, 2013. Publication	% <b>1</b>
10	www.wiloludjournal.com	%
11	portalgaruda.ilkom.unsri.ac.id	%1
12	libguides.niu.edu Internet Source	%
13	Shulin Fu, Lei Xu, Sali Li, Yinsheng Qiu, Yu Liu, Zhongyuan Wu, Chun Ye, Yongqing Hou, Chien-An Andy Hu. "Baicalin suppresses NLRP3 inflammasome and nuclear factor- kappa B (NF-κB) signaling during Haemophilus parasuis infection", Veterinary	% <b>1</b>

Research,	2016
-----------	------

Publication

14	Submitted to University of Queensland Student Paper	%1
15	www.rsu.ac.th Internet Source	%1
16	"Bergey's Manual® of Systematic Bacteriology", Springer Nature, 2012 Publication	%1
17	www.mysciencework.com	<‰1
18	www.affaveti.org	<‰1
19	veterinarymedicinejournal.usamv.ro	<‰1
20	file.scirp.org Internet Source	<‰1
21	doaj.org Internet Source	<‰1
22	M.S. Lorenson, M. Miani, J.A. Guizzo, B. Barasoul, S. Martínez-Martínez, E.F. Rodríguez-Ferri, C.B. Gutiérrez-Martín, L.C. Kreutz, R. Frandoloso. "Altered indirect hemagglutination method for easy serotyping	<% <b>1</b>

of Haemophilus parasuis", Arquivo Brasileiro

de Medicina Veterinária e Zootecnia, 2017 Publication



## 24 Bergey's Manual® of Systematic Bacteriology, 2005.

Publication

<%1

EXCLUDE QUOTES	ON	EXCLUDE MATCHES	OFF
EXCLUDE BIBLIOGRAPHY	OFF		