

ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 22 - Number 4 - April 2021

Front cover: Nycticebus javanicus É. Geoffroy, 1812
(PHOTO: ADENG BUSTOMI)

Published monthly

PRINTED IN INDONESIA

ISSN: 1412-033X

E-ISSN: 2085-4722



9 771412 033757



9 772085 472751

BIODIVERSITAS

Journal of Biological Diversity
Volume 22 – Number 4 – April 2021

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD:

Abdel Fattah N.A. Rabou (Palestine), **Agnieszka B. Najda** (Poland), **Ajay Kumar Gautam** (India), **Alan J. Lymbery** (Australia), **Annisa** (Indonesia), **Bambang H. Saharjo** (Indonesia), **Daiane H. Nunes** (Brazil), **Darlina Md. Naim** (Malaysia), **Ghulam Hassan Dar** (India), **Hassan Pourbabaei** (Iran), **Joko R. Witono** (Indonesia), **Kartika Dewi** (Indonesia), **Katsuhiko Kondo** (Japan), **Kusumadewi Sri Yulita** (Indonesia), **Livia Wannorp** (Sweden), **M. Jayakara Bhandary** (India), **Mahdi Reyahi-Khoram** (Iran), **Mahendra K. Rai** (India), **Mahesh K. Adhikari** (Nepal), **Maria Panitsa** (Greece), **Mochamad A. Soendjoto** (Indonesia), **Mohib Shah** (Pakistan), **Mohamed M.M. Najim** (Srilanka), **Nurhasanah** (Indonesia), **Praptiwi** (Indonesia), **Rasool B. Tareen** (Pakistan), **Sayed Aliakbar Hedayati** (Iran), **Sayed Mehdi Talebi** (Iran), **Shahabuddin** (Indonesia), **Shahir Shamsir** (Malaysia), **Shri Kant Tripathi** (India), **Subhash C. Santra** (India), **Sugeng Budiharta** (Indonesia), **Sugiyarto** (Indonesia), **Taufiq Purna Nugraha** (Indonesia), **Yosep S. Mau** (Indonesia)

EDITOR-IN-CHIEF:

S u t a r n o

EDITORIAL MEMBERS:

English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:

The Society for Indonesian Biodiversity

CO-PUBLISHER:

Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE:

biodiversitas.mipa.uns.ac.id; smujo.id/biodiv

.....



**Society for Indonesia
Biodiversity**



**Sebelas Maret University
Surakarta**

GUIDANCE FOR AUTHORS

Aims and Scope *Biodiversitas*, *Journal of Biological Diversity* or abbreviated as *Biodiversitas* encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of the gene, species, and ecosystem as well as ethnobiology.

Article types The journal seeks original full-length research papers, reviews, and short communication. Manuscript of original research should be written in no more than 8,000 words (including tables and picture), or proportional with articles in this publication number. Review articles will be accommodated, while, short communication should be written at least 2,000 words, except for pre-study.

Submission The journal only accepts online submission, through open journal system (<https://smujo.id/biodiv/about/submissions>) or email to the editors at unsjournals@gmail.com. Submitted manuscripts should be the original works of the author(s). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, a paragraph describing the claimed novelty of the findings versus current knowledge. Submission of a manuscript implies that the submitted work has not been published before (except as part of a thesis or report, or abstract); and is not being considered for publication elsewhere. When a manuscript written by a group, all authors should read and approve the final version of the submitted manuscript and its revision; and agree the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis; drafting of the manuscript and correcting of the revision. All authors must be responsible for the quality, accuracy, and ethics of the work.

Ethics Author(s) must obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If and when the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors or others are allowed to multiply article as long as not for commercial purposes. For the new invention, authors are suggested to manage its patent before published.

Open access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance The only articles written in English (U.S. English) are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers(double blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. The manuscript is rejected if the content does not in line with the journal scope, does not meet the standard quality, inappropriate format, complicated grammar, dishonesty (i.e. plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biodiversity significance. **Uncorrected proofs** will be sent to the corresponding author by email as .doc or .docx files for checking and correcting of typographical errors. To avoid delay in publication, corrected proofs should be returned in 7 days. The accepted papers will be published online in a chronological order at any time, but printed in the early of each month (12 times).

A charge Starting on January 1, 2019, publishing costs waiver is granted to authors of graduate students from **Least Developed Countries**, who first publish the manuscript in this journal. However, other authors are charged USD 250 (IDR 3,500,000). Additional charges may be billed for language editing, USD 75-150 (IDR 1,000,000-2,000,000).

Reprints The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by sending back the uncorrected proofs by email.

Manuscript preparation Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering size can be applied in presenting table and figure (9 pt). Word processing program or additional software can be used, however, it must be PC compatible and Microsoft Word based (.doc or .rtf, not .docx). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italic, except for italic sentence. Scientific name (genera, species, author), and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. Name of genera can be shortened after first mentioning, except generating confusion. Name of the author can be eliminated after first mentioning. For example, *Rhizopus oryzae* L. UICC 524, hereinafter can be written as *R. oryzae* UICC 524. Using trivial name should be avoided, otherwise generating confusion. **Biochemical and chemical nomenclature** should follow the order of the IUPAC - IUB. For DNA sequence, it is better used Courier New font. Symbols of standard chemical and abbreviation of chemistry name can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurement** use IS denomination, usage other system should follow the value of equivalent with the denomination of IS first mentioning. Abbreviations set of, like g, mg, mL, etc. do not follow by dot. Minus index (m⁻², L⁻¹, h⁻¹) suggested to be used, except in things like "per-plant" or "per-plot". **Equation of mathematics** does not always can be written

down in one column with text, in that case can be written separately. **Number** one to ten are expressed with words, except if it relates to measurement, while values above them written in number, except in early sentence. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent". Avoid expressing ideas with complicated sentence and verbiage, and used efficient and effective sentence.

Title of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Name of author(s) should be completely written. **Name and institution** address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscript written by a group, author for correspondence along with address is required. First page of the manuscript is used for writing above information.

Abstract should not be more than 200 words. **Keywords** is about five words, covering scientific and local name (if any), research theme, and special methods which used; and sorted from A to Z. All important **abbreviations** must be defined at their first mention. **Running title** is about five words. **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains for why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. **Concluding** sentence should be given at the end of the discussion. **Acknowledgments** are expressed in a brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest are noted.

Figures and Tables of maximum of three pages should be clearly presented. Title of a picture is written down below the picture, while title of a table is written above the table. Colored figures can only be accepted if the information in the manuscript can lose without those images; chart is preferred to use black and white images. Author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned source. **There is no appendix**, all data or data analysis are incorporated into Results and Discussions. For broad data, it can be displayed on the website as a supplement.

References Author-year citations are required. In the text give the authors name followed by the year of publication and arrange from oldest to newest and from A to Z. In citing an article written by two authors, both of them should be mentioned, however, for three and more authors only the first author is mentioned followed by et al., for example: Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation as shown with word "cit" should be avoided. Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in an alphabetical order (better, if only 20 for research papers). Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** (www.issn.org/2-22661-LTWA-online.php). The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning; a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158.

Book:

Rai MK, Carpinella C. 2006. Naturally Occurring Bioactive Compounds. Elsevier, Amsterdam.

Chapter in book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds) *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaad AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.) *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from internet:

Balagadde FK, Song H, Ozaki J, Collins CH, Barnet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4: 187. www.molecularsystemsbiology.com

The ethnobotany of *Ngusaba* ceremonial plant utilization by Tenganan Pegringsingan community in Karangasem, Bali, Indonesia

DEWA AYU SRI RATNANI^{1,✉}, I KETUT JUNITHA², ENIEK KRISWIYANTI², I NYOMAN DHANA³

¹Program of Biology Education, Faculty of Teacher Training and Education, Universitas Mahasaraswati Denpasar. Jl. Kamboja No. 11A, Denpasar 80233, Bali, Indonesia. Tel.: +62-361-227019, ✉email: sri.ratnani67@gmail.com

²Program of Biology, Faculty of Mathematics and Natural Sciences, Universitas Udayana. Jl. Raya Kampus Unud No. 9, Jimbaran, Badung 80361, Bali, Indonesia

³Program of Anthropology, Faculty of Humanities, Universitas Udayana. Jl. Nias No. 13, Sanglah, Denpasar 80114, Bali, Indonesia

Manuscript received: 22 March 2021. Revision accepted: 28 March 2021.

Abstract. Ratnani DA, Junitha IK, Kriswiyanti E, Dhana IN. 2021. The ethnobotany of *Ngusaba* ceremonial plant utilization by Tenganan Pegringsingan community in Karangasem, Bali, Indonesia. *Biodiversitas* 22: 2078-2087. Tenganan Pegringsingan is an ancient village in Bali, Indonesia, which often performs several ceremonies with high intensity. One of them is the *Ngusaba* ceremony, where many plants are utilized both in species and quantity. Hence, this study aimed to identify the species, family, local names, sources, and parts of plants, used for *Ngusaba* ceremonies by the Tenganan Pegringsingan community including the Index of Cultural Significance (ICS). Data analysis was qualitative and quantitative. Furthermore, the qualitative method was used to obtain data on the plants' local names, while snowball sampling was applied to select key informants through in-depth interviews and moderate participation. The results showed that the 130 species distributed in 56 families mostly belonging to the purchased source (34.61%). The Poaceae is the largest family, while the most widely used part of the plant is the leaf. Based on the ICS analysis results, a range of 2-114 values was obtained. The highest value is Base (*Piper betle* L.) and kangkung (*Ipomoea batatas* L.) as lowest.

Keywords: Ancient villagers, local knowledge, *Ngusaba* plant

INTRODUCTION

Bali is one of the tourism destinations in Indonesia has many attractions. Its distinctive feature is a unique blend of humans, nature, and culture, including customs and religious ceremonies where plants play an important role. Plants or their parts are the most important elements in material associated with the *Yadnya* ceremonies (Sujarwo 2020), including the *ngusaba* ceremony. The *Ngusaba* ceremony is a social activity to connect with the all mighty God (Ida Sang Hyang Widhi), which also includes banquets and *subak* village thanksgiving (Arwati 2007). It provides much information about the use of many plants or their parts, including leaves, flowers, fruits, seeds, and tubers (Adiputra 2011).

The utilization of *Ngusaba* by the Tenganan Pegringsingan community has some problems: which include (i) many of the ceremonial ingredients types and quantities needed exceed these plant's availability in nature; (ii) Only a few people are interested in traditional practices such as agriculture, because most of them rely on tenant farmers; (iii) The existence of plants, especially endemic species become increasingly hard to be found. Besides, plants are an important source of food, medicine, spice, construction materials, etc. in rural areas (Sujarwo et al. 2016; Sujarwo dan Caneva 2016; Sujarwo dan Keim 2017; Navia et al. 2020). They have many cultural sides, namely history, religion, language, art, politics, and social structure (Kakudidi 2004). They also have an important meaning, especially in various religious ceremonies

(Helida et al. 2015; Ristanto et al. 2020). Several plants are part of various ritual purposes (Sharma and Pegu 2011; Iskandar and Iskandar 2017) and a source of livelihood for the local people (Suwardi et al. 2020) that believe ritual is one of the most important instruments for understanding local communities and offering, to conserve nature (Geng et al. 2017). The conservation of plant resources is very important to combine with the understanding and awareness of local communities' cultural practices (Sheybani et al. 2015; O'Neill et al. 2017).

However, information technology development and modern lifestyle have led to a decline in local communities' traditional knowledge (Putri et al. 2017) and this condition also affects the Tenganan Pegringsingan community. In addition, the knowledge of ritual plant utilization is diminishing because it is only passed across generations orally and has remained unwritten (Anderson et al. 2011; Surata et al. 2015; Nisyapuri et al. 2018). The loss of local knowledge implicates plant resources' existence, as well as triggers disease and professional changes (Gomez et al. 2010; Cuadra et al. 2012; Ju et al. 2013; Vásquez et al. 2016; Aswani et al. 2018). The knowledge is very useful to conserve biodiversity, hence it needs to be maintained (Yusro et al. 2014) and documented for good management to halt the menace of biodiversity depletion (Adom 2018). There has been much effort in biodiversity conservation, such as plant preservation and documentation of their utilization through ethnobotany which is the study of utilitarian relationships between humans and plants in natural ecosystems and other social components (Hakim

2014). Ethnobotany data cover botany, taxonomy, and regional botanical knowledge. It is also essential for biodiversity conservation (Pieroni et al. 2014; Tapundu and Anam 2015), fulfillment of needs such as food, health, and culture (Setiawan and Qiptiyah 2014; Tamalene et al. 2016; Mesfin et al. 2018), construction, decoration, and other living necessities (Bosworth et al. 2011). Today, ethnobotany has become a crucial study area, which covers management resource development, biodiversity conservation at the genetic, species and ecosystem level, and regional socio-economic development (Caneva et al. 2017). Therefore, this study aims to identify plants used for *Ngusaba* ceremonies by the Tenganan Pegringsingan community.

MATERIALS AND METHODS

Study area

This study was conducted in Tenganan Pegringsingan community of Tenganan Village, Manggis Subdistrict, Karangasem District, Bali, Indonesia, from February to August 2020. The location is at positions 8000°.00' to 8041°.37.8' S and 115035° .9.8' to 115054° 8.9' E, at an altitude of 70-400 m asl. The village's temperatures ranging from 28-31°C.

General description of the study sites

Tenganan Pegringsingan is located in Manggis Subdistrict, Karangasem District, with a distance of ± 20 km from the District City, and ± 68 km from Denpasar. It is physiographically surrounded by three-quarters of a circle of hills forming borders in the north as Macang Village (*kaja* hill), east as Asak Village (*kangin* hill), and west as Ngis Village (*kauh* hill), but directly adjacent to Pesedahan Village in the south. According to usage the area includes paddy rice lands covering 255.85 ha, drylands covering 480.89 ha, and Adat forest lands covering 197.32 ha. (Monograph of Tenganan Village 2020). Tenganan Pegringsingan total population is 1022, with the family heads being 338, while the location map is shown in Figure 1.

Informant selection

Key informants were consulted with community leaders and selected using the snowball sampling technique, which was carried out in a chain by questioning those that have been interviewed or contacted previously (Hariyadi and Ticktin 2012). Furthermore, they had much information about the *Ngusaba* ceremony (Nurdiani 2014), including the offering expert, ceremony officials, and community leaders.




Figure 1. Map of the location of Tenganan Pegringsingan community (●) in Tenganan Village, Manggis Subdistrict, Karangasem District, Bali, Indonesia (Monograph of Tenganan Village 2020)

Data collection

Ethnobotany data were collected through semi-structured interviews and moderate participation in the form of species, family, local names, parts, sources, and the Index of Cultural Significance (ICS) of plants, which were analyzed qualitatively and quantitatively. A descriptive narrative was carried out for qualitative analysis through data reduction, display and analysis (Sugiyanto 2017). The quantitative analysis of the *Ngusaba* ceremonial plant was carried out through the ICS from Purwanto (2003). The ICS showed the importance values of each useful plant species based on the community's needs, and its calculation results showed each plant's importance level. The equation provided is to be employed to calculate ICS.

$$ICS = \sum_{i=1}^n (q \times i \times e) \times n_i$$

Because each species of plant has several uses, the equation is as follows:

$$ICS = \sum_{i=1}^n (q_1 \times i_1 \times e_1) \times n_1 + (q_2 \times i_2 \times e_2) \times n_2 + \dots + (q_n \times i_n \times e_n) \times n_i$$

Where:

ICS = the number of calculations the utilization of a plant species from 1 to n,

q : quality value calculated by giving a score or value on the quality value of a plant species: 3 = the main *Ngusaba* ceremony ingredient; 2 = additional *Ngusaba* ceremony materials + primary materials, 1 = other *Ngusaba* ceremony materials + secondary materials + primary materials

i : intensity value describes the intensity of utilization of useful plant species by giving values: value 3 = high intensity; 2 = moderate intensity; 1 = low intensity.

e : exclusivity value: value 2 = most important, is the first choice and is second to none; 1 = possibility of being a choice of secondary materials (Turner 1998; Purwanto 2003; modification of researchers).

The plants were collected with the informants and then identified by matching with the herbarium specimen of the Bali Botanical Garden, the picture on the flora book, and images on plantNet. Their scientific names were verified using online sources (e.g. The Plantlist 2019).

RESULTS AND DISCUSSION

Types of plants utilized for *Ngusaba* ceremony

The results showed 11 types of *Ngusaba* ceremonies carried out by the Tenganan Pegringsingan community, including *Ngusaba Kasa*, *Karo*, *Ketiga*, *Kelima* (*sambah*), *Kenem*, *Kepitu*, *Kaulu*, *Kesanga*, *Kedasa*, *Desta*, and *Sada*. The ceremonies are held almost monthly every year, and

each lasts for three days, except for *sambah* which lasts for one month. The *Ngusaba* plants in Tenganan Pegringsingan Village have a high diversity of 130 species belonging to 56 families among which the largest is Poaceae (16 species), followed by Fabaceae (9) and Musaceae (8). The percentage of the *ngusaba* plant families utilized by the community is shown in figure 2. The various species were collected from various habitats, mainly wild vegetation in the forest, roadsides, in front of the house, home gardens and drylands. The growth form indicated that the most widely used *ngusaba* are obtained from herbs (57 species or 43.84%), followed by trees (38 species or 29.23%), and shrubs (34 species or 26.15%) (Table 1).

This result is higher than 26 species representing 17 families found to be commonly used for performing the six main traditional rituals of the Karangwangi people (Erawan et al. 2018). The Baduy community uses 50 species representing 28 families for nine stages of their pure agricultural activity (Iskandar and Iskandar 2017), while the Aceh tribe in Peureulak uses 51 species consisting of 47 genera and 34 families (Sutrisno et al. 2020). Moreover, Bali Aga village uses 125 plant species for all the Panca yadnya ceremony (Sujarwo 2020) and based on these, cultural diversity shows biodiversity. The diversity of plants used for *Ngusaba* ceremonial offerings is an expression of the region's uniqueness which is a mountainous area surrounded by hills. A region's uniqueness determines biodiversity, including plants in a specific ecosystem. Each ethnic group grows according to regional uniqueness, culture, and natural resources' availability (Suryadarma 2017). Almost all the Poaceae family plants used for *Ngusaba* ceremonies are edible, staple foodstuffs, and the main agricultural product.

Plant parts utilized for *Ngusaba* ceremony by Tenganan Pegringsingan community

The plant parts used are in the form of leaf, stem, flower, fruit, seeds, tuber, and rhizome as presented in Figure 3. The most widely used are leaves, while the rhizome is the lowest.

The most utilized parts reported were leaves (45.52%), followed by fruits (38.80 %) and flowers (17.91%). Other studies such as Mesfin et al. (2013), Riadi et al. (2019), and Ristanto et al. (2020) also reported that leaves were the most commonly used. The high utilization of *Ngusaba* leaves appears to be associated with several advantages such as higher number or productivity of leaves that are easier to obtain than the other parts (Handayani 2015). *Piper betle* L. leaves are mostly utilized in all types of *Ngusaba* ceremonies. These are made in various forms that differentiate their names and are also irreplaceable (exclusive) and a must have in every offering. Furthermore, banana shoots are used almost equally as *Piper betle* L. and those having leaves that are useful to local people are included in a taste of sepia banana group, where the most widely used is *Musa acuminata* L. (*biyu keladi*).

Table 1. Species of *Ngusaba* plants utilization by Tenganan Pegringsingan community

Family/scientific name	Local name	Plants part	Habitus	ICS value	Category
Acanthaceae					
<i>Asystasia gangetica</i> L.	Loja	Leaf	Herb	6	Low
<i>Graptophyllum pictum</i> L.	Temen	Leaf	Shrub	24	Moderate
<i>Justicia adhatoda</i> L.	Dausa	Leaf	Shrub	102	Very high
<i>Thunbergia erecta</i> Benth	Terom Pelung	Flower	Shrub	4	Very low
Achariaceae					
<i>Pangium edule</i> Reinw.	Pangi	Seed	Tree	20	Moderate
Agavaceae					
<i>Dracaena marginata</i> Lam.	Sumenek	Leaf	Tree	24	Moderate
Amaranthaceae					
<i>Celocia cristata</i> L.	Kenyiwaan	Flower	Herb	6	Low
Amaryllidaceae					
<i>Allium sativum</i> L.	Kesuna	Tuber	Herb	42	Moderate
<i>Allium cepa</i> L.	Bawang	Tuber	Herb	30	Moderate
Anacardiaceae					
<i>Mangifera caesia</i> Jack.	Wani	Fruit	Tree	12	Low
<i>Mangifera indica</i> L.	Poh Arum Manis	Fruit	Tree	12	Low
<i>Mangifera indica</i> L.	Poh Madu	Fruit	Tree	12	Low
<i>Mangifera odorata</i> Griff.	Pakel	Leaf	Tree	6	Low
Annonaceae					
<i>Cananga odorata</i> Lamk.	Sandat	Flower	Tree	12	Low
Apocynaceae					
<i>Plumeria alba</i> L.	Jepun Bali	Flower	Tree	24	Moderate
<i>Plumeria alba</i> L.	Jepun Cenana	Flower	Tree	4	Very low
<i>Plumeria acuminata</i> L.	Jepun Merah	Flower	Tree	4	Very low
<i>Allamanda cathartica</i> L.	Kecubung Kuning	Flower	Shrub	4	Very low
Araceae					
<i>Colocasia esculenta</i> Schott.	Keladi	Leaf, tuber	Herb	12	Low
Araliaceae					
<i>Schefflera elliptica</i> (Blume) Harms.	Kayu Belang	Leaf	Shrub	20	Moderate
Arecaceae					
<i>Arenga pinnata</i> Merr	Jaka	Midrib, leaf, fruit	Tree	66	High
<i>Areca catechu</i>	Buah	Fruit, flower	Tree	66	High
<i>Cocos nucifera</i> L.	Nyuh Gadang	Midrib, leaf, fruit	Tree	84	High
<i>Cocos nucifera</i> L.	Nyuh Barak	Midrib, leaf, fruit	Tree	84	High
<i>Salacca zalacca</i> L.	Salak	Fruit	Tree	12	Low
Asclepiadaceae					
<i>Hoya australis</i> R.Br.ex.Trail.	Tebel-tebel	Leaf	Herb	6	Low
Asteraceae					
<i>Tagetes erecta</i> L.	Gumitir	Flower	Herb	4	Very low
<i>Tithonia aristrata</i> Oerst.	Sungenge	Flower	Herb	6	Low
Athyriaceae					
<i>Diplazium esculentum</i> (Retz.) Sw.	Paku Sayur	Leaf	Herb	24	Moderate
Bromeliaceae					
<i>Ananas comusus</i> Mer.	Manas	Fruit	Herb	12	Low
Cactaceae					
<i>Hylocereus polyrhizus</i> Britton&Rose	Buah Naga	Fruit	Herb	26	Moderate
Clusiaceae					
<i>Calophyllum inophyllum</i> L.	Camplung	Leaf	Tree	6	Low
<i>Mesua ferrea</i> L.	Nagasari	Leaf	Shrub	6	Low
<i>Garcinia mangostana</i> L.	Manggis	Fruit	Tree	12	Low
Combretaceae					
<i>Lummitzera littorea</i> Jack.	Padi-padi	Leaf	Tree	6	Low
Convolvulaceae					
<i>Ipomoea aquatica</i> Forssk.	Kangkung	Stem, leaf	Herb	2	Very low
Cucurbitaceae					
<i>Citrulus lanatus</i> (Thunb.)	Semangka	Fruit	Herb	24	Moderate
<i>Cucumis sativus</i> L.	Ketimun	Fruit	Herb	30	Moderate
Dioscoreaceae					
<i>Dioscorea bulbifera</i> L.	Ubiaung buluh	Tuber	Herb	6	Low

Euphorbiaceae					
<i>Aleurites moluccanus</i> L.	Tingkih	Seed	Tree	18	Low
<i>Codiaeum variegatum</i>	Kayu Mas	Leaf	Shrub	24	Moderate
<i>Phyllanthus boxifolius</i> Muell.Arg.	Kayu Sisi	Leaf	Shrub	24	Moderate
Fabaceae					
<i>Caesalpinia pulcherima</i> (L.) Sw.	Sumerak	Leaf, flower	Shrub	24	Moderate
<i>Casia glauca</i> Lamk.	Kembang Kuning	Leaf	Shrub	78	High
<i>Clitoria ternatea</i> L.	Teleng	Flower	Shrub	12	Low
<i>Erythrina abyssinica</i> Lam.	Dapdap	Leaf, stem	Tree	36	Moderate
<i>Psophocarpus tetragonolobus</i> L.	Kacang Botor	Seed	Herb	6	Low
<i>Tamarindus indica</i> Linn.	Cagi	Seed	Tree	6	Low
<i>Indigofera tinctoria</i> Mill.	Taum	Leaf	Shrub	6	Low
<i>Vigna unguiculata</i> L.	Kacang Barak	Seed	Herb	30	Moderate
<i>Vigna radiata</i> L.	Kacang Ijo	Seed	Shrub	6	Low
Heliconiaceae					
<i>Heliconia stricta</i> Huber.	Pisang Ikik	Leaf, fruit	Herb	6	Low
Hydrangeaceae					
<i>Hydrangea macrophylla</i> L.	Bunga Biru	Flower	Shrub	4	Very low
Lamiaceae					
<i>Ocimum gratissimum</i> L.	Sulasih	Leaf	Shrub	6	Low
Leeaceae					
<i>Leea aculeata</i> Burm.f.	Girang	Leaf, stem	Shrub	4	Very low
<i>Leea indica</i> Burm.f.	Kelawasan	Leaf	Shrub	24	Moderate
Leguminosae					
<i>Bauhinia purpurea</i> L.	Badya	Leaf	Tree	6	Low
<i>Indigofera tinctoria</i> Mill.	Kumaligi	Leaf	Herb	24	Moderate
<i>Mucuna pruriens</i> Wilmot.	Juleh	Seed	Herb	6	Low
<i>Lablab purpureus</i> L.	Komak selem	Seed	Herb	12	Low
Lygodiaceae					
<i>Lygodium circinatum</i> (Burm.f) Sw.	Ata	Stem+leaf	Herb	36	Moderate
Liliaceae					
<i>Cordyline fruticosa</i> L.	Andong	Leaf	Herb	30	Moderate
Limnocharitaceae					
<i>Limnocharis flava</i> L.	Biyah	Leaf	Herb	6	Low
Malvaceae					
<i>Durio zibethinus</i> L.	Duren	Fruit, leaf	Tree	12	Low
<i>Hibiscus rosasinensis</i> L.	Pucuk Bang	Flower	Shrub	84	High
Magnoliaceae					
<i>Michelia champaca</i> L.	Cempaka	Flower	Shrub	12t	Low
Marantaceae					
<i>Maranta ramosissima</i> Wall.	Kecandik	Leaf	Herb	6	Low
Meliaceae					
<i>Aglaiia odorata</i> Lour.	Kiulan	Flower	Shrub	4	Very low
<i>Azadirachta indica</i> Juss	Apah	Leaf	Tree	30	Moderate
<i>Lansium domesticum</i> L.	Langsat	Fruit	Tree	4	Very low
Moraceae					
<i>Ficus benyamina</i> L.	Bingin	Leaf	Tree	24	Moderate
<i>Ficus religiosa</i> L.	Ancak	Leaf	Tree	8	Low
Musaceae					
<i>Musa paradisiaca</i> L.	Biyu Gedang Saba	Leaf, fruit	Herb	30	Moderate
<i>Musa acuminata</i> L.	Biyu Ketip Tulang	Leaf, fruit	Herb	30	Moderate
<i>Musa acuminata</i> L.	Biyu Kunti	Leaf, fruit	Herb	84	High
<i>Musa acuminata</i> L.	Biyu Kayu	Leaf, fruit	Herb	48	Moderate
<i>Musa acuminata</i> L.	Biyu Bunga	Leaf fruit	Herb	54	High
<i>Musa acuminata</i> L.	Biyu Keladi	Leaf, fruit	Herb	99	High
<i>Musa acuminata</i> L.	Biyu Dak Sangket	Leaf, fruit	Herb	84	High
<i>Musa acuminata colla</i>	Biyu Alas	Fruit	Herb	32	Moderate
Myrtaceae					
<i>Psidium guajava</i> L.	Nyambu Kristal	Fruit	Shrub	4	Very low
<i>Syzygium polyanthum</i> Walp.	Don Juwet	Leaf	Tree	4	Very low
Nyctaginaceae					
<i>Bougenvillea spectabilis</i> L.	Bunga Kertas	Flower	Shrub	4	Very low
<i>Pisonia alba</i> Span.	Dagdag See	Leaf	Shrub	6	Low
Oleaceae					
<i>Nyctanthes arbotristis</i> L.	Srigading	Flower	Shrub	102	Very high

Oxalidaceae					
<i>Averrhoa carambola</i> L.	Belimbing Sayur	Leaf, fruit	Tree	6	Low
Pandanaceae					
<i>Pandanus amiryllicolius</i> Roxb.	Pandan Arum	Leaf	Shrub	8	Low
<i>Pandanus tectorius</i> Parkinson ex Du Roi	Pandan Duri	Leaf	Shrub	12	Low
Pinaceae					
<i>Pinus merkusii</i> Jungh.& de Vriese	Cemara	Leaf	Tree	6	Low
Piperaceae					
<i>Piper betle</i> L. var. <i>nigra</i>	Base Bali	Leaf	Herb	24	Moderate
<i>Piper betle</i> L.	Base biasa	Leaf	Herb	114	Very high
<i>Piper retrofractum</i> Vahl.	Tabia Bun	Fruit	Herb	24	Moderate
Poaceae					
<i>Brachiaria mutica</i> (Forssk.) Stapf.	Padang Guwun	Stem+leaf	Herb	18	Low
<i>Coix lacryma jobi</i> L.	Jali-Jali	Fruit	Herb	6	Low
<i>Cymbopogon citratus</i> DC	See	Stem	Herb	4	Very low
<i>Gigantochloa apus</i> (Schult.) Kurz	Tiying Tali	Stem	Tree	8	Low
<i>Hordeum scalinum</i> Schreb.	Ikuh bojog	Flower	Herb	12	Low
<i>Imperata cylindrica</i> L.	Ambengan	Leaf	Herb	18	Low
<i>Oryza sativa</i> L.	Beras	Seed	Herb	108	Very high
<i>Oryza nivara</i> L.	Beras merah	Fruit, seed	Herb	39	Moderate
<i>Oryza sativa</i> L. var. <i>glutinosa</i>	Ketan Putih	Fruit, seed	Herb	70	High
<i>Oryza sativa</i> L. var. <i>glutinosa</i>	Ketan barak	Fruit, seed	Herb	75	High
<i>Oryza sativa</i> L. var. <i>glutinosa</i>	Injin	Fruit, seed	Herb	75	High
<i>Oryza sativa</i> L.	Padi Gaga	Fruit	Herb	12	Low
<i>Oryza sativa</i> L.	Padi Bali	Fruit, seed	Herb	18	Low
<i>Saccharum officinarum</i> L.	Tebu Guwak	Stem	Herb	6	Low
<i>Sorghum bicolor</i> L.	Jagung Beleleng	Seed	Herb	6	Low
<i>Zea mays</i> L.	Jagung	Seed	Herb	6	Low
Pteridaceae					
<i>Adiantum pedatum</i> L.	Paku condong	Leaf	Herb	6	Low
Rubiaceae					
<i>Gardenia jasminoides</i> J.Ellis	Jempiring	Flower	Shrub	4	Very low
<i>Ixora coccinea</i> L.	Jaum-Jaum	Flower	Shrub	6	Low
<i>Psychotria micrantha</i> Kunth.	Wisnu	Leaf	Shrub	6	Low
Rosaceae					
<i>Malus domestica</i> Borkh.	Apel	Fruit	Tree	20	Moderate
<i>Pyrus communis</i> L.	Pir	Fruit	Tree	22	Moderate
Rutaceae					
<i>Citrus amblycarpa</i> Hassk	Limo	Fruit, leaf	Shrub	4	Very low
<i>Citrus grandis</i> L.	Jerungga	Fruit	Tree	12	Low
<i>Citrus reticulata</i> Blanco	Sumaga	Fruit	Shrub	24	Moderate
<i>Citrus sinensis</i> L.	Juuk	Fruit	Tree	24	Moderate
<i>Murraya paniculate</i> L.	Kemoning	Leaf	Shrub	24	Moderate
Santalaceae					
<i>Santalum album</i> L.	Cenana	Stem	Tree	6	Low
Sapindaceae					
<i>Cardiospermum halicacabum</i> Linn.	Kesuman Jai	Leaf	Herb	6	Low
<i>Nephelium lappaceum</i> L.	Buluan	Fruit	Tree	18	Low
Sapotaceae					
<i>Manilkara zapota</i> L.	Sabo	Fruit	Tree	4	Very low
Solanaceae					
<i>Solanum melongena</i> L.	Tuwung	Fruit	Shrub	6	Low
Urticaceae					
<i>Laportea stimulans</i>	Lateng Kebo	Leaf	Herb	6	Low
Vitaceae					
<i>Vitis vinifera</i> L.	Anggur	Fruit	Shrub	4	Very low
Zingiberaceae					
<i>Alpinia galanga</i> L.	Langkuas	Rhizome	Herb	57	High
<i>Curcuma longa</i> Linn.	Kunyit	Rhizome	Herb	18	Low
<i>Zingiber officinale</i> Rosc.	Jahe	Rhizome	Herb	30	Moderate




Figure 2. *Ngusaba* ceremony plant family used by Tenganan Pegringsingan community, Karangasem District, Bali, Indonesia




Figure 3. Parts of plant used for *Ngusaba* ceremony by Tenganan Pegringsingan community, Karangasem District, Bali, Indonesia




Figure 4. Sources of *Ngusaba* ceremonial plants by Tenganan Pegringsingan community, Karangasem District, Bali, Indonesia

Source of *Ngusaba* ceremonial plants

Ngusaba sources include the home gardens, in front of the house (*telajakan*), drylands, forests, roadsides, temples, other villages, and from purchase (Figure 4).

Thus, the purchase is the highest source (34.61%), followed by forest (16.92%) and *telajakan* (16.92%), and most of them are wild. This is in line with Sujarwo (2020) stated that most of the *Panca Yadnya* ceremonial plants in *Bali Aga Village* come from Balinese wild ethnoflora (Constant et al. 2018), and they are mostly distributed in various habitats. Their availability varies from one place to another among species. The majority used were harvested from the wild (35.38%), followed by semi-wild (23.84%), and cultivated (7.69 %). The community's efforts by planting in the settlements' vicinity including *telajakan*, drylands, and home gardens. However, there are many types and quantities of ceremonial ingredients needed that exceed this plant's availability in nature. Some of them have not been found in the Tenganan Pegringsingan Village, such as *Musa acuminata* L. (*biyu kunti*), *Musa acuminata* L. (*biyu kayu*), *Oryza nivara*, *Oryza sativa* var. *glutinosa* (red and black), *Citrus grandis* L., and *Hordeum scalinum* Schreb.

Index of Cultural Significance of useful plants (ICS)

The ICS calculation results showed various values with a range of 2-114. *Piper betle* L. has the highest (114), while the lowest value is noted for *Ipomoea aquatica* Forssk, and the ICS value categories (Figure 5).

The highest ICS value is noted for the plant species widely used by the Tenganan Pegringsingan community, especially those with high exclusivity and intensity levels. In fact, the intensity value is high because it is used in all *Ngusaba* ceremonies as a staple ingredient and is irreplaceable. Plants with more benefits often have a higher ICS value, which means to be more valuable and more exclusive (Hager 2008). The people of Tenganan Pegringsingan placed *Base* (*Piper betle* L.) plants at the highest level and as the most useful and valuable.




Figure 5. The percentage of the *Ngusaba* ceremony plants used by Tenganan Pegringsingan community, Karangasem District, Bali, Indonesia

This result showed that the Tenganan Pegringsingan community has the most interaction with the *Piper betle* L., meaning that this plant species will be used continuously in as much it is in line with the local community's cultural development. The variety of beneficial plants to a community group highly determines the conservation efforts made. The ICS results of Useful Plants as a quantitative ethnobotany analysis showed each useful plant species importance based on community needs (Munawaroh et al. 2011), hence determining the ones to be preserved (Supiandi 2019). The ICS plants' high index indicates a conservation stimulus, such as nature, benefits, and community willingness towards making efforts to develop it. The Tenganan Pegringsingan community tends to provide species that are often conserved as they are typical and cannot be replaced by other plants. In the beginning, local village communities made use of their natural resources and environment primarily based on local knowledge and/or beliefs embedded in their culture (Iskandar 2016). Therefore, human culture can be understood as the knowledge that contains several sets of models used effectively to interpret, understand, and guide behavior in adapting to the environment (Ahimsa-Putra 2012). This situation requires thoughts and efforts on plant

reintroduction, which the community continuously utilizes by creating a *Ngusaba* ceremonial plants' garden.

In conclusion, we indicated that Tenganan Pegringsingan people utilize a large number of plant species (130) named and explained for *Ngusaba* ceremonies. The plant's largest family (16) is Poaceae, while the most widely used part is the leaf, and the highest proportion was obtained by purchasing. Even though most of them are harvested from the wild vegetation, areas such as roadsides, forests, and dry lands are exposed to many threats. The Index of Cultural Significance of the *Ngusaba* in Tenganan Pegringsingan Village ranges from 2 to 114, and *Piper betle* L. has the highest value (114). There is an imbalance between their existence and the use of plants by the community. Many of them become increasingly hard to be found and some have not been found in Tenganan Pegringsingan, such as *Musa acuminata* L. (*biyu kunti*), *Musa acuminata* L. (*biyu kayu*), *Oryza nivara*, *Oryza sativa* var. *glutinosa* (red and black), *Citrus grandis* L., *Limncharis flava* L., *Pinus merkusii* L., and *Hordeum scalinum* Schreb. Therefore, urgent efforts on plant reintroduction are needed to be continuously utilized by the community by creating a *Ngusaba* ceremonial plants' garden.

ACKNOWLEDGEMENTS

The authors are grateful to all people in the studied village for their kind hospitality, the share of knowledge on plants and provision of the opportunity and facilities needed in Tenganan Pegringsingan, Manggis Subdistrict, Karangasem Bali, Indonesia.

REFERENCES

- Adiputra N. 2011. Medicinal plants, ceremonial plants and environmental conservation. *Bumi Lestari J Environ* 11 (2): 346-354. [Indonesian]
- Adom D. 2018. Traditional cosmology and nature conservation at the Bomfobiri Wildlife Sanctuary of Ghana. *Nat Conserv Res* 3 (1): 35-57. DOI: 10.24189/ncr.2018.005
- Ahimsa-Putra HS. 2012. Phenomenology approach to understanding religion. *Walisongo* 20 (2): 271-304. DOI: 10.21580/ws.20.2.200 [Indonesian]
- Anderson MJ, Christ TO, Chase JM, Vellend M, Brian D, Inouye, Freestone AL, Sanders NJ, Cornell HV, Comita LS, Davies KF, Harrison SP, Kraft NJB, Stegen JC, Swenson NG. 2011. Navigating the multiple meanings of diversity: A roadmap for ecologists. *Ecol Lett* 14 (1): 19-28. DOI: 10.1111/j.1461-0248.2010.01552.x.
- Arwati M. 2007. *Ngusaba* Ceremony. Office of Religion of Bali Province, Denpasar. [Indonesian]
- Aswani S, Lemahieu A, Sauer WHH. 2018. Global trends of local ecological knowledge and future implications. *PLoS ONE* 13 (4): e0195440. DOI: 10.1371/journal.pone.0195440
- Bosworth K, Lysbeth F, Diley HMA. 2011. School climate factors contributing to student and faculty perceptions of safety in select Arizona schools. *J School Health* 81 (4): 194-201. DOI: 10.1111/j.1746-1561.2010.00579.x
- Caneva G, Traversetti L, Sujarwo W, Zuccarello V. 2017. Sharing ethnobotanical knowledge in traditional villages: Evidence of food and nutraceutical B core groups in Bali, Indonesia. *J Econ Bot* 71 (4): 303-313. DOI: 10.1007/s12231-017-9395-x.
- Constant NL, Tshisikhawe MP. 2018. Hierarchies of knowledge: Ethnobotanical knowledge, practices and beliefs of the Vhavenda in South Africa for biodiversity conservation. *J Ethnobiol Ethnomed* 14 (56): 2-28. DOI: 10.1186/s13002-018-0255-2.
- Cuadra VP, Cambi V, Ruda MA. 2012. Consequences of the loss of traditional knowledge: The risk of injurious and toxic plants growing in kindergartens. *Ethnobot Res App* 10: 77-94.
- Erawan TS, Alillah AN, Iskandar J. 2018. Ethnobotany of traditional rituals in the Karangwani Village Cianjur District, West Java, Indonesia. *Asian J Ethnobiol* 1 (2): 52-60. DOI: 10.13057/asianjethnobiol/y010201.
- Geng Y, Hu G, Ranjitkar S, Shi X, Zhang Y, Wang Y. 2017. The implications of ritual practices and ritual plant use on nature conservation: a case study among the Naxi in Yunnan Province, Southwest China. *J Ethnobiol Ethnomed* 13 (1): 1-11. DOI: 10.1186/s13002-017-0186-3.
- Gomez BE, Mingoria S Reyes GV, Calvet L Montes C. 2010. Traditional ecological knowledge trends in the transition to a market economy: Empirical study in the Don Ana Natural Areas. *Conserv Biol* 24 (3): 721-729. DOI: 10.1111/j.1523-1739.2009.01401.x.
- Hager TF. 2008. Processing and storage effects on monomeric anthocyanins, percent polymeric color, and antioxidant capacity of processed blackberry products. *J Agric Food Chem* 56 (3): 689-695. DOI: 10.1021/jf071994g.
- Hakim L. 2014. Ethnobotany and home garden management: Food security, health, and agro-tourism. Selaras, Malang. [Indonesian]
- Handayani A. 2015. Utilization of medical plants by people around Mount Simpang Nature Reserve, West Java. *Pros Sem Nas Masy Biodiv Indon* 1 (6): 1425-1432. [Indonesian]
- Hariyadi B, Ticktin T. 2012. Uras: Medicinal and ritual plants of Serampas, Jambi Indonesia. *J Ethnobot Res Appl* 10: 133-149.
- Helida A, Zuhud EAM, Hardjanto, Purwanto, Hikmat A. 2015. Index of cultural significance as a potential tool for conservation of plants diversity by communities in The Kerinci Seblat National Park. *Jurnal Manajemen Hutan Tropika* 21 (3): 192-201. DOI: 10.7226/jtfm.21.3.192. [Indonesian]
- Iskandar J. 2016. Ethnobiology and cultural biodiversity in Indonesia. *Indones J Anthropol* 1 (1): 27-42.
- Iskandar J, Iskandar BS. 2017. Various plants of traditional rituals: Ethnobotanical research among Baduy community. *Biosaintifika* 9 (1): 114-125. DOI: 10.15294/biosaintifika.v9i1.8117. [Indonesian]
- Ju Y, Zhuo J, Lui B, Long C. 2013. Eating from the wild: Diversity of wild edible plants used by Tibetans in Shangrila-Region, Yunnan, China. *J Ethnobiol Ethnomed* 9 (1): 1-22. DOI: 10.1186/1746-4269-9-28
- Kakudidi EK. 2004. Cultural and social uses of plants from and around Kibale National Park, Western Uganda. *Afr J Ecol* 42 (1): 114-118. DOI: 10.1111/j.1365-2028.2004.00472.x.
- Mesfin K, Tekle G, Tesfay T. 2013. Ethnobotanical study of traditional medicinal plants used by indigenous people of Gemad District, Northern Ethiopia. *J Med Plants Stud* 1 (4): 32-37.
- Monograph of Tenganan Village, Karangasem. 2020.
- Munawaroh E, Saparita R, Purwanto Y. 2011. Community dependence on non-timber forest products in Malinau, East Kalimantan: an ethnobotany analysis and its implications for forest conservation. *Berkala Hayati Special Edition* 7A: 51-58.
- Navia ZI, Audira D, Afifah N, Turnip K, Nuraini, Suwardi AB. 2020. Ethnobotanical investigation of spice and condiment plants used by the Taming tribe in Aceh, Indonesia. *Biodiversitas* 21 (10): 4467-4473. DOI: 10.13057/biodiv/d211001.
- Nisyapuri FF, Johan I, Ruhyat P. 2018. Ethnobotany of medicinal plants in Wonoharjo Village, Pangandaran Regency, West Java. *Biodiversity* 4 (2): 122-132. DOI: 10.13057/psnmbi/m040205.
- Nurdiani N. 2014. Snowball sampling techniques in dry land research. *Comtech* 5 (2): 1110-1118.
- O'Neill AR, Badola HK, Dhyani PP, Rana SK. 2017. Integrating ethnobiological knowledge into biodiversity conservation in the eastern Himalayas. *J Ethnobiol Ethnomed* 13 (1): 1-14. DOI: 10.1186/s13002-017-0148-9.
- Pieroni A, Anely N, Avni H, Mustafa B, Bruno S, Kevin C, Cassandra LQ. 2014. Local knowledge on plant and domestic remedies in the mountain village of Peshkopia (Eastern Albania). *J Mt Sci* 11 (1): 180-194. DOI: 10.1007/s11629-013-2651-3.
- Purwanto Y. 2003. Ethnobotany Research Methods. Lab. Ethnobotany. Balitbang Botani, LIPI Biology Research and Development Center, Bogor, Indonesia. [Indonesian]
- Putri NS, Dewi R, Fitriana. 2017. Ceremony bridal process in Teubang Phui Baru village, Montasik sub-district, Aceh Besar district. *Jurnal*

- Ilmiah Mahasiswa Pendidikan Kesejahteraan Keluarga 2 (4): 42-57. [Indonesian]
- Ristanto RH, Suryanda A, Rismayati AI, Rimadana A, Datau R. 2020. Ethnobotany: a plant of Hindu-Balinese religious rituals. *Jurnal Pendidikan Biologi* 5 (1): 96-105. DOI: 10.31932/jpbio.v5i1.642 [Indonesian]
- Setiawan H, Qiptiyah M. 2014. Ethnobotany study of the Moronene tribe in the Rawa Aopa Watumohai National Park. *Wallacea J Res* 3 (2): 107-117. DOI: 10.18330/jwallacea.2014.vol3iss2pp107-117.
- Sharma UK, Pegu S. 2011. Ethnobotany of religious and supernatural beliefs of the Mising tribes of Assam with special reference to the 'Dobur Uje'. *J Ethnobiol Ethnomed* 7 (1): 1-13. DOI: 10.1186/1746-4269-7-16.
- Sheybani H, Charmchian M, Azadboni ZB. 2015. Factors affecting villagers' participation in forest conservation in the region of Miyandoroud. *Indian J Fundam Appl Life Sci* 5 (S1): 3724-3730.
- Sujarwo W, Caneva G. 2016. Using quantitative indices to evaluate the cultural importance of food and nutraceutical plants: Comparative data from the Island of Bali (Indonesia). *J Cul Her* 18: 342-348. DOI: 10.1016/j.culher.2015.06.006.
- Sujarwo W, Keim AP, Caneva G, Tonia C, Nicoletti M. 2016. Ethnobotanical uses of neem (*Acadiractha Indica* A. Juss.; Meliaceae) leaves in Bali (Indonesia) and the Indian subcontinent in relation with historical background and phytochemical properties. *J Ethnopharmacol* 189: 186-193. DOI: 10.1016/j.jep.2016.05.014.
- Sujarwo W, Keim AP. 2017. Ethnobotanical study of traditional building materials from the Island of Bali, Indonesia. *Econ Bot* 71 (3): 224-240. DOI: 10.1007/s12231-017-9385-z.
- Sujarwo W, Caneva G, Zuccarello V. 2020. Patterns of plant use in religious offerings in Bali (Indonesia). *Acta Bot Bras* 34 (1): 40-53. DOI: 10.1590/0102-33062019abb0110.
- Supiandi MI, Mahanal S, Zubaidah S, Julung HBEGE. 2019. Ethnobotany of traditional medicinal plants used by Dayak Desa community in Sintang, West Kalimantan, Indonesia. *Biodiversitas* 20 (5): 1264-1270. DOI: 10.13057/biodiv/d200516.
- Surata IK, Gata IW, Sudiana IM. 2015. Ethnobotanical study of Balinese Hindu ceremonial plants as an effort to protect local wisdom. *Jurnal Kajian Bali* 5 (2): 265-284. [Indonesian]
- Suryadarma IGP. 2017. Building the character of independence of an ethnobiological approach to local uniqueness in national perspectives and global reflections. *Biotic National Seminar Proceedings*. Yogyakarta, November 26, 2017. [Indonesian]
- Sutrisno IH, Akob B, Navia ZI, Nuraini, Suwardi AB. 2020. Documentation of ritual plants used among the Aceh Tribe in Peureulak, East Aceh District, Indonesia. *Biodiversitas* 21 (11): 4990-4999. DOI: 10.13057/biodiv/d211102.
- Suwardi AB, Navia ZI, Harmawan T, Syamsuardi, Mukhtar E. 2020. Ethnobotany and conservation of indigenous edible fruit plants in South Aceh, Indonesia. *Biodiversitas* 21 (5): 1850-1860. DOI: 10.13057/biodiv/d210511.
- Tamalene MN, Al Mudhar MHI, Suarsini E, Rahman F, Hasan S. 2016. Ethnobotany of Canarium plant species used by Tobelo Dalam (Togutil) ethnic community of Halmahera Island, Indonesia. *Biodiversitas* 17 (1): 61-69. DOI: 10.13057/biodiv/d170109.
- Tapundu AS, Anam S. 2015. Studi etnobotani tumbuhan obat pada Suku Seko di Desa Tanah Harapan Kabupaten Sigi, Sulawesi Tengah. *Jurnal Biocelbes* 9 (2): 66-86. [Indonesian]
- Turner NJ. 1998. The importance of a rose: Evaluating the cultural significance of plants in Thompson and Lillooet Interior Salish. *J Amer Anthropol* 90 (2): 272-290. DOI: 10.1525/aa.1988.90.2.02a00020.
- Yusro, Fathul, Yeni M, Farah D, Kazuhiro. 2014. Inventory of medicinal plants for fever used by four Dayak Sub Ethnic in West Kalimantan, Indonesia. *Kuroshio Sci* 8 (1): 33-38.
- Vásquez A, Vibrans H, Vergara SF, Caballero J. 2016. Intracultural differences in local botanical knowledge and knowledge loss among the Mexican Isthmus Zapotecs. *PLoS ONE* 11 (3): e0151693. DOI: 10.1371/journal.pone.0151693.