




David Publishing Company
www.davidpublishing.com

ISSN 1934-7391 (Print) ISSN 1934-7405 (Online)

JLS

Journal of Life Sciences

Volume 7, Number 7, July 2013



From Knowledge to Wisdom

Journal of Life Sciences

Volume 7, Number 7, July 2013 (Serial Number 63)



David Publishing Company
www.davidpublishing.com

Publication Information

Journal of Life Sciences is published monthly in hard copy (ISSN 1934-7391) and online (ISSN 1934-7405) by David Publishing Company located at 3592 Rosemead Blvd #220, Rosemead, CA 91770, USA.

Aims and Scope

Journal of Life Sciences, a monthly professional academic journal, covers all sorts of researches on molecular biology, microbiology, botany, zoology, genetics, bioengineering, ecology, cytology, biochemistry, and biophysics, as well as other issues related to life sciences.

Editorial Board Members

Dr. Stefan Hershberger (USA), Dr. Suiyun Chen (China), Dr. Farzana Perveen (Pakistan), Dr. Francisco Torrens (Spain), Dr. Filipa João (Portugal), Dr. Masahiro Yoshida (Japan), Dr. Reyhan Erdogan (Turkey), Dr. Grzegorz Żurek (Poland), Dr. Ali Izadpanah (Canada), Dr. Barbara Wiewióra (Poland), Dr. Valery Lyubimov (Russia), Dr. Amanda de Moraes Narcizo (Brasil), Dr. Marinus Frederik Willem te Pas (The Netherlands), Dr. Anthony Luke Byrne (Australia), Dr. Xingjun Li (China), Dr. Stefania Staibano (Italy), Dr. Wenle Xia (USA), Hamed Khalilvandi-Behroozyar (Iran).

Manuscripts and correspondence are invited for publication. You can submit your papers via Web Submission, or E-mail to life-sciences@davidpublishing.org or life-sciences@hotmail.com. Submission guidelines and Web Submission system are available at <http://www.davidpublishing.org>.

Editorial Office

3592 Rosemead Blvd #220, Rosemead, CA 91770, USA
Tel: 1-323-9847526, 1-302-5977046; Fax: 1-323-9847374
E-mail: life-sciences@davidpublishing.org, life-sciences@hotmail.com

Copyright©2011 by David Publishing Company and individual contributors. All rights reserved. David Publishing Company holds the exclusive copyright of all the contents of this journal. In accordance with the international convention, no part of this journal may be reproduced or transmitted by any media or publishing organs (including various websites) without the written permission of the copyright holder. Otherwise, any conduct would be considered as the violation of the copyright. The contents of this journal are available for any citation. However, all the citations should be clearly indicated with the title of this journal, serial number and the name of the author.

Abstracted / Indexed in

Database of EBSCO, Massachusetts, USA
Chemical Abstracts Service (CAS), USA
Database of Cambridge Science Abstracts (CSA), USA
Database of Hein Online, New York, USA
Ulrich's Periodicals Directory, USA
Universe Digital Library S/B, Proquest
Chinese Database of CEPS, American Federal Computer Library center (OCLC), USA
China National Knowledge Infrastructure, CNKI, China
Chinese Scientific Journals Database, VIP Corporation, Chongqing, China
Index Copernicus, Index Copernicus International S.A., Poland
Google Scholar (scholar.google.com)

Subscription Information

Price (per year): Print \$420, Online \$300, Print and Online \$560.

David Publishing Company
3592 Rosemead Blvd #220, Rosemead, CA 91770, USA
Tel: 1-323-9847526, 1-302-5977046; Fax: 1-323-9847374
E-mail: order@davidpublishing.org



David Publishing Company
www.davidpublishing.org

JLS

Journal of Life Sciences

Volume 7, Number 7, July 2013 (Serial Number 63)

Contents

Biochemistry and Biotechnology

- 677 **Dicamba Growth Regulator Promotes Genotype Independent Somatic Embryogenesis from Immature Zygotic Embryos of Tropical Maize Inbred Lines**

Joseck Akoyi, Allan J. Mgtutu, Jesse Machuka, Mieke van Lijsebettens, Catherine Taracha and Sylvester E. Anami

- 690 **Comparative Study of the Static Magnetic Field Effects on Growth Rate with Relative Antibiotic Susceptibility in *Escherichia coli***

Fouad Houssein Kamel, Ashti M. Amin, Khonaw Kader Salih and Saleem S. Qader

- 695 **Use of Wood Fibre Compost for the Cultivation of *Trichoderma* sp. (Isolate Td₂₂)**

Yan Ramona and Martin A. Line

- 700 **Evaluation of the Insecticidal Activity of the Aerial Part of *Pseudocytisus integrifolius* (Salisb) Rehder on Grain Borer, *Rhyzopertha dominica* Fab. (Bostrychidae) and Wheat Weevil, *Sitophilus granarius* Linn. (Curculionidae)**

Kassem Naima, Khelil Mohamed Anouar and Bendimerad Nassima

- 705 **Structure - Antimicrobial Activity Relationship Investigation of Some Butadiene and Chalcone Derivatives**

Hanoy AL-Amood, Hadeel T. AL-Hadithi and Ghazwan F. Fadhil

Plant Sciences

- 712 **Influence of Agricultural Inputs on Growth and Yield of *Jatropha curcas* (L.) in Cameroon**

Tehobsala, Mégueni Clautilde, Njintang Yanou Nicolas, Nenwôla Kona Bilele, Patrick Prudent, Joseph Wey, Lyana Jean and Djonbada Pouimo

- 722 **Restoration of *Posidonia oceanica* (L.) Delile Meadows: Is There An Effective Methodology?**

Bedini Roberto, Nannelli Annalisa and Batistini Francesca

- 727 **The Strategies of Grassland Management in Farms of Northeastern Part of Poland**

Jankowski Kazimierz, Kolczarek Roman, Jankowska Jolanta and Sosnowski Jacek

- 732 **Improving the Size and Market Value of an Underutilised Yam (*Dioscorea esculenta*) in Ghana: Implications for Crop Breeding and Production Choices**

Kwamina Banson and Kenneth Danso

Public Health

- 742 **Adverse Events Clustering with NAT2 Slow Metabolisers following Deparasitization in Children in Bangolan, NWR Cameroon**

Olivia Afa Achonduh, Babara Atogho-Tiedeu, Innocent Ali Mbulli, Jean Paul Chedjou, Mercy Achu, Akindeh Mbu Nji, Fokou Elie, Eric Kamgue, Vera Veyee, Orise Karana, Delphine Sahfe and Wilfred Fon Mbacham

- 749 **Diagnosis Traps in Polyarteritis Nodosa — Original Case Report**

Stoicescu Manuela

- 754 **Body Weight Models in a Population-Based Study of Albanian School Adolescents**

Enkelejda Shkurti, Diamant Shtiza and Diederik Aarendonk

- 760 **Meeting the ADA Guidelines of Diabetic Care at King Fahd Hospital of University, Khobar, Eastern Province, Saudi Arabia in 2012**

Waleed AlBaker, Fatemah AlFaraj, Reem AlArgan and Ammar Khamis

Biogeography

- 766 **Comparative Data of Millipedes' Distribution in Southern Region of Albania**

Hajdar Kicaj and Mihallaq Qirjo

- 773 **Estimation of Artificial Plantings of *Pinus sylvestris* in Kazakhstan according to Their Geographical Origin**

Nadezhda Konstantinovna Chebotko, Vitaliy Yurevich Kirillov and Bolat Mazhitovich Mukanov

- 781 **An Analysis of Wetland Degradation in the Dimoria Region of Assam, India**

Dharma Ram Deka and Pradip Sharma

Use of Wood Fibre Compost for the Cultivation of *Trichoderma* sp. (Isolate Td₂₂)

Yan Ramona¹ and Martin A. Line²

1. Integrated Laboratory for Biosciences and Biotechnology, Udayana University, Bali 80362, Indonesia

2. School of Agricultural Science, University of Tasmania, Tasmania 7001, Australia

Received: February 16, 2013 / Accepted: May 15, 2013 / Published: July 30, 2013.

Abstract: The main objective of this research was to investigate the ability of a *Trichoderma* sp. (Td₂₂), inhibitory to *Sclerotinia minor* Jagger, to grow and survive in mature wood fibre waste (WFW) compost of paper mill origin following nutrient amendment. The growth and survival of the fungus in the WFW compost was assessed by serial dilution plate count method followed by confirmation of the fungal identity using pectic enzyme analysis as described in Cruickshank and Pitt [1]. It was found in this study that the population densities of Td₂₂ achieved under non-sterile conditions in the WFW compost following nutrient amendment was approximately in the range of 7.0 log₁₀ CFU/g dw – 8.5 log₁₀ CFU/g dw after 28 days, depending on pre-treatment. The efficacy of this WFW compost-grown Td₂₂ for protection of lettuce from attack by *S. minor* was also demonstrated in glasshouse trials. This study indicates that cellulosic paper mill waste compost could provide an abundant low-cost growth medium for the large-scale cultivation of fungal antagonists, improving prospects for cost-competitiveness with chemical treatments.

Key words: Compost, biological control, *Trichoderma* sp., *Sclerotinia minor*.

1. Introduction

High concentration of pesticide usage often leads to increased pathogen resistance, resulting in ever higher pesticide applications in order to kill the same pathogen [2-4]. Furthermore, pesticide residue can have adverse effects on reproduction and developmental processes in a variety of animals including humans [5-7]. Due to these detrimental effects, reduced pesticide usage in agricultural practice has been championed [8, 9] and one of the best broad spectrum pesticide/fumigants (methyl bromide) is no longer used in developed countries. Alternatives have been proposed, but none appears to be as effective [10]. In anticipation of such future reduction, Painuly and Dev [11] proposed some possible alternative control, such as biological control, use of pathogen-resistant plant varieties, and integrated pest management.

Problems related to the large-scale production of biological control agents in low cost materials include difficulties in handling, transport, and storage. These problems have been largely overcome by maintenance of bacterial antagonists in carriers such as peat or vermiculite for periods of five months or more [12-14]. Cost of the cultivation medium, its transport, and field dispersal are critical factors in any assessment of biological control relative to chemical treatments, a problem sometimes exacerbated by the perceived need for proprietary media formulations.

Norske-Skog Paper Mill Limited produces approximately 33,000 tonnes of wood fibre waste (WFW) per annum, to be dumped as landfill [15].

This cellulosic waste was considered worthy of examination in view of its attributes of excellent water-holding and/or aeration capacity, its freedom from toxic elements, its free availability, and its potential utilization as a source of energy and carbon by cellulose-utilizing fungi, including the *Trichoderma* sp.

Corresponding author: Yan Ramona, Ph.D., research fields: environmental microbiology, composting and biocontrol.
E-mail: yan_ramona@yahoo.com.

isolate Td₂₂.

Based on the above background, the potential use of wood fibre waste compost as a possible substrate for the low-cost cultivation of a fungal biological control agent (*Trichoderma* sp. isolate Td₂₂) was investigated, together with assessments of the efficacy of the compost-grown Td₂₂ to protect lettuce seedlings from attack by *S. minor* in pot trials or glasshouse level experiments.

2. Materials and Methods

2.1 Isolate Td₂₂

Trichoderma sp., isolate Td₂₂ was kindly provided by Dr. Dean Metcalf (DPIWE Tasmania). For long term storage, the fungus was maintained at 4 °C in sterilized and moist millet seeds.

2.2 Wood Fiber Waste

The WFW used in this study was the same as that used by Ramona and Line [15]. It is a mix of eucalypt and *Pinus radiata*, comprising holocellulose as its primary constituent with very low levels of metal contaminants. It was also deficient (from a recycling perspective) in N and P. In the present trial, composting of this material was for three months (followed the method specified by Jackson and Line [16] and Jackson [17]) until its toxicity to radish seeds was eliminated.

2.3 Inoculum Preparation

The active starters used in this experiment were prepared according to the methods as specified in Ramona [18]. The potency of this inoculum, measured in CFU, was determined by serial dilution plating (in triplicates) before being used to inoculate the WFW.

2.4 Assessment of Growth of the Td₂₂

The WFW compost (at a self-generated temperature of 55-60 °C) was air-dried in a glasshouse, mixed with 2% (w/v) Phostrogen® solution, and placed into 750 mL flasks (100 g quantities each). These flasks were

then either briefly autoclaved for 5 minutes prior to inoculation (Compost A), or inoculated directly with inoculum (Td₂₂ suspension described above) (Compost B) to give an initial density of 5.5 log₁₀ CFU/g compost. Incubation was at 25 °C for 4 weeks with periodic assay of CFU on Potato Dextrose Agar (PDA) for Td₂₂ and on 0.1% Trypticase Soya Agar (TSA) for check for bacterial contaminants. The identity of the Td₂₂ was confirmed by analysis of pectic enzymes using the method as described by Cruickshank and Pitt [1].

2.5 The Efficacy of WFW Compost-Grown Td₂₂

Lettuce seeds were sown in pots containing field soil amended with suppressive compost (Compost A) at levels of 2.5, 5.0, 10 and 20% v/v.

The spore density of the antagonist (Td₂₂) in the compost was 9.21 log₁₀ spores/g wet weight of compost. Lettuce pathogen (*S. minor*) grown on millet seed was evenly inoculated approximately 20 mm below the soil surface at the rate of 2.0 g inoculum per pot. Soil without compost amendment, amended with pathogen only, or without pathogen, served as controls. Five replicate pots per treatment, each containing five seeds, were maintained for four weeks in a shade house. Pots were irrigated as required and the germinated seeds counted one week after sowing. Disease incidence on the lettuce seedlings was recorded from two weeks after sowing.

2.6 Data Analysis

The data obtained from this experiment was statistically analyzed by using analysis of variance (ANOVA) using Minitab software for Windows. The significance of differences between means was further tested using the least significant different (LSD) test at $P < 0.05$ following ANOVA.

3. Results

3.1 The Growth and Survival of Td₂₂ in WFW Compost

The growth and survival of Td₂₂ in compost over 28

days following different pre-treatments is presented in Fig. 1. Significant increases in CFU of Td₂₂ in steam-treated or untreated composts were observed in the first seven days, after which time the populations remained relatively constant at approximately 8.5 log₁₀ CFU/g dw in compost A or approximately 7.0 log₁₀ CFU/g dw in compost B.

The identity of Td₂₂ cultivated in non-sterile compost was confirmed by both morphological characteristics and isozyme profile (Fig. 2).

The initial bacterial populations (contaminants) in the mixes depended on the method of preparation. The population of the mesophilic bacteria in compost A was recorded at approximately one order of magnitude lower than that in compost B after two weeks incubation. However after four weeks of incubation, the growth rate of the indigenous mesophilic bacteria in compost A exceeded that of compost B, to result in a population of 9.3 log₁₀ CFU/g in compost A, in comparison with 8.5 log₁₀ CFU/g in compost B.

3.2 The Efficacy of Td₂₂-Grown Compost to Protect Lettuce from *S. minor* Attack

The Td₂₂-grown compost was found to be non toxic as between 88%-100% of the lettuce seeds were




Fig. 1 Population density of *Trichoderma* sp. (isolate Td₂₂) in three months old WFW compost, either briefly treated for 5 min at 121 °C prior to inoculation (Compost A), or directly inoculated following re-wetting (Compost B). Both compost types were brought to approximately field capacity with 2% (w/v) Phostrogen® solution (~350 mL/L). Each value in the graph is an average of three replicate determinations ± standard error.




Fig. 2 Contact print following polyacrylamide-gel electrophoresis to confirm survival of Td₂₂ in three-month old WFW compost. Dark bands indicate polygalacturonase and light bands indicate pectinesterase. Wells 1, and 9 – 14 contained the control reference, Td₂₂, while wells 2 – 8 contained re-isolated fungi from compost B on day 7 of sample collection.




Fig. 3 Protection of lettuce seedlings from attack by *S. minor*. Each value in the graph ± standard error bar is an average of 5 replicates. Bars with the same letter are not statistically significant at $P < 0.05$.

1. Nil pathogen and antagonist addition;
2. Pots amended with *S. minor* only;
3. Pots amended with 2.5% compost-grown Td₂₂ and *S. minor*;
4. Pots amended with 5% compost-grown Td₂₂ and *S. minor*;
5. Pots amended with 10% compost-grown Td₂₂ and *S. minor*;
6. Pots amended with 20% compost-grown Td₂₂ and *S. minor*.

observed to germinate in soil amended with 5-20% (v/v) of this compost. No inhibition of seed germination was evident even higher rate of Td₂₂-grown compost was applied 4 weeks after sowing.

The effectiveness of the Td₂₂-grown compost

amendment to protect lettuce seedlings from attack by *S. minor* is presented in Fig. 3. As indicated in Fig. 3, the rate of amendment of the compost/fungus was proportional to disease control, ranging from 32% provided by 2.5% (v/v) amendment to 100% protection provided by 20% (v/v) amendment at four weeks after sowing. This was statistically significant when compared to the control treatment (pots inoculated with *S. minor* only) where 100% mortality was observed at four weeks after sowing (Fig. 3).

4. Discussion and Conclusion

Air drying compost in a glasshouse for three weeks or brief steam-treatment (rather than sterilizing) with a view to minimizing the indigenous microbiota, prior to inoculation with Td₂₂, gave encouraging results. The CFU of this fungus cultured under non-sterile conditions increased by two to three orders of magnitude, although they were always lower than those obtained under sterile conditions (Fig. 1). The CFU were found to be higher in the briefly autoclaved compost (compost A) than those in the directly inoculated compost (compost B) following re-wetting. Application of heat treatment to the WFW compost (by briefly autoclaving at 121 °C) may have changed its composition. However the most probable reason for the better growth of Td₂₂ in this medium was the near-elimination of competition by mesophilic bacterial survivors in compost A relative to compost B. Similar results were also demonstrated by Nakasaki et al. [19] who found a reduced growth rate of a strain of *Bacillus subtilis* following inoculation of this antagonist into non-sterile (as compared with sterile), but freshly-cut grass clipping compost. This reinforces the notion that competition between the indigenous microbiota and the inoculated antagonists is important in the directed establishment of such antagonists in a compost matrix.

The ability of *Trichoderma* sp. to produce cellulase and pectolytic enzymes illustrated in Fig. 2 has been reported previously by Metcalf [20], Oksanen et al. [21], Domingues et al. [22], and Lee et al. [23]. These

and other enzymes are important in the degradation of complex carbon sources contained in WFW and the ability to produce them is advantageous when in competition for available major energy sources.

Assay of isoenzyme profiles (e.g. pectolytic enzyme profiles as undertaken in the present study) of fungal isolates provided a convenient tool to differentiate Td₂₂ from other isolates of *Trichoderma* sp. following re-isolation from non-sterile samples. This assay has been applied by many workers [24-26] to distinguish their fungal isolates from other related fungi.

The effectiveness of *Trichoderma* sp. (Td₂₂) grown in wood fibre waste compost to protect against *S. minor* in glasshouse trials has been demonstrated in the present study. Application of compost-grown Td₂₂ at rates up to 20% (v/v) was non-toxic (relative to controls) to lettuce seed in all treatments (Fig. 3).

The utilization of composted WFW as the authors described could be achieved in large scale at moderate cost while avoiding the problems of liquid cultures as outlined by Hadar et al. [27]. Application of compost-grown Td₂₂ after allowing growth for two weeks or more appeared to be advantageous, because it was mostly in the form of spores (unpublished data). This reduces concerns relating to viability in the field, because fungal spores will be relatively more resistant than mycelia to environmental stress [28].

Acknowledgment

The authors would like to acknowledge Dr. Dean Metcalf, Norske Skog Paper Mill Limited, and AusAid for the provision of *Trichoderma* sp. isolate Td₂₂, wood fiber waste (WFW), and financial support, respectively.

References

- [1] R.H. Cruickshank, J.I. Pitt, Identification of species in *Penicillium* subgenus *Penicillium* by enzyme electrophoresis, *Mycologia* 79 (1987) 614-620.
- [2] A.R. Jutsum, S.P. Heaney, B.M. Perrin, P.J. Wege, Pesticide resistance: Assessment of risk and the development and implementation of effective management strategies, *Pesticide Science* 54 (4) (1998)

- 435-446.
- [3] P. Margot, The challenge in chemical control of plant diseases, in: 8th International Congress of Plant Pathology, Christchurch, 1, New Zealand, 2003, p. 71.
- [4] R.E. Beever, Fungicide resistance: *Botrytis cinerea* versus dicarboximides as a case study, in: 8th International Congress of Plant Pathology, 1, Christchurch, New Zealand, 2003, p. 83.
- [5] M. Machala, J. Vondracek, Estrogenic activity of xenobiotics, *VeterinariMedicina* 43 (10) (1998) 311-317.
- [6] J. Miyamoto, W. Klein, Environmental exposure, species differences and risk assessment, *Pure and Applied Chemistry* 70 (9) (1998) 1829-1845.
- [7] V.S. Poovala, V.K. Kanji, H. Tachikawa, A.K. Salahudeen, Role of oxidant stress and antioxidant protection in acephate-induced renal tubular cytotoxicity, *Toxicological Sciences* 46 (2) (1998) 403-409.
- [8] C. Perrin-Ganier, Soil and environment: The fate of pesticides and organic pollutants, *Ocl-Oleagineux Corps Gras Lipide* 5 (5) (1998) 341-343.
- [9] A.J. Mmochi, R.S. Mberek, Trends in the types, amounts, and toxicity of pesticides used in Tanzania-Effort to control pesticide pollution in Zanzibar, *Ambio* 27 (8) (1998) 669-676.
- [10] T.J.J. DeCeuster, H.A.J. Hoitink, Prospect for composts and biocontrol agents as substitutes for methyl bromide in biological control of plant diseases, *Compost Science and Utilization* 7 (3) (1999) 6-15.
- [11] J.P. Painuly, S.M. Dev, Environmental dimensions of fertilizer and pesticide use, relevance to Indians agriculture, *International Journal and Pollution* 10 (2) (1998) 273-288.
- [12] P. Vidhyasekaran, R. Rabindran, M. Muthamilan, K. Nayar, K. Rajappan, N. Subramanian, et al., Development of a powder formulation of *Pseudomonas fluorescens* for control of rice blast, *Plant Pathology* 46 (1997) 291-297.
- [13] P. Vidhyasekaran, K. Sethuraman, K. Rajappan, K. Vasumathi, Powder formulations of *Pseudomonas fluorescens* to control pigeonpea wilt, *Biological Control* 8 (1997) 166-171.
- [14] L. Gazoni, J. Cozzi, K. Kobayashi, Survival of potential biocontrol bacteria in various formulations and their ability to reduce radish damping-off caused by *Rhizoctoniasolani*, *Journal of Plant Diseases and Protection* 105 (1) (1998) 41-48.
- [15] Y. Ramona, M.A. Line, Potential for the large scale production of a bio-control fungus in raw and composted paper mill waste, *Compost Science and Utilization* 10 (1) (2002) 57-62.
- [16] M.J. Jackson, M.A. Line, Composting pulp and paper mill sludge-effect of temperature and nutrient addition method, *Compost Science and Utilization* 5 (1) (1997) 74-81.
- [17] M.J. Jackson, Compost-recycling of pulp and paper mill sludge, Ph.D. Thesis, School of Agricultural Science, The University of Tasmania Australia, 1998.
- [18] Y. Ramona, Assessment of some antagonists to fungal plant pathogens, Ph.D. Thesis, Tasmania University, Australia, 2003.
- [19] K. Nakasaki, S. Hiraoka, H. Nagata, A new operation for producing disease-suppressive compost from grass clippings, *Applied and Environmental Microbiology* 64 (10) (1998) 4015-4020.
- [20] D.A. Metcalf, Biological control of onion white root rot, Ph.D. Thesis, School of Agricultural Science, The University of Tasmania-Australia, 1997.
- [21] T. Oksanen, J. Pere, L. Paavilainen, L. Viikari, Treatment and recycled kraft pulps with *Trichoderma reesei* hemicellulases and cellulases, *Journal of Biotechnology* 78 (1) (2000) 1785-1790.
- [22] F.C. Domingues, J.A. Querioz, J.M.S. Cabral, L.P. Fonseca, The influence of culture condition on mycelial structure and cellulase production by *Trichoderma reesei* Rut C-30, *Enzyme and Microbial Technology* 26 (5-6) (2000) 394-401.
- [23] I. Lee, B.R. Evans, J. Woodward, The mechanism of cellulase action on cotton fibres: Evidence from atomic force microscopy, *Ultramicroscopy* 82 (1-4) (2000) 213-221.
- [24] S.M. Neate, R.H. Cruickshank, A.D. Rovira, Pectic enzyme patterns of *Rhizoctoniasolani* isolates from agricultural soils in south Australia, *Trans. Br. Mycol. Soc.* 90 (1) (1998) 37-42.
- [25] G. Leone, *In vivo* and *in vitro* phosphate-dependent polygalacturonase production by different isolates of *Botrytis cinerea*, *Mycol. Res.* 94 (8) (1990) 1039-1045.
- [26] D.A. Metcalf, C.R. Wilson, Histology of *Sclerotiumcepivorum* infection of onion roots and spatial relationships of pectinases in the infection process, *Plant Pathology* 48 (1999) 445-452.
- [27] Y. Hadar, I. Chet, Y. Henis, Biological control of *Rhizoctoniasolani* damping-off with wheat bran culture of *Trichoderma harzianum*, *Phytopathology* 69 (1979) 64-68.
- [28] A. Tronsmo, *Trichoderma harzianum* in biological control of fungal diseases, in: R. Hall (Ed.), Principles and Practice of Managing Soil-Borne Plant Pathogens, APS Press, St. Paul, Minnesota, 1996, pp. 213-236.