Annals of Medical Research

E-ISSN 2636-7688



Volume 28 Number 8 Year 2021 Pages 1430-1610

www.annalsmedres.org

E-ISSN: 2636-7688

Journal Abbreviation: Ann Med Res

Published Monthly

Web page: https://www.annalsmedres.org

E-mail: annalsmedres@inonu.edu.tr

Formerly Journal of Turgut Ozal Medical Center - ISSN 1300-1744

Owner

On Behalf of Inonu University Faculty of Medicine

Mahmut Durmus (Dean) Inonu University, Faculty of Medicine, Department of Anesthesiology and Reanimation, Malatya, Turkey

Editor-in-Chief

Nevzat Erdil Inonu University Faculty of Medicine, Cardiovascular Surgery, Malatya, Turkey

Co Editors

Murat Kara Inonu University, Faculty of Medicine, Department of Medical Parasitology, Malatya, Turkey

Editorial Board

Adel Hamed Elbaih Suez Canal University Faculty of Medicine, Department of Emergency Medicine, Ismailia, Egypt

Ahmet Sigirci Inonu University, Faculty of Medicine, Department of Radiology, Malatya,Turkey

Ayse Seval Ozgu Erdinc Ministry of Health, Ankara City Hospital, Gynecology and Obstetrics, Ankara, Turkey

Cemşit Karakurt Inonu University, Faculty of Medicine, Department of Pediatric Cardiology Malatya, Turkey

Caner Ediz University of Health Sciences, Sultan Abdulhamid Han Education and Training Hospital, Urology, Istanbul, Turkey

Ender Gedik Baskent University, Faculty of Medicine, Anaesthesiology and Reanimation, Ankara, Turkey

Erdem Topal Inonu University Faculty of Medicine, Pediatric, Malatya, Turkey

Ertan Yetkin Derin Dere Hospital Cardiology, Istanbul, Turkey

Gokce Simsek Kirikkale University, Faculty of Medicine, Otorhinolaryngology, Kirikkale, Turkey

Hakan Parlakpinar Inonu University Faculty of Medicine, Medical Pharmacology, Malatya, Turkey

Aysegul Taylan Ozkan Hitit University, Faculty of Medicine, Medical Microbiology, Çorum, Turkey

Mehmet Hamamci Bozok University, Faculty of Medicine, Neurology, Yozgat, Turkey

Mehmet Kilic Firat University Faculty of Medicine, Pediatric Immunology and Allergy, Elazig, Turkey

Sertac Usta Inonu University, Faculty of Medicine, Department of General Surgery, Malatya, Turkey

Resit Sevimli Inonu University, Faculty of Medicine, Department of Orthopedic Surgery, Malatya, Turkey

Meltem Kurus Katip Celebi, University, Faculty of Medicine, Histology and Embology , Izmir, Turkey

Mustafa Canpolat Inonu University, Faculty of Medicine, Department of Anatomi, Malatya, Turkey

Neslihan Yucel Inonu University, Faculty of Medicine, Emergency Medicine, Malatya, Turkey

Numan Karaarslan Namik Kemal University Faculty of Medicine, Department of Neurosurgery, Tekirdag, Turkey

Ozkan Ozger Rumeli University, Medicalpark Canakkale Hospital, Neurosurgery, Istanbul, Turkey

Siho Hidayet

Inonu University, Faculty of Medicine, Department of Cardiology, Malatya, Turkey

Rizaldi Taslim Pinzon Universitas Kristen Duta Wacana | UKDW, Department of Neurology, Yogyakarta, Indonesia

Tuba Bayindir Inonu University Faculty of Medicine, Otorhinolaryngology, Malatya,Turkey

Yucel Duman Inonu University, Faculty of Medicine, Department of Clinic Microbiology, Malatya, Turkey

Language Editors

Murat Kara Inonu University, Faculty of Medicine, Department of Parasitology, Malatya, Turkey

Tayfun Guldur Inonu University, Faculty of Medicine, Department of Biochemistry, Malatya, Turkey

Biostatistics Editors

Saim Yologlu Inonu University, Faculty of Medicine, Department of Biostatistics and Medical Informatics, Malatya, Turkey

Cemil Colak Inonu University, Faculty of Medicine, Department of Biostatistics and Medical Informatics, Malatya, Turkey

Harika Gozde Gozukara Bag Inonu University Faculty of Medicine, Biostatistics and Medical Informatics, Malatya,Turkey

Publications Coordinator

Ozden Keser Inonu University Faculty of Medicine, Annals of Medical Research/Publications Coordinator, Malatya,Turkey

Derya Yılmaz Inonu University Faculty of Medicine, Annals of Medical Research/Publications Coordinator, Malatya,Turkey

This journal is indexed in TÜBİTAK/ULAKBİM Tıp Veri Tabanı, Türkiye Atıf Dizini, Türk Medline, EBSCO, Google Scholar, Akademik Dizin, Biblio Med. EuroPub, Sobiad, CABI, Indexs Copernicus, Advanced Sciences Index

Material published in the Journal is covered by copyright ©2021 Annals of Medical Research. All rights reserved.

Ann Med Res

August 2021 Volume 28/ Number 8 CONTENTS

Original Article

1430-4	The microbiology of acute cholangitis and associated in-hospital mortality Elmas Biberci Keskin, Gulay Okay
1435-9	Retrospective analysis of autologous stem cell transplantation outcomes in multiple myeloma patients with renal insufficiency Deniz Goren Sahin, Olga Meltem Akay, Neslihan Andic, Hava Uskudar Teke, Eren Gunduz
1440-4	YKL-40 and fibronectin levels in patients with placental invasion anomaly Oya Soylu Karapinar, Hanifi Sahin, Ilay Gozukara, Eda Adeviye Sahin, Oguzhan Ozcan, Burak Sezgin, Arif Gungoren
1445-9	Evaluation of clinicopathological characteristics and prognosis of lung ýnvasive mucinous adenocarcinomas based on computed tomography (CT) findings Running title: Invasive mucinous carcinomas according to CT findings Nurcan Unver, Neslihan Akanil Fener, Halide Nur Urer
1450-6	Evaluation of post-transplant complications, patient and graft survival in patients with autosomal dominant polycystic kidney disease after renal transplantation; A single center experience Cigdem Bugday, Muhammet Serdar Bugday, Ersoy Oksuz, Aydin Turkmen
1457-61	Relationship of neuropathic pain component and osteoarthritis stage in patients with knee osteoarthritis Raikan Buyukavci, Ibrahim Catma
1462-6	Are noninvasive biochemical parameters an alternative to liver biopsy in patients with chronic hepatitis B? Sevki Konur, Neslihan Surmeli, Adnan Ozkahraman, Ramazan Dertli, Yusuf Kayar
1467-70	Our sentinel lymph node experience in patients diagnosed with DCIS and microinvasive breast carcinoma Zeynep Bayramoglu, Ethem Omeroglu, Hande Koksal, Mehmet Ali Eryilmaz, Yasar Unlu

AIIII Meu nes	Ann	Med	Res
---------------	-----	-----	-----

August 2021 Volume 28/ Number 8 CONTENTS

August 2(021 Volume 28/ Number 8 CONTENTS	1520-4	resistance in COPD patients with exacerbation Burcu Yormaz, Duygu Findik, Hatice Turkdagi, Salih Macin, Kazim Korez, Mecit Suerdem
1471-6	Serum adiponectin levels in children with nonspecific infections Banu Aydin, Birgul Kirel, Basar Sirmagul	1525-31	Investigation of piwil2 expression as a biomarker in solid tumors of childhood Yilmaz Secilmis, Musa Karakukcu, Ozlem Canoz
1477-82	Does GSM-like 1800 MHz radiofrequency cause KRAS and p53 mutations in colon? : Histopatologically and microbiologically changes in colon Ezgi Ece Sag, Saadet Celikozlu, Hayri Dayioglu, Sibel Kokturk, Cengiz Ozzaim, Sinan Darcan	1532-9	Decreased expression of villin has adverse effect on the prognosis of patients with colorectal cancer Muhammet Calik
1483-9	Assessing of the forensic reports documented by forensic medicine polyclinic of Adiyaman education and research hospital in 2017-2018 Kerem Sehlikoglu, Huseyin Kafadar	1540-4	The association of lymph node involvement with lenfovascular invasion, perineural invasion and peritumoral lymphocytic reaction in surgically treated colon cancer patients Tunc Eren, Mustafa Demir, Tamer Akay, Ibrahim Ali Ozemir, Suleyman Orman, Haydar Yalman, Pefet Vigithagi, Orban Alimoglu
1490-5	Effect of pentoxifylline administration on an experimental rat model of intraperitoneal adhesion Omer Lutfi Akgul, Erkan Dalbasi, Mesut Gul, Ulas Alabalik	1545-50	A comparison of cyanoacrylate closure to radiofrequency ablation techniques for treatment of truncal venous insufficiency Fatih Uzunkaya, Aysegul Idil Soylu
1496-9	The healing effects of prp and thymoquinone on dss induced inflammatory bowel disease in rats Huseyin Ozden, Gokhan Karaca	1551-7	Histopathological investigation of the effects of a combination of parenteral nutrition and hunger on rabbit splenic tissue Semra Gurunluoglu, Elif Kayhan Kustepe, Harika Gozukara Bag
1500-7	Evaluation of the relationship between sociodemographic characteristics, quality of life, depression, drug compliance and biochemical parameters in patients with thalassemia major in Isparta Aslihan Badur, Demircan Ozbalci, Emine Guchan Alanoglu	1558-64	A comparison of the platelet index, neutrophil- lymphocyte and platelet-lymphocyte ratios of patients with differentiated thyroid cancer and nodular hyperplasia Mehmet Aslan, Mehmet Turan Cicek
1508-14	2D/4D proportion and preoperative anxiety: Is there an association? Ebru Canakci, Ufuk Arslan, Onur Bukagikiran, Ilker Coskun, Ali Altinbas, Tuba Catak, Ali Ozgul Saltali, Zubeyir Cebeci, Huseyin Erdinc Basar	1565-71	TROP2 and SOX-10 expression in the differential diagnosis of bladder tumors Ismail Harmankaya, Nevzat Serdar Ugras, Sumeyye Nur Tataroglu
1515-9	Menopausal transition may affect the onset of Herpes Zoster: An update on herpes zoster vaccine Servet Ozturk, Ebru Unal Akoglu, Eylem Emel Arikan, Canan Agalar	1572-6	Death anxiety of doctors and nurses with levels of depression related to death and the factors affecting them Gulsum Karaoglu, Mehmet Ali Kurcer, Zeynep Erdogan

Short term results comparison of antibiotic

August 2021 Volume 28/ Number 8 CONTENTS

1577-82	The association of non-motor symptoms with motor symptoms in parkinson's disease Seyda Cankaya

1583-8	Negative correlation between TNF-α and triglyceride levels in patients with cerebral infarction Hikmet Sacmaci, Murat Cakir, Seda Sabah Ozcan
	HIKINEL Sacinaci, Mulat Cakii, Seua Saban Ozcan

1589-95	Effect of enteral nutritional support on
	nutritional status of patients with gastric cancer
	receiving chemoradiotherapy after gastrectomy
	Ozlem Elmas, Ramazan Kozan

Case Reports

	A 19-day neonatal case with accidental alcohol
1596-8	intoxication during umbilical care
	Oksan Derinoz Guleryuz

Review Article

	Prolonged use of statins and peripheral neurop			
	athy: A systematic review			
1599-604	I Putu Eka Widyadharma, Eric Hartono Tedyanto,			
	Dewa Putu Gede Purwa Samatra, Anak Agung			
	Ayu Putri Laksmidewi, I Made Oka Adnyana			

1605-10	Glucose metabolism and oncogenes in cancer
	Cuma Mertoglu

Prolonged use of statins and peripheral neuropathy: A systematic review

I Putu Eka Widyadharma, Eric Hartono Tedyanto, Dewa Putu Gede Purwa Samatra, Anak Agung Ayu Putri Laksmidewi, I Made Oka Adnyana

Department of Neurology, Sanglah General Hospital and Udayana University, Bali, Indonesia

Copyright@Author(s) - Available online at www.annalsmedres.org Content of this journal is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.



Abstract

Peripheral neuropathy is one of the most common neurological disorder. Traditionally, diabetes mellitus is the most common aetiology of peripheral neuropathy. However, there are many other aetiologies of peripheral neuropathy, such as drug-induced, spinal trauma, alcoholism, vitamin B12 deficiency, and many more. In previous studies, long-term statin use was also associated with peripheral neuropathy.

We conducted a systematic search through PubMed and Embase using the protocol following the PRISMA statement in making a systematic review. W e included article that published in the last 10 years (from 2011 until July 2020). Two reviewers extracted data from the articles that have been selected. We extracted the study population, study design, type of intervention, control variable, p-value, and the outcome of the study

We found 250 journals in our search strategy, and at the end, 4 journals included. Some studies revealed that peripheral neuropathy was observed during the long-term treatment of statin. However, some studies also revealed that statin does not increase the risk of peripheral neuropathy.

The possible mechanism of statin-induced peripheral neuropathy in humans are disturbing the function and integrity of cell membrane, the role of alfa-tocopherol or Vitamin E, and the role of ubiquinone (Coenzyme Q10). In contrast, it is believed that statins lowering the risk of peripheral neuropathy through the anti-inflammatory effects of statins. Different result from each study can be caused by various factors. Besides the peripheral neuropathy side effect that described in the literature, statins are well tolerated. Awareness the side effects would lead to better treatment. Future study about association between statins and peripheral neuropathy is needed.

Keywords: Peripheral neuropathy; prolonged use; statins

INTRODUCTION

Peripheral neuropathy is one of the most common neurological disorder. Symptoms of peripheral neuropathy such as tingling sensation, pain, numbness, and also weakness. The symptoms usually first affect the leas. then hands, and the other part of body. Sometimes, it is called stocking-gloves patterns because the boundaries of the symptoms are clear like we wear the stockinggloves (1). Peripheral neuropathy happens at any age, but most common found over 55 years of age (2). Traditionally, diabetes mellitus is the most common aetiology of peripheral neuropathy. However, there are many other aetiologies of peripheral neuropathy, such as drug-induced, spinal trauma, alcoholism, vitamin B12 deficiency, and many more. About 26% of people over 65 years of age, develop peripheral neuropathy without any aetiology and predisposing factors (3).

Drug-induced neuropathy is one of the main causes of peripheral neuropathy. The prevalence of drug-induced neuropathy is about 2-4% (4,5). Chemotherapy drugs, metronidazole, allopurinol, phenytoin, and colchicine may develop peripheral neuropathy. In previous studies, long-term statin use was also associated with peripheral neuropathy. Several statin patients report that they experienced neuropathic pain, thus creating the question of whether statins cause peripheral neuropathy or not (6). Statins are the drugs most common used to lower blood cholesterol. Statin is a Hydroxymethyl-glutaryl-CoA (HMG-CoA) reductase inhibitor, that lowering plasma cholesterol, especially Low-density Lipoprotein (LDL) (7). Due to role of statin that lowering cholesterol production, it may disturb the function and integrity of cell membrane. (8) Statins also inhibit ubiquinone (Coenzyme Q10), an important enzyme for mitochondrial respiration. Inhibiting

Received: 28.07.2020 Accepted: 04.11.2020 Available online: 20.08.2021

Corresponding Author. I Putu Eka Widhyadharma, Department of Neurology, Sanglah General Hospital and Udayana University, Bali, Indonesia **E-mail:** eka.widhyadharma@unud.ac.id

ubiquinone may reduce the produce of energy that is required for nerve cells. Therefore, statins may have side effects such as central nervous system disorders and peripheral neuropathy (9,10).

Since statin is the most widely used lipid-lowering drugs, awareness the side effects would lead to better treatment. The purpose of this review is to perform a systematic review to identify association between prolonged use of statins and peripheral neuropathy.

Searching and Screening

We conducted a systematic search through PubMed and Embase using the protocol following the PRISMA statement in making a systematic review. We included article that published in the last 10 years (from 2011 until July 2020). We used specific keywords 1. "STATIN"; AND 2. "PERIPHERAL NEUROPATHY" on search database.

Inclusion and Exclusion Criteria

We included original and published articles with a cohort, cross-sectional, experimental, randomized-controlled

trial, and case-control study designs. We exclude articles that are descriptive studies, case reports, or qualitative studies. Table 1 presents the eligibility criteria in the PICOS model. First, the title and abstract were screened by two-reviewer, then full-text screening was performed based on the eligibility criteria.

Data Extraction

Two reviewers extracted data from the articles that have been selected. We extracted the study population, study design, type of intervention, control variable, p-value, and the outcome of the study. The disagreement of data that have been extracted by two reviewers will be resolved by the third reviewer

Quality Assessment

Fourteen criteria will be identified in each of the included journals. See Table 2 for more info about methodological quality assessment.

Ethical Approval

Ethical approval for this review was not required.

Table 1. Eligibility criteria in PICOS model						
	Inclusion	Exclusion				
Population	Human, adults (>18 years) with history of statin use	Non-human				
Intervention	Do not observed patients with lo long-term statins use use (only with a short-term stat agent of lipid-lowering					
Control	Healthy patients who do not take statins					
Outcomes	peripheral neuropathy as measured by certain method and assessment	besides peripheral neuropathy				
Study Design	Cohort	Case Report				
	Case-control	Case Series				
	Cross-sectional	Descriptive Studies				
	Randomized-Controlled Trial	Qualitative studies				
Language	English	Non-English				
Statistics	An adequate method of statistics that explain the association between 2 variables (p value)	Just only descriptive studies				

Study Selection

We found 250 journals in our search strategy. At the first screening, we excluded 50 duplicated journals. Second, the title and abstract were reviewed based on the eligibility criteria, as many as 158 were excluded, 42 remained. At the third screening, full-text was reviewed, as many as 38 journals were excluded. Finally, 4 studies met the eligibility criteria and were included in this review. The flow diagram of systematic searching strategy is showed on Figure 1.

Quality Assessment

Methodological quality of the studies was high. All of the studies described the eligibility criteria, study population, and population characteristic. For the quality of statistical analysis, all of the studies described statistical model appropriately. All of other item score were 4, except the participation rate, frequencies of most important determinants, and controlled for confounding or effect modification, the score were 3. The quality assessment is presented in the Table 2.

Study Report Characteristics

The studies were conducted in Malaysia, Iran, Turkey, and Czech Republic. There were 2 experimental studies, 1 cohort study, and 1 case-control study. The number of study subjects ranged from 39 to 757. Otruba et al revealed that of the 42 patients who used statins for a long time for more than 2 years, they did not have symptoms of

peripheral neuropathy. But, in the ENMG examination, there was a prolongation of F-waves. So, Otruba et al. concluded that prolonged use of statins can cause silent neuropathy, because there are no symptoms, but found abnormalities on ENMG examination. Ozdemir et al revealed that of the 50 study subjects who used statins for a long time period, 33 out of 50 experienced peripheral neuropathy symptoms. Study results from Mohammad Reza are different from the other 3. Mohammadreza revealed that there was no significant difference between the intervention and control groups (p=0.055), but there was a difference between the amplitude of the sural and peroneal nerves which showed that there was peripheral neuropathy. This might be due to the small sample size, which is only 39. In contrast, a study from Warendorf revealed that statin do not increase the risk of polyneuropathy. Whereas, statin use decreased the odds ratio of polyneuropathy as duration of statin exposure increased (p=0,03, OR 0,5 95%; CI 0,34-0,9). More info about study characteristic is presented in Table 3.



Figure 1. Flow diagram. The search strategy for systematic review

Table 2. Methodological quality assessment					
	Otruba, 2011	Ozdemir, 2019	Warendorf, 2019	Mohammadreza, 2018	Total per item
Study design and object					
1.objective of the study was clear	1	1	1	1	4
Study population					
2.Description of eligibility criteria.	1	1	1	1	4
3. Description of selection of study population	1	1	1	1	4
4. Description of population characteristics	1	1	1	1	4
5. Case-control: definition of cases and control were mentioned	-	-	1	1	2
6. Participation rate is more than 80%	0	0	0	0	0
Measure of instruments					
7.Appropriate definition and assessment of outcome	1	1	1	1	4
8.Appropriate definition and assessment of determinants	1	1	1	1	4
Data					
9. The measurements of most important outcome	1	1	1	1	4
10.Frequencies of most important determinants	1	1	0	1	3
11.Measures of association presented	1	1	1	1	4
Analysis					
12.Appropriate statistical model	1	1	1	1	4
13.Controlled for confounding or effect modification	0	1	1	1	3
Other					
14.No identification of funding sources and conflicts of interest	1	1	1	1	4
15.0ther sources of bias (that not mention)	0	0	0	0	0
Total score for each study and/or well performed	11	12	12	13	

1, and/or well performed; 0, has not been reported, has not been well performed or unclear; -, not have been applicable yet

Table 3. Study Chara	cteristics					
Author Otruba (2011) (11)	Design RCT	Subject 42 patients	Intervention Long-term statin (simvastatin 20mg) use (>2 years)	Control -	p <0.001	Result In this study, there were no peripheral neuropathy symptoms found on neurological examination. However, the Electroneuromyography (ENMG) examination found prolongation of F-wave mean latency on peroneal and tibial nerves. The study confirmed that long- term treatment with statins caused a clinically silent but still definite damage to peripheral nerves when the treatment lasts longer than 2 years
Mohammadreza, 2018(4)	Case-control	78 aged 35-55 years	39 patients who received statin for 6 months	39 healthy controls	0.055	No significant difference between two groups (statin and control group), but there was a significant difference between the amplitude of sural nerve and peroneal nerve, indicates peripheral neuropathy
Ozdemir (2019) (9)	RCT	100 patients	50 patients with long term statin use (>1 year)	50 healthy people who do not take a statin that match for age and gender	<0.01	In the group using long-term statins, the ENMG results revealed polyneuropathy in 33 out of 50 patients (34 on atorvastatin, 16 on rosuvastatin). No polyneuropathy was revealed in control group. There was a significant difference between intervention and control group (p<0.01), but not between atorvastatin and rosuvastatin group (p=0.720). This study showed that long-term use of statin (>1 year) could induced peripheral neuropathy based on ENMG examination
Warendorf, 2019 (12)	Case-control	616 patients	333 with cryptogenic axonal polyneuropathy	283 healthy controls	P=0.03; OR 0.5 95% CI 0.34- 0.95	Statin use decreased the odds ratio of polyneuropathy as duration of statin exposure increased

In this systematic review, we tried to identify the association between prolonged use of statins and peripheral neuropathy. The most common neurological manifestation of prolonged use of statin is muscular disorder, from mild to severe like rhabdomyolysis and myoglobinuria. The second neurological manifestation is peripheral neuropathy. Some studies revealed that peripheral neuropathy was observed during the long-term treatment of statin (13). However, some studies also revealed that statin does not increase the risk of peripheral neuropathy (12). In human studies, statins may induce peripheral neuropathy. But in animal studies, statins reduced the risk of peripheral neuropathy.

The possible mechanism of statin-induced peripheral neuropathy in humans are disturbing the function and

integrity of cell membrane, the role of alfa-tocopherol or Vitamin E, and the role of ubiquinone (Coenzyme Q10). Vitamin E or alfa-tocopherol is a fat-soluble vitamin and has a function in maintaining nerve health. It is transported to tissue by LDL.(4) Thus, lowering LDL means lowering the transport of alfa-tocopherol, and may play a role in peripheral neuropathy (14). The association between ubiquinone, statin, and peripheral neuropathy is not well explained. Ubiquinone is an enzyme for mitochondrial respiration, and statins inhibit this enzyme. Inhibiting ubiquinone may reduce the production of energy that is required for nerve cells. Therefore, statins may have side effects such as central nervous system disorders and peripheral neuropathy.

While peripheral neuropathy is associated indirectly with poor glycaemic control in diabetes, peripheral neuropathy is also found in people who do not have diabetes, but hyperlipidemia (15). Association between hyperlipidemia and peripheral neuropathy has been discussed. People with hyperlipidemia have high amounts of LDLs and oxysterols, that promotes the proliferation of pro-inflammatory cytokines. Thus, it may cause local inflammatory lesions on the peripheral neuron (16). In animal studies, it is believed that statins lowering the risk of peripheral neuropathy through the anti-inflammatory effects of statins. In in-vitro study, statins decrease the proliferation of monocyte and pro-inflammatory cytokines such as interleukin-2, interleukin-4, interleukin-6, and interferon-w(17).

Different result from each study can be caused by various factors. One of which is the methods and tools for diagnosing peripheral neuropathy. Some study uses ENMG for diagnosing PN, while some study only uses a questionnaire to diagnose PN. It is known that the gold standard to diagnose PN is using ENMG, because the symptoms may be silent. It is proven by the study of Otruba. There were no peripheral neuropathy symptoms found on neurological examination but found the prolongation of F-wave in ENMG examination. (11) Comorbid factors like cardiovascular disease and metabolic syndrome could lead to over-diagnose statin-induced peripheral neuropathy since these conditions are also associated with cryptogenic axonal polyneuropathy (18).

There are some limitations to this review. First, the small number of studies included in this review (only 4 studies). Second, there were different diagnostic criteria, modality, and another comorbid factor in each included study. Third, not all study explained the dosage and type of statin that used for intervention. So, there might be under or overdiagnose. Hopefully, this systematic review could give update information about the related topic.

CONCLUSIONS

Statin is the most widely used lipid-lowering drugs and often used for a long time period. Their efficacy and effectiveness for treating hypercholesterolemia were well established. Besides the peripheral neuropathy side effect that described in the literature, statins are well tolerated. The benefits and disadvantages must be considered. Awareness the side effects would lead to better treatment. Future study about association between statins and peripheral neuropathy is needed.

Competing interests: The authors declare that they have no competing interest.

Financial Disclosure: There are no financial supports.

REFERENCES

1. Mohamed A Hammad, Syed Azhar Syed Sulaiman, Saleh Alghamdi, et al. Statins-related Peripheral Neuropathy Among Diabetic Patients. Diabetes & Metabolic Syndrome: Clinical Research and Reviews. Elsevier,2020;14:341-6.

- St Charles JL, Bell JA, Gadsden BJ, et al. Guillain Barré Syndrome is induced in Non-Obese Diabetic (NOD) mice following Campylobacter jejuni infection and is exacerbated by antibiotics. J Autoimmun 2017;77:11-38.
- 3. West B. The implications of statin induced peripheral neuropathy. J Foot Ankle Res 2011;4:57.
- 4. Emad M, Arjmand H, Farpour HR, Kardeh B. Lipidlowering drugs (statins) and peripheral neuropathy. Electron Physician 2018;10:6527-33.
- 5. de Langen JJ, van Puijenbroek EP. HMG-CoAreductase inhibitors and neuropathy: Reports to the Netherlands Pharmacovigilance Centre. Neth J Med 2006;64:334-8.
- McCarberg B, D'Arcy Y, Parsons B, et al. Neuropathic pain: a narrative review of etiology, assessment, diagnosis, and treatment for primary care providers. Current Medical Research and Opinion. 2017;33:1361-9.
- Svendsen T de K, Nørregaard Hansen P, García Rodríguez LA, et al. Statins and polyneuropathy revisited: case-control study in Denmark, 1999–2013. Br J Clin Pharmacol 2017;83:2087-95.
- 8. Bosomworth NJ. Statin therapy as primary prevention in exercising adults: Best evidence for avoiding Myalgia. J Am Board Fam Med 2016;29:727-40.
- 9. Ozdemir IH, Copkiran O, Tikiz H, et al. Peripheral polyneuropathy in patients receiving long-term statin therapy. Turk Kardiyol Dern Ars 2019;47:554-63.
- Apostolopoulou M, Corsini A, Roden M. The role of mitochondria in statin-induced myopathy. Eur J Clin Invest 2015;45:745-54.
- 11. Otruba P, Kanovsky P, Hlustik P. Treatment with statins and peripheral neuropathy: Results of 36-months a prospective clinical and neurophysiological followup. Neuroendocrinol Lett 2011;32:688-90.
- 12. Warendorf JK, Vrancken AFJE, Van Eijk RPA, et al. Statins do not increase risk of polyneuropathy A case-control study and literature review. Neurology. 2019;92:2136-44.
- 13. Golomb BA, Evans MA. Statin adverse effects: A review of the literature and evidence for a mitochondrial mechanism. American Journal of Cardiovascular Drugs 2008;8:373-418.
- 14. Muller DPR. Vitamin E and neurological function. Molecular Nutrition and Food Research 2010;62:107-12.
- 15. Rajabally YA. Neuropathy and impaired glucose tolerance: An updated review of the evidence. Acta Neurologica Scandinavica 2011;124:1-8.
- 16. Vincent AM, Hinder LM, Pop-Busui R, et al. Hyperlipidemia: A new therapeutic target for diabetic neuropathy. Journal of the Peripheral Nervous System. 2009;14:257-67.
- 17. Gustavsson C, Agardh CD, Zetterqvist A V., et al. Vascular cellular adhesion molecule-1 (VCAM-1) expression in mice retinal vessels is affected by both hyperglycemia and hyperlipidemia. PLoS One. 2010; 5:12699.

18. Visser NA, Vrancken AFJE, Van Der Schouw YT, et al. Chronic idiopathic axonal polyneuropathy is associated with the metabolic syndrome. Diabetes Care 2013;36:817-22.