

Bali Medical Journal



INDONESIAN PHYSICIAN FORUM &
INDONESIA COLLEGE OF SURGEONS,
INDONESIA

Home > Editorial Board & Reviewer

Editor-in-Chief

Prof. Dr. Sri Maliawan, SpBS

(Scopus ID), (Google scholar)

srimaliawan@unud.ac.id / maliawans@yahoo.com

Department of Neuro Surgery, Udayana University

Sanglah General Hospital

Bali - Indonesia

Associate Editor

Prof. Putra Manuaba, M.Phil

(Scopus ID), (Google Scholar)

putramanuaba@unud.ac.id / putramanuaba28@yahoo.com

Biomedicine Postgraduate Program, Udayana University

Bali - Indonesia

Prof. Ketut Suwiyoga, SpOG

(Scopus ID)

suwiyoga@unud.ac.id

Faculty of Medicine, Udayana University, Sanglah Hospital Denpasar, Bali-Indonesia

Editorial Board for Regional America

Ankit Sakhuja, M.B.B.S., F.A.C.P., F.A.S.N.

(Scopus ID)

asakhuja@med.umich.edu

Nephrology and Hypertension Cleveland Clinic (United States)

Editorial Board for Regional Australia

Prof. John Svigos, MB. BS. DRCOG., FRCOG., RANZCOG

(Scopus ID)

john@svigos.com.au

Ashford Hospital & Faculty of Health Sciences, University of Adelaide, Australia

dr Deasy Ayuningtyas Tandio MPH-MBA.

(OrcidID)

deasytandio@yahoo.com

James Cook University Australia Master of Public Health Master Of Business Administration, Indonesia

Editorial Board for Regional Europa

Prof. Harald Hoekstra

(Scopus ID)

jsvigos@primus.com.au

Editorial Board for Regional Asia

Prof Huang Qin

(Scopus ID)

qhuang@cqu.edu.cn

Chairman Dept. of Neurosurgery, Guangdong 999 Hospital Guangzhou China

Assoc. Prof. Mohammad Amin Bahrami

(Scopus ID)

aminbahrami1359@gmail.com

Head of healthcare management department, Shahid Sadoughi University of Medical Sciences, Yazd,Iran

Dr. Tanveer Beg, PhD

(Scopus ID)

tbmirza@jazanu.edu.sa

Assistant Professor, Department of Biology, Faculty of Science, Jazan University, Jazan, Saudi Arabia.



WEB OF SCIENCE™



ELSEVIER
Scopus



DIRECTORY OF
OPEN ACCESS
JOURNALS



Full Indexing List

In Press


Submit An Article

Scopus Citedness



HP Spectre x360 13

Uncompromising power
to fuel your passions



Editorial Board Members

Prof. Andi Asdul Islam

(Scopus ID), (Google Scholar)

undee@med.unhas.ac.id

Faculty of Medicine Hasanudin University, Makasar-Indonesia

Prof. Dr. dr. Abdul Hafid Bajamal, Sp.BS

(Scopus ID)

hfbajamal@gmail.com

Faculty of Medicine Airlangga University, Surabaya-Indonesia

Dr. dr. I Wayan Sudarsa, Sp.B(K) Onk, FINACS, FICS.

(Scopus ID), (Google Scholar), (Researchgate)

dr. I.B. Amertha P. Manuaba, SKed, MBIomed.

(Scopus ID), (Google Scholar), (ORCID), (Researcher ID) (Researchgate)


AmerthaManuaba@gmail.com / Amertha_Manuaba@unud.ac.id

Faculty of Medicine, Universitas Udayana, Indonesia

Editorial inquiries to be addressed to:

email 1: editorbalimedicaljournal@gmail.com

email 2: editor@balimedicaljournal.org



Vol 10, No 3 (2021): (Available online: 1 December 2021)

Table of Contents

ORIGINAL ARTICLE

Convalescent plasma as an adjunctive treatment for severe and critically ill COVID-19

Damai Santosa, Muchlis Achsan Udji Sofro, Retnaningsih Retnaningsih, Eko Adhi Pangarsa, Budi Setiawan, Nur Farhanah, Fathur Nur Kholis, Thomas Handoyo, Sofyan Rahardjo, Like Rahayu Nindhita, Iva Puspitasari, Ridho M. Nalbaho, Retty Kharismasari, Ika Kartiyani, Vina Yunarvika, Daniel Rizky, Suhartono Suhartono

Online First: September 14, 2021
| DOI: 10.15562/bmj.v10i3.2590

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Root cause analysis of maternal deaths at Dr. Soetomo General Academic Hospital Surabaya, Indonesia in 2019

Dimas Ryan Desetyaputra, Budi Prasetyo, Hanifa Erlin Damayanti

Online First: October 07, 2021
| DOI: 10.15562/bmj.v10i3.2723

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Functional Movement Screen (FMS) as a determinant factor to assess the risk of injuries in young athletes at Jakarta Province, Indonesia

Anthony Anthony, Trevino Aristarkus Pakasi, Jull Kurniarobbi

Online First: October 23, 2021
| DOI: 10.15562/bmj.v10i3.2638

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

RUMI DOPING application development as Anti-doping information media for Indonesian national paralympic committee Athletes

Rumi Iqbal Doewes, Gunathevan Elumalal, Siti Hartini Azmi, Singgih Hendarto, Rina Nurhudi Ramdhani

Online First: November 05, 2021
| DOI: 10.15562/bmj.v10i3.2781

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Short cervix, wide anterior cervical angle and low cervical consistency index as a predictor of preterm birth: a case-control study

Andy Susanto, Anak Agung Gede Putra Wiradyana, Ketut Suwiyoga, I Gusti Putu Mayun Mayura, Ida Bagus Fajar Manuaba, I Nyoman Bayu Mahendra, I Wayan Megadhana

Online First: November 18, 2021
| DOI: 10.15562/bmj.v10i3.2767

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Type of anemia correlate with ferritin levels in patients with Chronic Kidney Disease (CKD): a cross sectional study

Linda Rosita, Nurrin Jannatin, Barkah Djaka Purwanto, Tri Ratnaningsih, Usi Sukorini, Zulkifli Supranto Simanjuntak, Rahma Yuantari

Online First: September 15, 2021
| DOI: 10.15562/bmj.v10i3.2418

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Expression of Growth Associated Protein-43 (GAP-43) and Brain-derived Neurotrophic Factor (BDNF) from Adipose Mesenchymal Stem Cells (ASC) with fresh frozen nerve scaffold in 5% hypoxic condition

I Putu Gede Pradnyadewa Pradana, Asra Al Fauzi, Heri Suroto

Online First: October 23, 2021
| DOI: 10.15562/bmj.v10i3.2575

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

A new innovation in topical diabetic foot ulcer; hyaluronic acid platelet-rich fibrin (HAPRF) gel - a study in inflammation and angiogenesis

Ronald W. Kartika, Idrus Alwi, Em Yunir, Sarwono Waspadji, Saptawati Bardosono, Suzzana Immanuel, Todung Silalahi, Saleha Sungkar, Jusuf Rachmat, Franciscus D. Suyatna, Mirta Hedyati Reksodiputro

Online First: November 04, 2021
| DOI: 10.15562/bmj.v10i3.2317

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Changes in Ki67 expression and clinical response to neoadjuvant chemotherapy in locally advanced breast cancer (LABC)

Desak Gede Agung Pramesti Devi, Heru Purwanto, Willy Sandhika

Online First: November 12, 2021
| DOI: 10.15562/bmj.v10i3.2700

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Implementation of child nutrition anthropometry pocketbook for early childhood education to increase the ability of teachers to assess nutritional status of early childhood education students in Gianyar Regency, Bali, Indonesia

Ni Nengah Ariati, Ni Komang Wiardani, Anak Agung Ngurah Kusumajaya, Arma Fetria

Online First: December 04, 2021
| DOI: 10.15562/bmj.v10i3.2662

[Abstract](#) | [PDF](#) |



WEB OF SCIENCE™



ELSEVIER
Scopus



DIRECTORY OF
OPEN ACCESS
JOURNALS



SINTA^{S1}
Science and Technology Index

Full Indexing List

In-Press

Submit An Article

Scopus Citedness



Install & Chrome

Google

Learn More

ORIGINAL ARTICLE

Intravenous Wharton's Jelly stem cell increased the number of β cells pancreas and reduced the fasting blood glucose level in diabetes mellitus Wistar rat male (*Rattus norvegicus*)

Nadia Permatasari, Wimpie Pangkahila, Anak Agung Gde Budhiarta, J. Alex Pangkahila, I Gusti Made Aman, Desak Made Wihandani, Indra Bachtiar

Online First: December 02, 2021
| DOI: 10.15562/bmj.v10i3.2306

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Comparative efficacy of topical Adipocyte-derived Mesenchymal Stem Cells-Conditioned Medium (ADMSC-CM) and Amniotic Membrane Mesenchymal Stem Cells-Conditioned Medium (AMSC-CM) on chronic plantar ulcers in leprosy: a randomized controlled trial

Medhi Denisa Alinda, Paulus Mario Christopher, Muhammad Yulianto Listiawan, Anang Endaryanto, Heri Suroto, Fedik Abdul Rantam, Esti Hendradi, Hari Basuki Notobroto, Sri Linuwih Susetyo Wardhani Menaldi, Cita Rosita Sigit Prakoeswa

Online First: December 10, 2021
| DOI: 10.15562/bmj.v10i3.2747

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Effects of sodium bicarbonate ingestion on prolonged running performance and recovery in trained runners

Lew Ka Shing, Ler Hui Yin, Gee Yong Yong, Wee Eng Hoe

Online First: December 14, 2021
| DOI: 10.15562/bmj.v10i3.2741

ORIGINAL ARTICLE

Correlation between Left Ventricular Ejection Fraction (LVEF) and six-month mortality risks after hospital discharge following myocardial infarction in patients with Non-ST-Elevation Myocardial Infarction (NSTEMI) at Prof. Dr. R. D. Kandou General Hospita

Christian Yosaputra, Reginald Leopold Lefrandt, Agnes Lucia Panda, Janry Pangemanan

Online First: December 22, 2021
| DOI: 10.15562/bmj.v10i3.2749

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

The Comparison of intravesical hyaluronic acid and other modalities for the improvement of interstitial cystitis symptoms: a systematic reviews and meta-analysis

St Cahyo Ariwicaksono, Harrina Erlianti Rahardjo

Online First: December 21, 2021
| DOI: 10.15562/bmj.v10i3.2510

ORIGINAL ARTICLE

The effectiveness of multiflora honey to prevent hepatotoxicity in invasive ductal breast cancer patients with FAC chemotherapy

Dwi Feris Martua Sidabutar, Selamet Budijitno, Awal Prasetyo

Online First: December 22, 2021
| DOI: 10.15562/bmj.v10i3.2660

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

The differences of lactate dehydrogenase and activin A levels among thalassemia major and non-thalassemia

Ryan Alexander Gunawan, Nyoman Sucl Widyastiti, Ariosta Ariosta, Rina Pratiwi, Dwi Retnoningrum, Dwi Ngestiningsih, Yetty Movieta Nancy

Online First: December 28, 2021
| DOI: 10.15562/bmj.v10i3.2241

ORIGINAL ARTICLE

Ethyl p-methoxycinnamate isolated from *Kaempferia galanga* L. rhizome reduces airway remodeling in asthmatic rat models

Ismiranti Andarini, Harsono Salimo, Bambang Purwanto, Setyo Sri Rahardjo, Brian Wasita, Vitri Widyaningsih

Online First: December 30, 2021
| DOI: 10.15562/bmj.v10i3.2701

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Correlation between Bone Mineral Density (BMD) with blood phosphorus levels in elderly women: study using Dual Energy X-ray Absorptiometry (DXA)

Adhikarmika Aripriandari, Hermina Sukmaningtyas, Dwi Ngestiningsih, Lusiana Batubara

Online First: December 28, 2021
| DOI: 10.15562/bmj.v10i3.2848

ORIGINAL ARTICLE

Relationship between plasma atherogenic index and xanthelasma palpebrarum

Sri Karunia Setio Wati, Imam Budi Putra, Nelva Karmila Jusuf

Online First: December 30, 2021
| DOI: 10.15562/bmj.v10i3.2813

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

Comparison of the sacral ratio between children with Hirschsprung disease and non-Hirschsprung

Harry Galuh Nugraha, Pramesti Indri Miranti

Online First: September 15, 2021
| DOI: 10.15562/bmj.v10i3.2352

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

The Association between Medical History-based Risks and Sepsis Events in Immunocompromised Patients according to Type III Stratification of the Indonesian Regulation on the Prospective Antimicrobial System (Regulasi Antimikroba Sistem Prospektif / RASPRO)

Ronald Irwanto Natadidjaja, Armi Setia Kusuma, Gede Bangun Sudradjad, Lies Nugrohowati

Online First: December 30, 2021
| DOI: 10.15562/bmj.v10i3.2561

[Abstract](#) | [PDF](#) |

ORIGINAL ARTICLE

The correlation between sebum levels and pityriasis versicolor

Arzia Mustika, Meidina Kusuma, Lukmanul Hakim Nasution

Online First: December 30, 2021
| DOI: 10.15562/bmj.v10i3.2859

[Abstract](#) | [PDF](#)

ORIGINAL ARTICLE

Relationship between serum IgE levels and psoriasis area and severity index (PASI) score in psoriasis vulgaris

Fitra Aina Hidayat, Kristo Alberto Nababan, Irma Damayanti Roesyanto

Online First: December 21, 2021
| DOI: 10.15562/bmj.v10i3.2861

[Abstract](#) | [PDF](#)

ORIGINAL ARTICLE

Comparison between the diagnosis of acne vulgaris by teledermatology and face-to-face consultations

Anggita Dwi Puteri Rangkuti, Nelva Karmila Jusuf, Imam Budi Putra

Online First: December 30, 2021
| DOI: 10.15562/bmj.v10i3.2792

[Abstract](#) | [PDF](#)

ORIGINAL ARTICLE

Correlation between short-term heart rate variability and QT-Interval with left ventricular systolic function in patients with stable Coronary Artery Disease (CAD): a cross-sectional study

Agnes Lucia Panda, Natalia Christina Intania Polli, Victor Giovannie Xaverison Rooroh, Andreas Erick Haurissa, Agustinus Mahardhika Sarayar, Filipus Michael Yofrido

Online First: December 30, 2021
| DOI: 10.15562/bmj.v10i3.2862

[Abstract](#) | [PDF](#)

ORIGINAL ARTICLE

The difference in the effectiveness of warm compress and active stretching exercise in reducing dysmenorrhea pain

Ni Wayan Tianing, Made Hendra Satria Nugraha, Agung Wiwiek Indrayani, I Putu Eka Widyardharma

Online First: December 30, 2021
| DOI: 10.15562/bmj.v10i3.2301

[Abstract](#) | [PDF](#)

REVIEW

Adipose-Derived Stem Cells (ADSCs): a review article

Ida Bagus Putra Pramana, Anak Agung Gde Oka, I Nyoman Mantik Astawa, Tjokorda Gde Bagus Mahadewa

Online First: October 23, 2021
| DOI: 10.15562/bmj.v10i3.2301

[Abstract](#) | [PDF](#)

REVIEW

Management of Spinal Cord Injury During COVID-19 Era

Azharuddin Azharuddin, Diaz Novera

Online First: November 04, 2021
| DOI: 10.15562/bmj.v10i3.2430

[Abstract](#) | [PDF](#)

REVIEW

Assessment and Detection of male Reproductive Outcome (ANDRO) model as a concept for improving the contribution of general practitioners in Indonesia for male infertility management

Dicky Moch Rizal

Online First: November 04, 2021
| DOI: 10.15562/bmj.v10i3.2670

[Abstract](#) | [PDF](#)

CASE REPORT

A rare case of porokeratosis mibelli in 3-year-old boy

Wahyu Lestari, Sitti Hajar

Online First: September 14, 2021
| DOI: 10.15562/bmj.v10i3.2424

[Abstract](#) | [PDF](#)

CASE REPORT

Sclerosing rhabdomyosarcoma of the gingiva mimicking malignant peripheral nerve sheath tumor: a case report

I Wayan Juli Sumadi, Dandy Citra, Elysanti Dwi Martadiani, I Nengah Wiadnyana Steven Christian

Online First: December 07, 2021
| DOI: 10.15562/bmj.v10i3.2679

[Abstract](#) | [PDF](#)

CASE REPORT

Cervical myelopathy due to an endodermal intraspinal cyst in a two-year-old child: a case report

Jainal Arifin, Karya Triko Biakto, Putu Acarya Nugraha Winangun

Online First: December 07, 2021
| DOI: 10.15562/bmj.v10i3.2785

[Abstract](#) | [PDF](#)

CASE REPORT

Intractable singultus: a case report

Anita Hartono, Tomy Lesmana

Online First: December 22, 2021
| DOI: 10.15562/bmj.v10i3.2786

[Abstract](#) | [PDF](#)

CASE REPORT

Functional outcome of a patient with upper type brachial plexus injury treated by Saha and Steindler procedure: a case report

Andi Dhedie Prasatia Syam, M. Ruksal Saleh, Indra Harianto Rante, I Gusti Putu Yoga Kusmawan

Online First: December 27, 2021
| DOI: 10.15562/bmj.v10i3.2667

[Abstract](#) | [PDF](#) |

CASE REPORT

Hemorrhagic transformation in SARS-CoV-2 infected patients: case reports from Indonesia

Retnaningsih Retnaningsih, Dodik Tugaworo, Yovita Andhitara, Rahmi Ardhini, Aditya Kurnianto, Jacob Bunyamin, Fatiha Sri Utami, Ida Ayu Sogata, Hairuzaman Hairuzaman

Online First: December 27, 2021
| DOI: 10.15562/bmj.v10i3.2254

[Abstract](#) | [PDF](#) |

EDITORIAL

The importance of histological appearance in a preclinical test of the plant extracts effectiveness on dermal wound healing

David Tjahyadi, Edy Parwanto

Online First: December 09, 2021 | DOI: 10.15562/bmj.v10i3.2765

[Abstract](#) | [PDF](#) |

The difference in the effectiveness of warm compress and active stretching exercise in reducing dysmenorrhea pain

Ni Wayan Tianing^{1*}, Made Hendra Satria Nugraha²,
Angung Wiwiek Indrayani³, Eka Widyadharna⁴

ABSTRACT

Introduction: Dysmenorrhea (menstrual pain) usually occurs at the beginning of menstruation, can be sharp, blunt, cyclic or persistent until the end of the menstrual phase and can last for several hours to one day.

Objective: This study aims to determine the difference in the effectiveness of warm compresses and active stretching exercises in reducing pain in dysmenorrhoea patients.

Method: The design of this study is an experimental method. Pre-test and post-test control group designs were the designs in this study. This study used 20 subjects, divided into two groups, in which the control group (n = 10) received a warm compress while the treatment group (n = 10) received an active stretching exercise. Interventions were given three times a week for four weeks. The sampling technique is simple random sampling. Pain biomarkers were measured by prostaglandin levels, while the pain was measured through a menstrual distress questionnaire (MDQ).

Results: The data analysis showed no difference in prostaglandin level in the control or treatment groups. In contrast, the MDQ value shows a significant value for both the control and treatment groups, with a mean value that is more significant in the treatment group.

Conclusion: Therefore, it can be concluded that active stretching exercise is more effective at reducing pain than warm compresses in dysmenorrhea.

Keywords: warm compress, active stretching exercise, dysmenorrhea, pain

Cite This Article: Tianing, N.W., Nugraha, M.H.S., Indrayani, A.W., Widyadharna, E. 2021. The difference in the effectiveness of warm compress and active stretching exercise in reducing dysmenorrhea pain. *Bali Medical Journal* 10(3): 1041-1044. DOI: 10.15562/bmj.v10i3.2301

¹Biochemistry Department, Faculty of Medicine, Universitas Udayana

²Physiotherapy Department, Faculty of Medicine, Universitas Udayana

³Pharmacology and Therapy Department, Faculty of Medicine, Universitas Udayana

⁴Neurology Department, Faculty of Medicine, Universitas Udayana

*Corresponding to:

Ni Wayan Tianing; Biochemistry Department, Faculty of Medicine, Udayana University;
w.tianing@gmail.com

Received: 2021-03-03

Accepted: 2021-11-05

Published: 2021-12-30

INTRODUCTION

Adolescence is a transition from childhood to adulthood. During this period, changes occur both in biological and physiological rapid, marked by the maturation of primary sex signs and secondary sex signs. The most notable changes in terms of maturity of the reproductive organs are marked by menstruation/menarche in women.¹ Menstruation is one of the signs in the maturation of reproductive organs, usually occurring in women at the age of 15-17 years.² Dysmenorrhea (pain during menstruation) usually occurs at the beginning of menstruation / first day, can be sharp, blunt, cyclic or persist until the end of the menstrual phase and can last for several hours to one day. Sometimes symptoms can last up to 72 hours. Other common symptoms are nausea, emotional changes, diarrhea, headaches.

The incidence of dysmenorrhea globally is reported to be very large, especially in Indonesia. The incidence rate is 55% and occurs in the age range of 15-17 years.^{3,4} As many as 50-70% with symptoms such as cramps and pain in the pelvic area that can spread to the back to the legs, dizziness, fatigue, increased appetite, nausea, vomiting, and diarrhea. Cramps that occur will result in the release of inflammatory mediators into the systemic circulation, resulting in women experiencing increased uterine contractions, nausea, vomiting, and diarrhea that occurs in nearly 60% with primary dysmenorrhea.^{5,6} Generally, dysmenorrhea interferes with the activities experienced by a woman and often results in activities being undertaken disrupted and sometimes stopped completely. Dysmenorrhea is the most commonly reported reproductive disorder

and impacts functional activities.

Several studies showed that if a woman experiences dysmenorrhea and then does stretching movements focused on the abdominal area, the intensity of the pain can be lessened while increasing strength and elasticity of the pelvic muscles and spine⁶. A non-randomized study conducted by Anugraheni and Wahyuningsih (2013) compared a woman who had dysmenorrhea who actively performed stretching movements compared with women who had never done stretching exercises and had significant results in improving mood and increasing a person's functional activity.⁷

The etiology of primary dysmenorrhea is not yet fully understood but is generally related to the ovulatory cycle. Several factors are associated with the emergence of pain in primary dysmenorrhea, one of

which is a prostaglandin. Prostaglandin is a hormone that comes from the body, synthesized from essential acids. There are two types of prostaglandins: PGE2 and PGF2 α , each binds to 2 aliphatic chains. Prostaglandins and their PGE2 and PGF2 α particles, synthesized from arachidonic acid, are derived from the hydrolysis of phospholipid membrane cells by the enzyme lysosomal phospholipase A2. Arachidonic acid is synthesized via cyclooxygenase or lipoxygenase⁸. PGF2 α is the derivatives of arachidonic acid metabolism by the enzyme cyclooxygenase, which causes vasoconstriction and contraction of the myometrium, causing ischemic pain. PGF2 α and PGE2 have a high ratio in endometrium and menstrual blood in women who experience primary menstrual pain (primary dysmenorrhea). However, PGF2 α and PGE2 have the opposite effect, where PGF2 α has the function to stimulate contractions of the uterus during all phases of the menstrual cycle. Meanwhile, PGE2 can inhibit myometrial contractility during menstruation and stimulate it during the proliferative and luteal phases. Research on the relationship of prostaglandins with the appearance of pain during menstruation has not been done much, so it is very necessary to do the research, which can then be used in the management of pain during menstruation.

MATERIALS AND METHODS

Study design

The design of this study was an experimental study with a randomized pre-test and post-test group control design. Group 1 was given intervention in the form of Active Stretching Exercise, and group 2 as a control group was given intervention in the form of warm water compress.

Place and Time

The study was conducted at the Biochemistry Laboratory of the Faculty of Medicine, Udayana University, for four months, beginning August to September 2017. Therapeutic interventions for each study sample were conducted three times a week for four weeks.

Populations and Samples

The target population for this study was female students with primary dysmenorrhea. The affordable population in this study were female students in semester I and semester III Physiotherapy Study Program, Faculty of Medicine, Udayana University, who had primary dysmenorrhea. Based on sample calculations, the minimum sample needed in this study was 9 participants. The number of study participants was added by 10% to become 10 participants in each treatment group to anticipate the subject drop-out.

Research procedure

The two groups were both measured for pain using the Menstrual Distress Questionnaire in the pre-test and post-test. The subject gives answers to each parameter. Before measuring the pain level, subjects were given an explanation of the Menstrual Distress questionnaire. Subjects were asked to answer several questions with criterion values: never, rarely, sometimes, often, always. Then, when the pain appeared, the blood was taken from each respondent to measure prostaglandin levels.

Procedure for Warm Water Compress Implementation

Warm compresses are applied to the front and side of the abdomen and back. This intervention was carried out 3 times

a week for 4 weeks with a duration of 15 minutes per session.

Procedure for Implementing Active Stretching Exercise

Active stretching exercise is applied through the movement of stretch paint, lower trunk rotation, buttock hip stretch, abdominal strengthening (curl up), and bridging. Each movement is held for 20 seconds and repeated three times. This intervention was carried out three times a week for four weeks.

RESULTS

Subject Baseline Characteristics

Table 1 above describes the sample characteristics based on sex and age. The total number of samples in the control group and the treatment group is 20 peoples

Prostaglandin Decrease Test Mean Difference Before and After Intervention in the Control and Treatment Groups. The test results are listed in **Table 3**.

Comparison of Prostaglandin Level between Groups

Based on **Table 2**, the results of a decrease in prostaglandin values were analyzed by Wilcoxon sign rank test before and after intervention in the control group with a value of $p = 0.241$ or $p > 0.05$, which means that there was no significant difference in the decrease in prostaglandin level before and after the intervention.

Table 1. Gender distribution of the research subjects.

CHARACTERISTICS	CONTROL GROUP (n=10) f (%)	TREATMENT GROUP (n=10) f (%)
GENDER (%)		
WOMEN	10(100)	10(100)
AGE (year)		
Mean \pm SD	19.60 \pm 0.516	19.50 \pm 0.527

Table 2. The Results of Prostaglandin Value Analysis with the Wilcoxon Sign Rank Test Data

Data of Groups	P value
Control groups	0.241
Treatment groups	0.093

Evaluating the Effect of Warm Compress and Active Stretching between The Research Groups

Based on Table 3 shows a decrease in MDQ values analyzed by paired sample t-test before and after the intervention in the control group with a value of $p = 0.002$ or $p < 0.05$, which means that there was a significant difference in the decrease in MDQ values before and after giving warm water compresses. Likewise, in the treatment group, $p = 0.000$ or $p < 0.05$ was obtained, which means a significant difference in the decrease in MDQ values before and after administration of active stretching exercise.

Post-Test Difference Test For MDQ Impairment Before And After Intervention Between Control And Treatment Groups

The Comparison Of MDQ Post-Test Impairment Between Before And After The Intervention In The Two Groups That Were Given Warm Water Compress Intervention In The Control Group And The Active Stretching Exercise Intervention In The Treatment Group Was Tested With Independent Sample T-Test. The Test Results Are Listed In The Following Table 4.

Table 4 shows the calculations regarding the difference in mean impairment of MDQ values, i.e., $p = 0.004$ or $p < 0.05$, between the two intervention groups. This indicates that there are significant differences between the two pieces of training.

DISCUSSION

The warm water compress decreased MDQ values were analyzed by paired sample t-test before and after intervention in the control group with $p = 0.002$ or $p < 0.05$, which means a significant difference in the decrease in MDQ values before and after giving warm water compresses. The results of this study were supported by Anugraheni and Wahyuningsih and Dahlan and Syahminan, who stated that giving warm water compresses can reduce pain in dysmenorrhoea.^{7,9} In addition, Lowdermilk *et al.* explain that dysmenorrhoea pain can be reduced by non-pharmacological therapy in the form of warm compresses because it provides a sense of security for patients by using fluids or devices that cause warmth in

Table 3. MDQ Value Analysis Results with Paired Sample t-test.

Data Group	Before intervention	after intervention	P-value
	Mean±SD	Mean±SD	
Control groups	86.7±3.23	79.00±3.40	0.002
Treatment groups	87.40±4.55	72.00±6.83	0.000

Table 4. Comparison between MDQ value between Groups in Pre-Intervention and Post-Intervention

	Data groups	Mean±SD	P-value
Pre-Test	Control groups	86.70±3.23	0.696
	Treatment groups	87.40±4.55	
Post-Test	Control groups	79.00±3.40	0.010
	Treatment groups	72.00±6.83	
Difference	Control groups	7.70±5.67	0.004
	Treatment groups	15.40±4.86	

parts of the body that need it.¹⁰ The results in the heat transfer to the stomach so that the compressed stomach becomes warm, there is a dilation of blood vessels in the area that is experiencing pain and increased blood flow in the area so that the perceived dysmenorrhoea pain will decrease or disappear. According to Lowdermilk *et al.*, non-pharmacological arm compresses are very beneficial in reducing dysmenorrhoea pain.¹⁰ Muscle relaxation occurs and reduces uterine ischemia so that pain can be reduced. Active stretching exercise also decreased MDQ values. It was analyzed by paired sample t-test before and after intervention in the control group with a value of $p = 0,000$ or $p < 0.05$, which means significant differences in the decrease in MDQ values before and after administration of active stretching exercise.

The results of this study support the theory conveyed by Maidartati *et al.*, which states that active stretching exercise can increase muscle strength while increasing joint mobility and flexibility of the muscle; when muscle groups are given stretching, then stretch reflex works automatically to contract by stretching the muscles to protect it from excessive stretching, the

reflex that occurs in the Golgi tendon will be activated and inhibit tension by relaxation through muscle lengthening when there is increased tension (tension) in the muscle group.¹¹

According to Bobak, women with dysmenorrhoea who have warm compresses when experiencing dysmenorrhoea respond by vasodilation from target specific tissues such as muscles, which is the mechanism performed when giving warm compresses to the body¹². Where the body will respond by sending signals to the hypothalamus, the hypothalamus has an important role in controlling bodily functions, including the release of hormones from the pituitary gland and homeostasis function of the body, when the hypothalamus responds to signals from the internal environment in the form of temperature changes the effector gives signals to the body in the form of dilation of blood vessels (vasodilation) which is regulated by the vasomotor system that is in the medulla oblongata under the control of the hypothalamus so that vasodilatation of the tissue occurs, the mechanism will affect the connective tissue such as muscles to relax in certain muscle groups. However, the effect produced on the warm compress

is pseudo only about 10-20 minutes after using the warm compress. The target-specific tissue will experience increased back pain is a drawback of a therapy for some cases related to connective tissue such as muscles, ligaments, and tendons.

From the results of this study it can be concluded that, the two interventions have different results in which active stretching exercise has more effective and significant results in reducing dysmenorrhea pain by working targets to reduce muscle tension caused by the release of substance P produced by the brain with a mechanism muscle lengthening of the golgi tendon. Decreased muscle mobility due to reduced physical activity causes changes in the muscle structure. It can change the pain receptors produced by a particular muscle group so that it causes motion limitations due to increased muscle stiffness. Mobility of a specific muscle group can be improved through various therapies and rehabilitation, such as active stretching, passive stretching or some other exercise therapy. Based on the results of studies on active stretching exercise and warm compresses have the advantage of reducing the level of pain produced by muscle tension.^{12,13}

However, active stretching exercise has the advantage of actively influencing changes in muscle length and tendons which will overall affect changes in anatomy, physiology and biomechanics which will result in biomechanical function in joints and tissue metabolism and active mobilization will affect the movement of primary muscles in the body involved in the mobility of a muscle particle.¹⁴ Meanwhile, warm compresses do not cover the results and the mechanism produced by the active stretching exercise. Still, rather short-lived results are more generated by the mechanism of the warm compresses.

Based on the results of previous studies and studies, it can be concluded that the reduction in pain levels can be achieved significantly by both types of therapy

through the components that have been described. However, active stretching exercise is more effective in reducing pain compared to warm compresses because several important components are involved and successfully obtained during the implementation of vigorous stretching exercise, which is very useful and applied in cases of primary dysmenorrhea.

CONCLUSION

Based on the results of this study, it can be concluded both compresses warm water or provision of active stretching exercise is effective in reducing pain in dysmenorrhoea. Active stretching exercise is more effective in reducing pain than warm water compresses in dysmenorrhoea. For the future study, we recommended assessing other therapy to compare with active stretching exercise, which can be used to reduce pain in dysmenorrhea

CONFLICT OF INTEREST

All authors declared that there is no conflict of interest regarding this article.

AUTHOR CONTRIBUTION

All authors contributed equally to the research process and the writing of this article.

FUNDING

This study was self-funded

ETHICS APPROVAL

This research was ethically approved with ethical clearance number 1734/UN14.2.2.VII.14/LT/2020.

REFERENCES

- Narayan KA, Srinivasa DK, Peltó PJ, and Veeramal S. Puberty Rituals, Reproductive Knowledge and Health of Adolescent Schoolgirls in South India. *Asia-Pacific Population Journal*. 2001;16(2):225-238

- Mashvash N, Eidy, A, Mehdi, K, Toorzani, Z, Mani, M, Shahla, H. The Effect of Physical Activity on Primary Dysmenorrhea of Female University Students. *World Applied Sciences Journal*, 2012;17(10): 1246-1252
- Pedron- Nuevo N, Gonzalez-Unzaga LN, De Celis-Carillo R, Reynoso Isla M, De La Torre-Romeral L. Incidence of Dysmenorrhea and Associated Symptoms in Women Aged 12-24 Years. *Ginecol Obstetric Mex*. 1998; 66:492-494
- Dawood, M.Y. Primary dysmenorrhea: Advances in Pathogenesis and Management. *Obstetrics & Gynecology* 2006.; 108 (2):428-441
- Cakir M, Mungan I, Karakas T, Giriskan I, Okten A. Menstrual Pattern and Common Menstrual Disorders Among University Students in Turkey. *Pediatr Int*. 2007;49(6):938-942
- Patel, V. The Burden and the Determinants of Dysmenorrhea: a Population- Based on Survey of 2262 Woman in Goa, India. *International Journal Obstetrics and Gynecology*. 2006;113(4) 453-463
- Anugraheni, VMD and Wahyuningsih, A. The Effectiveness of Warm Compresses in Reducing the Intensity of Dysmenorrhoea Pain in Students. *Baptist Kediri Hospital.STIKES Journal*, 2013; 6(1)
- El-Gilany, A-H., K. Badawi and El-Fedawy. S. Epidemiology of Dysmenorrhea among Adolescents Students in Mansoura, Egypt. *Eastern Mediterranean Health Journal*, 2005; 11(1/2): 155-163
- Dahlan, A and Syahminan, TV. The Effect of Warm Compress Therapy on Menstrual Pain (Dysmenorrhoea) in Simpang Haru Vocational School Students in Padang. *Applied Science and Technology Journal*, 2016; 10 (2):141-147
- Lowdermilk D, Perry S, and Cashion MC, and Alden KR. *Nursing Maternitas*. Elsevier Mosby. 2013.
- Maidartati, M, Hayati, S, Hasanah, AP. The Effectiveness of Warm Compress Therapy Against Decreasing The Pain of Dysmenorrhea in Adolescents in Bandung. *Nursing BSI Journal*, 2018; 6 (2),156-164
- Bobak L. *Maternity nursing textbooks*. Jakarta:EGC. 2005.
- Maimunah, S, Sari, RDP, Prabowo, AY. Comparison of The Effectiveness of Warm Compresses and Cold Compresses as A Non Pharmacological Treatment of Dysmenorrhea in Adolescents. *Medula Jurnal*, 2017; 7(5):79-83
- French, L., *Dysmenorrhea American family physician*, 2005; 71(2):285-291



This work is licensed under a Creative Commons Attribution